covered with granulation tissue, and the foot and leg fixed in plaster of Paris. The healing of both wounds was quickly accomplished and the plaster-of-Paris splint dispensed with about three months after the operation. Some time afterwards, as there was still a tendency to equinus, the tendo Achillis was again divided and the foot put into plaster-of-Paris for a fortnight. The result now—ten months after the operation—is very satisfactory, the foot being in good position, with more strength in it than one dared hope for after the division of the tendons of so many important muscles.

Great Ormond-street, W.C.

CASE OF FRACTURE OF THE BASE OF THE SKULL, WITH OUTWARD DISPLACEMENT OF BONE.

By C. B. DALE, M.R.C.S. ENG., L.R.C.P. LOND.,
LATE HOUSE SURGEON, BUCKINGHAMSHIRE GENERAL
INFIRMARY, AYLESBURY.

ELEVATED fractures of the vertex of the skull are described as the result of oblique cuts. In the following case a similar condition affected the base. On Sept. 26th, 1894, a man aged fifty-one years was working on the Great Northern Railway line when an iron plate measuring 13 in. $\times$ 7½ in. $\times$ ½ in. flew off the engine of a passing express and struck him behind the left ear. When seen a few minutes afterwards he was in a semiconscious condition, but rapidly became comatose. There was an irregular, lacerated wound about 5 in. in length, extending downwards and forwards behind the left ear as far as the angle of the jaw. The skull was perforated just behind the mastoid process, and a circular piece of bone about 2 in. in diameter was cut out and turned back in the posterior flap. There were some loose splinters of bone lying in the wound, but the edges of the opening in the skull were not depressed, and no fissures could be detected extending from it. The surface of the brain was lacerated. The wound was washed out with carbolic acid lotion and brought together with carbolised silk sutures, the piece of bone being replaced and kept in position by the flap to which it was attached. The patient made a good recovery. The wound healed by first intention, with the exception of the lower angle, from which a small quantity of cerebro-spinal fluid continued to escape for some days. Four days after the accident he developed left-sided facial paralysis, but this has since almost completely disappeared. He is now back at work and complains only of slight weakness of the right leg.

Hatfield.

A Mirror OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proœmium.

KING'S COLLEGE HOSPITAL.

A CASE OF FLOATING KIDNEY WITH INTERMITTENT HYDRO- NEPHROSIS CURED BY FIXING THE KIDNEY IN THE LOIN.

(Under the care of Mr. WATSON CHEYNE.)

THIS case illustrates a condition of the kidney which is much more common than was supposed a few years ago; in fact, the relationship between movable kidney and hydro-nephrosis was not mentioned in our text-books and was perhaps not recognised. Probably the most interesting and important contribution to this subject (intermittent hydro-nephrosis) is that by Terrier and Baudouin,¹ who collected a series of eighty-three cases. They found that the most

common cause of this condition is a movable kidney, which when displaced causes a kink in the ureter, the escape of urine being thus prevented until the kidney has returned to its normal position. They state that such a condition usually becomes a permanent one, because in consequence of inflammatory changes about the kidney adhesions form which hold it in its displaced position. The symptoms produced by intermittent hydro-nephrosis were well shown in the case which follows, whilst the treatment carried out has proved very successful. It is, of course, advisable that the operation should be performed as early as possible, so that the inflammatory changes which are only too apt to ensue and produce destruction of kidney substance, and may possibly necessitate the operation of nephrectomy, may be prevented.

A widow aged thirty-five years was admitted to King's College Hospital on Nov. 16th, 1893. Four years before admission she first began to suffer from attacks of pain in the left side. These attacks lasted from two or three days to a fortnight, and from that time she had had attacks of this kind every week or ten days. At first she noticed that the attacks occurred especially after exertion, such as lifting heavy weights, and relief was sometimes speedily obtained if she lay down on her side. Latterly, however, the attacks had occurred without exertion, and even on turning in bed. The pain was of an aching character, beginning mildly at first, gradually increasing in severity, and subsiding either gradually or sometimes pretty quickly. For the last year, coincidentally with each attack of pain she had noticed the appearance of a lump in front and above the left anterior iliac spine, which was very tender and was the seat of the pain. It appeared and disappeared coincidentally with the pain. Tenderness remained for some days after the attack. During these attacks she did not notice any frequency of micturition or alteration in the quantity of urine passed, nor had she noticed any increase in the quantity of urine when the tumour disappeared. She had, however, frequently noticed blood in her urine, both during and in the intervals between the attacks. She was admitted to hospital during an attack, but on her being examined two days later the pain and tumour were found to have completely disappeared. An elongated mass which seemed to be the remains of the kidney was, however, felt and was freely movable. The patient's general health was good. The urine was acid, of sp. gr. 1030, and with a considerable deposit of urates, but no albumen, sugar, or blood. As no lump was present she went to the ophthalmic wards on account of a keratitis from which she had suffered for some months. She afterwards went out for a few days and returned on Dec. 20th with a fresh attack of pain in the lumbar region. On examination a large, soft, rounded swelling about the size of two fists was felt in the left lumbar region, extending as low as the anterior iliac spine and passing up under the ribs. There was also bulging in the left lumbar region behind. There was dulness over this area, and the swelling was very tender to the touch. On considering the history and symptoms Mr. Watson Cheyne came to the conclusion that this was one of the cases which have been recently described by Mr. Clement Lucas, where hydro-nephrosis had followed on a movable kidney, and that the attacks of pain &c. coincided with a kinking of the ureter from an alteration in the position of the kidney. As it was evident that considerable kidney substance still remained, Mr. Watson Cheyne decided to wait till the attack passed off—that is to say, till the position of the kidney which led to kinking of the ureter had given place to a position in which the ureter was free—and then to try to fix the kidney in that position. The swelling and other symptoms had completely disappeared by Jan. 12th, 1894, and accordingly on Jan. 17th the usual oblique posterior incision was made and the kidney exposed. Considerable remains of kidney substance were found, of a horse-shoe shape, and in front and below a large empty sac, the dilated pelvis. The kidney having been placed as nearly as possible in its normal position, a portion of the capsule was stripped off and stitched in with stitches which united the divided layers of muscles somewhat after the manner described by Mr. Arbuthnot Lane. There is nothing special to note about the after-progress of the case except that two of the superficial stitches found their way out through an angle of the wound left open for drainage. The patient left the hospital on Feb. 24th, not having had any further accumulation or attack of pain. She was seen quite lately, when she stated that she had remained perfectly well and free

¹ *Revue de Chirurgie*, Paris, 1891, vol. ii. *Sajous' Annual of the Universal Medical Sciences*, vol. iii., E. 23.