Operation.—Sept. 2, 1911, the patient was given ether and a tourniquet applied at the wrist. A curved linear incision was made in the palm of the hand and the small tumor was found at the center of the deep palmar arch. On loosening up the tourniquet the tumor could be seen to pulsate plainly, and to increase to more than twice its size in the collapsed state. It was an aneurysm with a small pedicle, which was easily ligated, and the whole sac removed.

The patient made an uneventful recovery, and now has no difficulty at all in using this hand.

A NEW TONSIL HOOK

L. F. LONG, M.D.
ZANESVILLE, O.

I have devised a new tonsil hook which seems to me an improvement over any that I have used. Its advantages are the following:

A snare can be slipped on or off the tonsil without changing the hold on the tonsil after the hook is once placed.

The hook, once placed, will stay in place and not drop out when pressure is relaxed, as the slight thickening on the lower two-thirds of the inner surface of the prongs holds it in place.

601 Market Street.

KNEE-JOINT AMPUTATIONS

REPORT OF A NEW METHOD

CHARLES EATON PHILLIPS, M.D.
ANCON, CANAL ZONE

To overcome the disadvantages of the thigh amputation, and to combine the good points and disadvantages of disarticulation with those of the osteoplastic operation, I have devised the following operation:

An oval anterior and posterior flap are turned up, the former line of incision extending from the lower portion of the internal condyle downward, and anterior as far as the tuberosity of the tibia. The patella tendon is cut without dissecting the skin from the patella and also leaving a portion of the patella tendon attached to the patella. The posterior flap is cut off approximately equal length or a little longer, if possible. The hamstring muscles are cut long. Disarticulation is then completed by cutting the remaining structures. Vessels are ligated, nerves pulled down and cut short, and allowed to retract.

The anterior flap is next turned up and the posterior surface of the patella sawed off. This is easily done by grasping the patella tendon by a pair of strong holding forceps when the patella can be sawed without difficulty, care being taken to remove as small a part of the patella as possible, so as to preserve nearly its original thickness.

Next, a small piece is sawed off each side (Fig. 1 A A) in such a manner as will leave the patella narrower toward the upper end, or quadriceps attachment, and also narrowed toward the anterior surface.

If it has been possible to preserve sufficient skin flaps to cover the stump, the next step is to saw into each side of the intercondyloid notch (Fig. 2 A A) in such

a manner that, when the piece of bone has been removed by means of a mallet and chisel (Fig. 2 B), there is left a depression, or modified intercondyloid notch, which is broader at the bottom and also toward the popliteal space, or, in other words, narrower at the end of the bone and also toward the anterior surface. The bevel of the notch is made to conform as nearly as possible to the bevel of the patella, but as there is of necessity a slight error, our aim should be to make the bevel of the patella greater, or that of the notch less, so as to insure good apposition when the patella is fitted into position.

The patella is next grasped by a pair of strong bone-holding forceps and drawn down forcibly until it slips into the place prepared for it and dovetails into it (Fig. 3). The greater the muscular pull, the more firmly it holds its place.

Next, the hamstring muscles are sutured to the patella tendon. The skin is sutured, a small drain inserted and the leg put up in dressing. It will be found that the forcible drawing down of the patella brings the line of skin suture well to the popliteal space and away from all pressure.

If, however, it is impossible or inexpedient to save enough skin to cover the stump without too great tension—and this can be determined by grasping and drawing

Figs. 1, 2 and 3.—Diagrams of patella and joint, illustrating new method of knee-joint amputation.

the patella down before anything has been done toward preparing the notch—equally good results may be obtained by sawing through the condyles and then cutting out a notch in the sawn condyles so that the patella will dovetail into it as before. Nearly 2 inches may be gained in this way without sacrificing any of the advantages of the operation.

In either case we get a stump that is very satisfactory. The patella forms the center of it, and laterally the two condyles share their portion of the weight in such a manner that the normal distribution of weight, as in the flexed knee, is preserved.

The large bearing surface formed by the tough skin taken from the anterior aspect of the knee is able to take the full weight of the body without abrasion or discomfort.

Another advantage, which it shares in common with disarticulation, or amputations below the knee, is the enlargement furnished by the condyles, or even portions of the condyles. With the enlargement, an artificial leg can be adjusted so that it will not slip up and down when walking or sitting.

Still another advantage which this operation possesses is the short flap which is sufficient to cover the stump

this gain is made by forcibly drawing down the patella, bringing with it the skin fascia and muscles, which, if allowed to retract, would fail to cover a stump several inches shorter.

A representative of one of the largest artificial limb manufacturing in the country remarked, after careful examination of one of these amputations, that the stump was ideal for the fitting and wearing of an artificial leg, and that it was by far the best end-bearing stump he had ever seen, and urged its adoption in suitable cases, from a prosthetic standpoint.

The operation is not at all difficult and requires very little additional time. The first time I ever attempted the operation it took about a half hour with closure and dressing.

The limits of usefulness of this operation, as with any, are narrow. No one would sacrifice $1\frac{1}{2}$ or 2 inches of tibia needlessly, but where it is impossible to save the knee, by this amputation all the benefits of a knee-bearing stump may be preserved.

Ancon Hospital.

CASE OF INTUSSUSCEPTION COMPLICATED BY VOLVULUS

C. S. LAWRENCE, M.D.
WINSTON-SALEM, N. C.

A careful search of the literature at my command has failed to reveal a case of intestinal obstruction in which the symptoms and pathologic findings coincide with the following case:

Aug. 21, 1911, I first saw Mr. B., white, aged 22, laborer. Family history was negative. Patient had had usual diseases of childhood. He had had attacks of severe pains in abdomen with nausea and vomiting at intervals of from one to six months for the past six years. Four years ago his appendix was removed, but this gave him no relief, and one of his attacks came on a few days after the operation. For the past six months these attacks have increased in frequency and severity.

Present Illness.—About 9 a. m. Aug. 20, 1911, while at breakfast, he was taken with one of his usual attacks of abdominal pain, he did not eat much breakfast, thinking the pain would pass off, as usual, but it grew worse. He became nauseated and vomited. A physician was called and gave morphin to relieve the pain, but it failed; chloroform was given, but as soon as the effect of the chloroform would wear off, the pain would come on again with great severity. An enema of soap and water was given with no result. The pain, vomiting and constipation continued all that day and the next until 10 p. m., when I saw him in consultation with Dr. V. M. Long. His condition was as follows: He was lying in bed, on his right side, thighs flexed on abdomen, anxious facial expression, pulse 110, temperature 101 F.; abdomen rigid, slightly distended in upper part. On palpation a mass could be felt below and slightly to the left of the umbilicus. On rectal examination the mass could be felt at the brim of the pelvis, but it was not in the rectum. The patient had had large quantities of castor oil during the day, and an enema in the evening, which was returned with a large quantity of blood, but no fecal matter or flatus. The patient had also vomited a dark fluid, which the father said smelled bad. He had voided a small amount of urine before I saw him.

Treatment.—Operation was decided on at once. Under ether anesthesia a median incision about three inches long, between the umbilicus and symphysis, was made. On opening the peritoneal cavity a large quantity of seropurulent fluid escaped. On passing the finger down to the brim of the pelvis a mass could be felt attached by what seemed to be a hard cord, the mass hanging over the pelvic brim. There were a few soft adhesions which were separated with the finger, and

the mass delivered, together with several coils of distended ileum. On examination, the mass was found to be an iliac intussusception which had become twisted on its mesenteric axis. The involved intestine was very black, so that it was considered unsafe to leave it. The entire mass was resected and an end-to-end anastomosis made with the Murphy button, reinforced by silk Lembert sutures. Gauze drains were placed in the upper and lower ends of the incision. The patient left the table in about thirty minutes; pulse 160.

Postoperative History.—He received strychnin, gr. 1/20, and 700 c.c. normal salt solution under the skin while on the table. After being placed in bed morphin, gr. $\frac{1}{4}$, and atropin, gr. 1/150, was administered. One hour later the pulse was 140, respiration 34. The pulse gradually came down to 110 at 6 a. m., temperature 99 F. Both began to rise; morphin, gr. $\frac{1}{4}$, atropin, gr. 1/150, was given and proctolysis started. Patient passed gas. The stomach was distended and patient vomited a dark fluid. The stomach-tube was passed and a large quantity of gas and dark fluid mixed with oil was removed. The patient continued to improve from this time on, the stomach being washed at intervals of from six to eight hours for the first forty-eight hours. The bowels moved freely thirty-six hours after the operation. The patient voided urine freely and never had to be catheterized. Everything went smoothly until the fifth day when I dressed the wound and removed the drain from the upper part of the incision and placed in a fresh piece of gauze in a very small opening. In the afternoon the patient complained of pain in the abdomen and his temperature was found to be 103.2, pulse 120. On removal of the gauze a small quantity of purulent material was found. A small rubber tube was inserted and the temperature promptly returned to normal. When the dressings were removed from the abdominal wound sixteen days after operation they were completely saturated with a dark fluid, due to a fecal fistula discharging from the upper drainage tract. The small tube was removed; the fistula discharged profusely all day, but soon closed and by the third day there was not enough discharge to soil the dressing through more than four layers of gauze. On September 18 patient went home and was up and about the house; September 26 he passed the button. At present, September 28, he is about, looks well and feels well.

208 Masonic Temple.

New and Nonofficial Remedies

THE FOLLOWING ADDITIONAL ARTICLES HAVE BEEN ACCEPTED BY THE COUNCIL ON PHARMACY AND CHEMISTRY OF THE AMERICAN MEDICAL ASSOCIATION. THEIR ACCEPTANCE HAS BEEN BASED LARGELY ON EVIDENCE SUPPLIED BY THE MANUFACTURER OR HIS AGENT AND IN PART ON INVESTIGATION MADE BY OR UNDER THE DIRECTION OF THE COUNCIL. CRITICISMS AND CORRECTIONS ARE ASKED FOR TO AID IN THE REVISION OF THE MATTER BEFORE PUBLICATION IN THE BOOK "NEW AND NONOFFICIAL REMEDIES."

THE COUNCIL DESIRES PHYSICIANS TO UNDERSTAND THAT THE ACCEPTANCE OF AN ARTICLE DOES NOT NECESSARILY MEAN A RECOMMENDATION, BUT THAT, SO FAR AS KNOWN, IT COMPLIES WITH THE RULES ADOPTED BY THE COUNCIL.

W. A. PUCKNER, SECRETARY.

PROPAESIN—Propylis Aminobenzoas.—Propyl Aminobenzoate.—Paramidobenzoic Acid Propyl Ester.—Propaesin, $C_6H_4.NH_2.COO(C_2H_5)_3$:4, is the propyl ester of paraminobenzoic acid, $C_6H_4.NH_2.COOH$:1:4.

Propaesin is prepared by esterification of paraminobenzoic acid with propyl alcohol.

Propaesin is a fine, white or colorless, odorless, nearly tasteless powder, which produces numbness when placed on the tongue. Propaesin is very slightly soluble in water and is not readily wetted by this solvent. It is soluble in alcohol, benzene, chloroform and ether. Propaesin melts at 73° C. (163.4° F.).

When heated on platinum foil Propaesin burns without leaving any ash; when heated with an excess of potassium hydroxide test solution it melts before boiling, forming an oily layer. By boiling this mixture the melted Propaesin is saponified with the formation of propyl alcohol, which may be recognized by its odor. A clear,