


This article was downloaded by: [University of Alberta]
On: 02 January 2015, At: 05:14
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer Street,
London W1T 3JH, UK



Annals and Magazine of Natural History: Series 8

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah14>

LII.—A synopsis of the fishes of the subfamily Salanginæ

C. Tate Regan M.A.

Published online: 08 Sep 2009.

To cite this article: C. Tate Regan M.A. (1908) LII.—A synopsis of the fishes of the subfamily Salanginæ, *Annals and Magazine of Natural History: Series 8*, 2:11, 444-446, DOI: [10.1080/00222930808692507](https://doi.org/10.1080/00222930808692507)

To link to this article: <http://dx.doi.org/10.1080/00222930808692507>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly

or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

beneath black; legs ochraceous; head shorter than pronotum, ocelli a little less removed from each other than from eyes; tegmina distinctly thickly punctate; posterior tibiae with two strong spines.

Long. 6 mm.

E. Ruwenzori, 6000–13,000 feet.

LII.—A *Synopsis of the Fishes of the Subfamily Salanginæ.*

By C. TATE REGAN, M.A.

SALANGINÆ.—Argentinidæ with the dorsal fin placed far behind the pelvics, the head depressed, flat above, and the body elongate, subcylindrical anteriorly and compressed posteriorly. Teeth conical; scales deciduous; 4 branchiostegals; pelvic fins 7-rayed*.

Synopsis of the Genera.

I. Præmaxillaries subnormal; lower jaw projecting.

A. Dorsal fin entirely in advance of the anal; tongue with 2 series of teeth; 2 series of teeth on each side of the palate; teeth in the jaws small, subequal 1. *Protosalanx*, gen. nov.

B. Dorsal fin partly above the anal; tongue toothless; 1 series of teeth on each side of the palate.

Teeth in the jaws small, subequal 2. *Salangichthys*, Bleek.

A pair of canines near the symphysis of the lower jaw; præmaxillary teeth somewhat enlarged, strongly recurved 3. *Hemisananx*, gen. nov.

II. Præmaxillaries forming an anterior triangular expansion; lower jaw not projecting; a single series of teeth on each side of the palate; 1 or more anterior canines in the lower jaw, perforating the roof of the mouth behind the præmaxillary expansion; præmaxillary teeth strong, recurved, set rather far apart.

A. Dorsal fin entirely in advance of the anal; tongue with a single series of teeth; lower jaw ending in a short fleshy appendage. 4. *Leucosoma*, Gray.

B. Dorsal fin wholly or partly above the anal; tongue toothless.

Lower jaw ending in a short fleshy appendage 5. *Salanx*, Cuv.

Lower jaw ending in a distinct, more or less movable præsymphyseal bone, with a double series of teeth 6. *Parasalanx*, gen. nov.

* In *Protosalanx*, *Salangichthys*, *Hemisananx*, and *Parasalanx*, and probably throughout the group, the males have the outer pectoral ray produced and the anal fin modified, the middle rays being set close together and curved backwards, whilst on each side of the body a single longitudinal series of adherent scales, decreasing in size posteriorly, is placed above the base of the fin.

*Synopsis of the Species.*1. *Protosalanx hyalocranius*, Abbott, 1901*.

D. 17. A. 30-32. P. 23-25. Depth of body 10-13 in the length (♀), length of head $5\frac{1}{4}$ - $5\frac{1}{2}$.

Three specimens, 115-130 mm. in total length, from Shanghai (*Swinhoe*).

2. *Salangichthys microdon*, Bleek., 1860.

D. 12-14. A. 25-27. P. 15-16. Origin of anal below posterior rays of dorsal.

Numerous specimens, up to 85 mm. in total length, including the types of the species from Yeddo (*Bleeker*), and examples from Vladivostock (*Hook*), Yokohama ('*Challenger*'), and the Inland Sea of Japan (*Gordon Smith*).

3. *Hemisalanax prognathus*, sp. n.†

D. 13. A. 26. P. 10. Depth of body 14 in the length, length of head $6\frac{1}{3}$.

A single specimen, 120 mm. in total length, from Shanghai (*Swinhoe*).

4. *Leucosoma chinense*, Osbeck, 1765 (*Synodus macrocephalus*, Lacep., *Leucosoma reevesii*, Gray).

D. 10-11. A. 29-31. P. 10-12.

Four specimens, 160-180 mm. in total length, from China, including the type of *L. reevesii*.

5. *Salanax cuvieri*, Val., 1849.

D. 13. A. 26-27. P. 9-10. Depth of body 13 to 14 in the length, length of head $5\frac{1}{4}$. Head 3 times as long as broad; snout as long as postorbital part of head. Origin of pelvics nearer to that of anal than to base of pectoral; anal origin below fourth or fifth ray of dorsal.

* Abbott gives D. 16-18. A. 28-31. P. 27 for specimens from Tientsin. His figure of a male shows depth of body 9 in the length.

† *Salanax ariakensis* (Kishinouye), Jord. & Snyder. Proc. U.S. Nat. Mus. 1902, p. 592, from the Ariaka Sea (Kiushiu), is an insufficiently described species which may belong to *Hemisalanax*.

D. 13. A. 26. P. 10. Dorsal opposite anal. Teeth subequal; tongue toothless.

Two specimens, 150 mm. in total length, from Ichang (*Pratt*).

I am indebted to Dr. Pellegrin for notes on the type of the species, which leave little doubt that these examples should be referred to *Salanæ cuvieri*.

6. *Parasalanæ gracillimus*, sp. n.

D. 12. A. 27. P. 10. Depth of body 18 in the length, length of head $5\frac{1}{2}$. Head a little more than 3 times as long as broad; snout as long as postorbital part of head. Origin of pelvics equidistant from head and origin of anal, which is below the eighth or ninth dorsal ray.

A single specimen, 120 mm. in total length, from Shanghai (*Shanghai Museum*).

7. *Parasalanæ longianalis*, sp. n.

D. 12-13. A. 30-32. P. 10. Depth of body 13 to 15 in the length, length of head $5\frac{1}{2}$. Head 3 times as long as broad; snout shorter than postorbital part of head. Origin of pelvics equidistant from head and origin of anal, which is below the fifth to the seventh dorsal ray.

Four specimens, 110-125 mm. in total length, from Liao-ho, Northern China (*Morrison*).

8. *Parasalanæ acuticeps*, Regan, 1908.

D. 13-14. A. 26-27. P. 9-10. Depth of body 11 in the length, length of head $5\frac{1}{2}$ to $5\frac{2}{3}$. Head 3 times as long as broad; snout shorter than postorbital part of head. Origin of pelvics nearer to anal than to base of pectoral; origin of anal below second dorsal ray.

Two specimens, 115 mm. in total length, from Lake Candidius, Formosa (*Moltrecht*).

9. *Parasalanæ angusticeps*, sp. n.

D. 14. A. 28. P. 10. Depth of body 16 in the length, length of head $5\frac{1}{2}$. Head nearly 4 times as long as broad; snout a little longer than postorbital part of head. Origin of pelvics equidistant from base of pectoral and origin of anal, which is below that of the dorsal.

A single specimen, 153 mm. in total length, from China (*Haslar*).