NOTES FROM THE U. S. BUREAU OF CHEMISTRY.*

THE VOLATILE REDUCING SUBSTANCE IN CIDER VINEGAR.¹

By R. W. Balcom.

[ABSTRACT.]

A HISTORICAL discussion is followed by experimental results which include details of the preparation and purification of the phenylosazone obtained from the distillate from cider vinegar. Such distillates contain a reducing substance which reduces Fehling's solution at room temperature. The melting-point of the phenylosazone and the amount of nitrogen it contained indicated that it was diacetyl phenylosazone. Diacetyl ($\text{CH}_3\text{-CO-CO-CH}_3$) and acetylmethylcarbinol ($\text{CH}_3\text{-CHOH-CO-CH}_3$), the two substances from which this osazone might be formed, were prepared and the behavior of dilute solutions of these two substances was compared with that of the cider vinegar distillate. It is shown that the reducing substance in the cider vinegar distillate is largely, if not wholly, acetylmethylcarbinol. This substance is not formed during the distillation of the vinegar. It is present as such in the vinegar and appears to be a normal constituent of cider vinegar.

SOME PROTEINS FROM THE JACK BEAN (*CANAVALLIA ENSIFORMIS*).*

By D. Breese Jones and Carl O. Johns.

From the Protein Investigation Laboratory, Bureau of Chemistry,
Department of Agriculture, Washington, D. C.

[ABSTRACT.]

Two globulins, *canavalin* and *concanavalin*, and an albumin have been isolated from the jack bean. The air-dried jack bean meal contained 23 per cent. of protein, 15 per cent. of which is extracted from the meal by distilled water. A two per cent. sodium chloride solution extracted 18.5 per cent., and a 0.2 per cent. potassium hydroxide solution extracted nearly all of the

* Communicated by the Chief of the Bureau.

¹ *J. Am. Chem. Soc.*, **39**, 309-315 (February, 1917).

protein, or 22.3 per cent. By mixing the meal with three times its weight of 10 per cent. sodium chloride solution, and this mixture ground in a mill to break up the cells, 20.5 per cent. of protein was extracted ($N \times 6.25$). Dialysis of salt extracts of the jack bean against distilled water yielded about 10 per cent. of pure, dry globulin, based on the weight of the meal used. The globulin was so soluble in salt solutions that it could not be precipitated by diluting these solutions with water. After the globulins had been removed, about 1.5 per cent. of an albumin was isolated as a flocculent coagulum from the dialysate by heating the latter to 75 to 80°. The globulin of the jack bean is not identical with phaseolin, isolated by Osborne from the kidney bean (*Phaseolus vulgaris*). Concanavalin, the globulin present in the jack bean in the smaller amount, and which is the less soluble, was precipitated from a one per cent. salt extract of the meal by adding ammonium sulphate to 0.6 of saturation. The precipitate was filtered off, redissolved in water, and dialyzed until free from sulphates. Canavalin was obtained by making the filtrate from the concanavalin completely saturated with ammonium sulphate. The chief difference between canavalin and concanavalin lies in their sulphur content; namely, 0.48 and 1.10 per cent. respectively. It is evident, from the analyses of these two globulins, that only a small amount of concanavalin can be present in the mixture of globulins obtained by dialysis, since the sulphur content of canavalin and the mixture of globulins are practically the same. Canavalin contains 3.17 per cent. of basic nitrogen. The albumin, which contains 3.73 per cent. of basic nitrogen, resembles the legumelins which have been described by Osborne and his co-workers.

THE PROTEINS OF THE PEANUT (*ARACHIS HYPOGÆA*).¹

By Carl O. Johns and D. Breese Jones.

From the Protein Investigation Laboratory, Bureau of Chemistry,
Department of Agriculture, Washington, D. C.

[ABSTRACT.]

THE recent increase in the production of peanuts in the United States has stimulated a special interest in the peanut from an economic standpoint. The proteins of the peanut have received but little attention. The only published experiments seem to be

¹ *The Journal of Biological Chemistry*, vol. 28, 77 (1916)