THE EFFECTS OF ALCOHOL UPON MENTAL AND NERVOUS DISEASES.*

By H. S. WARD, M.D.
Birmingham, Ala.

In presenting this subject to a scientific body like this, if we had the power to lay aside all preconceived ideas and view facts and not prejudices, what a pleasure would be our task. In reviewing the massive literature, we find much of it argumentative rather than academic. Very few permit the truth to develop, as the facts are evolved; but facts, partial or speculative, are used to prove some preconceived theory. Alcohol is a biochemical substance as old as carbohydrates and the yeast plant, *saccharomyces cerevisiae*. Alcoholic fermentation appears to be an extremely common process. The belief is prevalent that alcohol is produced in the human body by carbohydrate fermentation under the influence of the *bacillus coli communis*:

Historical—In the mythical past, before the dawn of history, man used some substance to help increase the joys of life, or deaden the keen edge of sorrow, and alcohol in some form has been most extensively used. Egyptian monuments and Chinese manuscripts contain the records of the use and abuse of wine. The Vedas contain prayers to the Deity: beseeching him to condescend to come and get drunk with his worshipers, that he might grant their requests and bestow favors upon them, which, when sober, he would refuse. The Old Testament gives graphic accounts of its abuse, and is full of warnings against drunkenness. As civilization has advanced we have learned to make drinks with a higher content of alcohol, and have taught the uncivilized to substitute these for the simpler fermented drinks. Everywhere the use has been bound up in the routine of man's daily toil, in the

expression of his emotions, and in the performance of his religious rites.

Pharmacological, Psycho-Physical—The leading pharmacologists class alcohol, when once in the blood and tissues, with the anesthetics and narcotics, such as ether, chloroform, chloral, paraldehyde, etc. Its use as an anesthetic is much older than that of ether or chloroform. Among bonesetters it was their custom to give large quantities of alcohol to dull the pain and produce relaxation. It was also resorted to in capital operations in the pre-etherization days. Even now it is often used to dull pain for tooth extraction. Is it a stimulant, muscular or mental? This question has called forth a massive and ingenious amount of experimentation. In popular language it is still referred to as a stimulant, but the best experimental evidence goes to prove it to be in no sense a stimulant. In small quantities it generally produces a feeling of well being and good fellowship, along with increased confidence in the powers, both mental and physical. In speaking of a certain type of individual, James says:

"It is the absence of scruples, or consequences, of consideration, extraordinary simplification of each moment's outlook, that gives to the explosive individual such motor energy and ease."

This description applies to the individual who is under the influence of a moderate "quantity of alcohol."

"It tends to turn the inhibitive type of mind into the 'hair-trigger' type. We have said that the speech and bearing of men, the play of their features, all bear witness to the action of alcohol on the brain; that it removes restraints, blunts too acute sensibility, dispels sensations of fatigue, causes a certain type of ideas and mental images to follow each other with greater rapidity and gives a 'cerebral sense of richness.'" (Abel.)

All forms of mental and muscular activity have been tested on individuals accustomed and unaccustomed to the use of alcohol, with and without, and the results seem to be almost unanimous that every form of mental activity is decreased by its use in small or large quantities. Prof. Kraepelin of Munich,

*Read before Alabama State Medical Association, Montgomery, Ala., April, 1914.

one of the greatest psychologists in the world, has made the most extended and exhaustive line of experiments, and is very frequently quoted. He selected a number of accountants of different ages and measured the amount of work for a certain number of days without alcohol. Then gave four cupfuls of claret, to be drunk in small quantities throughout the day. The work fell off three and one-tenth per cent on the first day, and by the end of the week efficiency was reduced 15.3 per cent. Then he took away the wine and at the end of four days they were restored to their former efficiency. Then typesetters were given a measured amount of work, using the printed page as copy, and capacity noted. They were then given alcohol, and at the end of one week their efficiency had decreased 9.6 per cent. Alcohol drunk with meals is less injurious than taken between meals or late at night. Men abstaining from drink during the day, but taking heavy drinks upon retiring, lost on the following day 27 per cent of their coordinating power. Kraepelin was unable to find any time at which alcohol was taken without reducing efficiency. The ability to reason quickly, the power to grasp facts, and to bring complicated ideas into interrelation, these and all other mental processes are slowed down or rendered less active. Among Arctic explorers or soldiers on long marches, those receiving alcohol fall out first.

Cushny, one of the world's leading pharmacologists, says:

"Even the smallest quantities of alcohol tend to lessen the activity of the brain, the drug appearing to act most strongly, and therefore in the smallest quantities, on the most recently acquired faculty, to annihilate these qualities that have been built up through education and experience, the power of self-control and the sense of responsibility."

Helmholtz, that master mind, says that the smallest quantity of alcohol sufficed to dispel from his mind every idea of the creative order when he was trying to give form and being to some dimly seen conception.

Schenedeberg refers every alteration in the

field of consciousness to a primary paralytic action of alcohol, however elusive to analysis and detection the paralysis may be. The stimulation of the emotions and faculties is but fictitious, and is in reality due to the removal of the barriers of self-restraint by the paralysis of the higher functions.

Pathology—The injuries of alcohol upon the body are represented only in part by known anatomical lesions. The burden of the pathological changes in the tissues of alcoholics does not always fall upon the nervous system. Some individuals are much more susceptible than others. The vascular system is often most seriously affected and the other changes are secondary to this. In acute alcoholic intoxication Welch noted the following:

"Two different kinds of lesions of the nerve cells have been found in acute poisoning of animals by alcohol, the one revealed by the Golgi method being of the cellular processes, and the other, shown by Nissl's stain, being of the body of the cells. The former, described as 'The Moniliform change,' is characterized by the appearance of irregular swellings or varicosities in the course of the protoplasmic processes of some of the nerve cells, associated with partial loss of the delicate bud-like or spinous projections normally present on the processes. The other change, designated 'chromatolysis,' is the disintegration of the small, stainable granules, known as the Nissl bodies, which can be demonstrated by certain methods of hardening and staining within normal nerve cells. . . . Nerve cells altered in the ways described have been found in the cerebral hemispheres, the cerebellum, the medulla oblongata, the spinal cord, and the sympathetic ganglia."

I will epitomize some of the changes in the nervous system of 125 autopsies of patients dying with a definite history of excessive alcoholism, as noted by Lambert at Bellevue Hospital (90 men and 35 women). The lesions in the central nervous system seem to be brought about either from the degeneration of the cerebral arteries or from the direct action of alcohol on the nerve cells. In examinations of given brains, it is difficult, or often impossible, to differentiate how much the

changes are due to one or other of these causes. Oedema and congestion of the membranes and cerebral tissue are usually present. Adhesions of the dura to the skull, with increase in the Pachionian bodies, are common and very frequently there is thickening, opacity, and adhesions of the pia. Chronic meningitis combined with atrophy of the convolution was common.

Cerebral degeneration seems to be more common among women and corresponds with the clinical experience that women degenerate mentally from alcoholic excesses more quickly and more completely than men. Pachymeningitis is not uncommon and pachymeningitis hemorrhagica is not infrequently seen in alcoholics, probably from trauma acting on the diseased vessels. All inflammations in alcoholics seem to be prone to the hemorrhagic type and hemorrhagic meningitis was present in three men.

Microscopical examinations of the cerebral tissues show an intense degree of atheromatous degeneration of the minute vessels, which are enlarged, often tortuous, unevenly distended, usually by fusiform dilatations, and their tissues covered with nuclear proliferations. About these vessels the spider or glia cells are crowded in great abundance. These so-called scavenger cells form, with their branching processes, a thick connective tissue felting, densest just underneath the pia, converting the outer fourth of the cortex into a closely matted layer, much diminished from the normal thickness, and often clearly mapped off from the layers beneath. The perivascular spaces are often filled with lymphoid cells. The second and third layers of the cortex show but little change, but according to Bevan Lewis a few of the low, pyramidal cells may be degenerated; the fifth layer of motor cells show extensive fatty degeneration, and, with spider cells beneath, is undergoing disintegration and absorption. The cells show a marked degeneration of their apical processes, and, in many, these have disappeared. The white

matter of the cortex shows equally extensive atheromatous changes in its vessels. The changes found in the spinal cord are similar to those of the cerebral cortex. The paralysis produced by peripheral neuritis was formerly described as of two types—the first, in which the degeneration of the axis cylinder was primary; the second, the interstitial type, in which hyperplasia of the endoneurium and perineurium were primary, pressure atrophy following in the axis cylinder processes. The acceptance of the neuron theory causes us in the vast majority of cases to believe the neuritis of alcoholics to be of central origin.

Insanity.—“Besides the alcoholic deliria, we may have mental disturbances simulating a whole host of other affections, mania-like conditions, melanchollas, chronic persecutory insanity, counterfeiting paranoia, epilepsy, stuporous conditions, morbid impulsiveness, and progressive dementias. The essential characteristic of the whole series of alcoholic insanities, is to be found in the progressive mental weakening with consequent deterioration of the finer ethical and intellectual attainments.” (Berkley.)

In a collective report from a large number of insane asylums of the U. S. A., the percentage of inmates directly or indirectly traceable to the effects of alcohol, was about 20 per cent out of 30,000 inmates. Many of these individuals not quite up to the standard, were unable to bear the stress of life and competition, fell a victim to alcohol, then into the different types of insanity. Without becoming victims to alcohol many might have escaped the asylum. Then indirectly many of alcoholic inheritance become insane in substitution for alcoholism. Thus, alcohol works in a vicious circle. One alcoholic may be the starting point of a family of criminals, degenerates, epileptics, inebriates, and insane.

“The most striking feature is an unreasonable irritability, which frequently leads to outlets of passion of a blindly impulsive character, of which their families or their associates are often the victims. Wife-beating, inhuman treatment of children during the stage of inebriety, attacks upon associates upon the slightest provocation, are of

daily occurrence. The brutalizing influence of the drug, too well known to need minute description here, is shown in the loss of moral sense, indifference to the feelings and sufferings of the family, and to the loss of honor, or position, which inevitably follows the continued indulgence. The alcoholic is wrapped in himself; he becomes a brutal egotist, careless of his work, his standing, of everything that he formerly prized." (Berkley.)

The end results of all types, if life is prolonged, is dementia—to them kind nature's blessing.

Epilepsy—Epilepsy can be produced directly by alcohol in heavy drinkers, proof of which is that when alcohol is given up the convulsions cease. As high as eight to ten per cent of alcoholics eventually have epileptic seizures the first usually after a prolonged debauch. At old Charite in Paris a large per cent of alcoholics were epileptics, supposedly due to chronic alcoholism. Periodic drinkers in ancestors often shows itself in the offspring by epilepsy. There is much evidence to prove that many epileptics were conceived when one or both of the parents were intoxicated. In France, Fere states that among 308 male epileptics, 118 were descendants of alcoholics; and of 286 females, 130 were from alcoholics. Statistics everywhere vary from 30 to 70 per cent of epileptics being descendants of alcoholic forbears.

Heredity—That alcoholism is inherited is a matter of every day observation, and scarcely admits of scientific doubt. The children of abstaining parents, when compared with drinkers' children, give some very astonishing figures. Howe, in Massachusetts, found 145 of 300 idiots were descendants of drinkers. In England, Beech found in 430 idiots that 31.6 per cent were children of alcoholics. In Paris, of 1,000 idiotic epileptics and weak-minded children, 471 had alcoholic fathers, 84 alcoholic mothers, and in 65 both were drunkards. In 209 neither parent was alcoholic. In Normandy, Dahl found that 50 to 60 per cent of idiots had one or both parents alcoholics.

Alcohol being a cell poison, it is not surpris-

ing that the germ plasma becomes so much weakened that the offspring becomes sterile in a few generations and nature thus rids herself of these undesirables.

Alcoholism—An alcoholic is one who either continuously or at certain periods suffers from a craving for alcohol or one of its substitutes.

If you can agree with me that alcoholism is a definite disease, is not an inborn or cultivated "cussedness," then we have a common basis of reasoning. To acquiesce in the popular belief that an alcoholic is one who drinks because he enjoys the sensation, and can give it up or continue it at his will, then we are treating an unfortunate class very unjustly. Old beliefs are given up reluctantly; but the teaching that alcoholism is a disease is not new. The ancient Egyptians recognized alcoholism as a disease and treated it as such. Heroditus, 500 B.C., wrote that "drunkenness showed that both soul and body were sick." Diodorus and Plutarch claim "that drink madness is an affection of the body which hath destroyed many kings and noble people." The Greeks passed laws forbidding women to use wine and restricting young boys. Saint John Chrysostom in the first century asserted with emphasis that alcoholism was a disease like dyspepsia. Other early Christian writers made like statements. The Romans took a similar view. In 1747 Condillac wrote of alcoholism as a disease, and urged its proper treatment. He contended that the impulse to drink was like insanity, an affection of the brain, which could not be reached by law or religion. To Dr. Benjamin Rush, in the latter part of the eighteenth century, is due the honor of giving a clear statement as to this disease. In two essays, entitled "The Influence of Physical Causes Upon Moral Faculties," and "An Inquiry into the Effects of Ardent Spirits upon the Human Body and Mind," he described the disease of alcoholism, dividing it into acute and chronic forms, giving many causes, among which heredity was prominent, and he urged the necessity of

treatment in hospitals for that purpose. During the nineteenth century French, German, English and American writers have produced a mass of literature and agree that alcoholism is a disease, and should everywhere be recognized and treated as such. Heredity and environment are the chief causes. It is a humiliating acknowledgment, but is apparently true that all who are not alcoholics by heredity can become so by its prolonged use.

Prof. Howell says:

"Those who employ it in excess are in danger of acquiring an alcoholic thirst or habit, toward which the body possesses no counteracting regulation. When food is eaten in excess we experience a feeling of satiety which destroys the desire for more food, and some regulation prevails in the case of water. With alcoholic drinks however, the desire may continue long after the alcohol taken has begun to exert an injurious action upon the tissues."

The psychologist, James, has well said:

"The reason for craving alcohol is, that it is an anaesthetic, even in moderate quantities. It obliterates a part of the fevered consciousness and abolishes collateral trains of thought."

Let no one deceive himself into thinking that he can use alcohol with impunity and give it up at his pleasure. Your cells have the same affinity for alcohol as your fallen brother's (or his ancestors once had), and you and yours can fall into the same gutter to curse yourself and be cursed by your abstaining but unsympathetic brother.

Classes—(1) The constant drinker; (2) the periodical drinker; (3) the dipsomaniac; (4) the voluntary drinker; and (5) mixed cases. Then the many types of delirium, and lastly, dementia.

Constant—Those who feel the constant need of alcohol and indulge in it daily. In this type craving is present whenever the supply is stopped. Many who think themselves only social drinkers belong to this class. The test—abstain for one month, and if you find your nerves just as steady without it as with it, then you have not contracted the disease. Second,

those who have no craving between attacks and may even abhor alcohol, and those who indulge to excess. They are often people most fastidious in their taste, correct in dress, without reproach in morals, chaste in speech, and punctilious in their choice of companions. Suddenly desire for drink overwhelms them and they sink into the lowest depths, seeking the vilest companions, becoming filthy in dress and odious in language. They revel in drink and viciousness for a time, and just as suddenly become satiated, completely disgusted with drink, and their companions. They claim a change of heart—never to return to the mire again. Be not deceived; this is only temporary, lasting only a few years or months. The old desire will return; he has not reformed; his tissues are saturated and his stomach nauseated. The intervals become shorter, and the sprees longer, until finally they overlap and the individual is a hopeless inebriate. Third, dipsomaniacs—those who become mentally excited of a maniacal type every time they drink alcohol in any form. These have no craving between attacks, but are always insane during attacks. Many who have given their most serious study consider dipsomania as a form of mental epilepsy. Voluntary drinkers belong to the "cussed" type—those who drink from mere vice, and not to satisfy a physical craving; they drink or leave it alone, at their pleasure. They are not physical but moral alcoholics. These, before they realize it, may become the constant drinker.

Any of these on a prolonged spree may go into one of the forms of delirium. The treatment for the above mentioned types is institutional. Any of these can be sobered up and put upon their feet again; but to be of lasting value, many things are to be considered—and the chief one is the patient's full and unequivocal mental consent to give up and forego all the pleasures of alcoholic drink and its environment. To remove the craving I am of the opinion that there is some specific effect se-

cured by the combined use of atropin and strychnine nitrate. Others use the allied drug, hyoscine. Either of these properly administered will relieve them. But like all types of drug habitues, deep down in their hearts they do not want to give it up. On these one would not care to waste time; but, to the regenerated, the cure should be permanent. To this class, I believe, the state should aid and control. The ideal would be an institution to give a few weeks' treatment to relieve the craving; then some form of outdoor work, either on a farm or public highway for several months to restore them physically and give them moral tone, and time to acquire mental confidence.

There sat two glasses, filled to the brim,
On a rich man's table, rim to rim;
One was ruddy, and red as blood,
And one as clear as the crystal flood.

Said the glass of wine to his paler brother,
"Let us tell tales of the past to each other.
I can tell of banquet, and revel, and mirth,
Where I was king, for I ruled in might,
And the proudest and grandest souls on earth
Fell under my touch, as though struck with blight.
From the heads of kings I have torn the crown,
From the heights of fame I have hurled men down;
I have blasted many an honored name;
I have taken virtue, and given shame;
I have tempted the youth with a sip, a taste,
Which has made his future a barren waste.
For greater than any king am I,
Or than any army beneath the sky:
I have made the arm of the driver fail,
And sent the train from its iron rail;
I have made good ships go down at sea,
And the shrieks of the lost were sweet to me;
For they said, Behold, how great you be!
Fame, strength, wealth, genius, before you fall,
And your might and power are over all.
Ho! Ho! pale brother," laughed the wine,
"Can you boast of deeds as great as mine?"

Said the water glass: "I cannot boast
Of a king dethroned, or a murdered host;
But I can tell of hearts that were sad,
By my crystal drops made light and glad;
Of thirst I have quenched and brows I've layed;
Of hands I have cooled, and souls I've saved.
I have leaped through the valley, dashed down the
mountain,

Slept in the sunshine, and dripped through the
fountain;
I have burst my cloud fetters and dropped from
the sky,
And everywhere gladdened the landscape and eye.
I have eased the hot forehead of fever and pain,
I have made the parched meadow grow fertile with
grain;
I can tell of the powerful wheel of the mill
That ground out the flour, and turned at my will;
I can tell of manhood, debased by you,
That I have uplifted and crowned anew.
I cheer, I help, I strengthen and aid,
I gladden the heart of man and maid;
I set the chained wine-captive free,
And all are better for knowing me."

These are the tales they told to each other,
The glass of wine and its paler brother,
As they sat together, filled to the brim,
On a rich man's table, rim to rim.

—Ella Wheeler.

SOME PHYSICAL REASONS WHY WE ARE WHAT WE ARE.*

By G. H. BENTON, M.D.,
Miami, Fla.

How frequently, in reviewing mentally the distinctive personalities of the collective mass of human beings who continually pass through the busy physician's sphere of more or less critical observation, do we, especially the alienist and neurologist, find ourselves particularly interested in the phenomena responsible for certain personal attitudes, either or both physical or mental, which give to the individual a personality individually distinctive from the general class to which he may belong. These features may lie within or without the range of symptomatology of a given pathological condition which one may at the time be concerned in analyzing, still there often comes to one the desire to ascertain the physical or psychic or combined process through which such end results accrue. We meet continually many numerous

*Read by title in the Section of Medicine of the Southern Medical Association, seventh annual meeting, Lexington, Ky., November 17-20, 1913.