whole affair and mentions certain salient points which must always be taken into account when estimating the actual cost of a troop-horse—such a point, for example, as transport and its necessary accompaniments in regard to forage, and the like. There is also a complaint regarding the cost of working the Indian Remount Department, as, upon the authority of the *Times*, each troop-horse passed into the ranks costs on an average £200. An inspection of the remount farms has been recently made by a distinguished body of army officers and as a result there is a suggestion that an Imperial Stud Department shall be formed to absorb the Remount and Civil Veterinary Departments and thus to bring them under one head. Clinical articles are given on Sarcoptic Mange of the Ox and Demodactic Mange of the Dog and there is an interesting illustrated case of an Actinomycotic Tumour removed from a Cow's Mouth. The remainder of the pages are devoted to extracts from foreign journals, reviews of books, and papers read before various veterinary medical societies during the previous month.

The *Veterinarian* for March has a dearth of clinical articles which seems to show either a very great apathy amongst the members of the veterinary profession or else that the usual recrudescence of disease which comes in the spring has made practitioners so busy that they cannot find time to put their pens to paper. The journal commences with a lengthy extract from a recently published work on Surgery, and is followed by others on Liver Fluke, the Development of Knowledge Concerning Animal Diseases, and Live Stock in Russia. These are followed in turn by a few extracts from foreign journals, some remarks on Current Topics, and Science Notes. The dearth of original matter is a subject for regret.

New Inventions.

SET OF TROCARS AND CANNULÆ FOR EMPYEMA.

MESSRS. ARNOLD AND SONS, London, have made for me a set of special trocars and cannulæ for use in certain cases of empyema. They are flattened from side to side and not circular as usual in other trocars, to enable them to pass through an intercostal space with greater ease. The cannulæ

are made of silver and they can be left *in situ* for 10 days if necessary. The trocars are of very hard steel and in consequence of their shape the end and the cutting edge project further than in the ordinary form. To minimise the danger to deep parts when the instrument is used each cannula is fitted with a moveable shield, being fixed at any depth by a screw as in Durham's tracheotomy tube. The shape of the shield is similar to that of a bivalve tracheotomy tube, but it is perforated by two long slots placed in such a manner that whatever be the inclination of the ribs the tube can be kept in position by tapes threaded through these slots and passed straight round the chest. The set consists of three, with one handle which screws into all the trocars. The sizes are as follows: 1. Small, $\frac{5}{16}$ inch $\times$ $\frac{1}{8}$ inch and $1\frac{1}{8}$ inches long. 2. Medium, $\frac{7}{16}$ inch $\times$ $\frac{1}{8}$ inch and $1\frac{1}{8}$ inches long. 3. Large,

$\frac{3}{8}$ inch $\times$ $\frac{1}{4}$ inch and $1\frac{9}{16}$ inches long. The above are inside measurements; the length is the effective length—viz., from the shield to the end of the cannula. The chief use which I have put this instrument to, and in which I have found it to act extremely well, is to drain an empyema after incision of the skin under eucaïne administered hypodermically in those cases where the administration of a general anæsthetic is especially dangerous on account of the condition of the patient.

WILLIAM TURNER, M.S. Lond., F.R.C.S. Eng.,
Senior Assistant Surgeon, Westminster Hospital; Surgeon,
Royal Hospital for Diseases of the Chest, City-road, E.C.

Looking Back.

FROM

THE LANCET, SATURDAY, APRIL 10, 1824.

LECTURE 49.¹

Thursday Evening, April 1.

On the Different Amputations.

Operations are now much less frequently performed than they were in the days of our ancestors, owing to the great improvements which have taken place in surgical science. Many of the diseases which were formerly considered incurable can now be easily cured by modes of treatment corresponding with our increased pathological knowledge. Many accidents, for example, where the parts are much lacerated, and for which the ancients would have operated, we leave to nature, by whose influence the different reparative processes will be set in action and the injured limb restored to health and utility. When amputation is necessary, nature will occasionally even perform this operation unassisted by art; in mortification of the feet it often happens that the leg will be amputated by nature as effectually as though it had been accomplished by the amputating knife. At Guy's Hospital there is at the present moment a case of this description. You have seen, in the case which I allude to, first, a division of the skin, then the division of the muscles shorter than the skin, and lastly, the division of the bones; the fibula has already separated, and the exfoliation of the tibia is rapidly going on. Nature in this case if left to herself, would, without doubt, accomplish the amputation of the leg, but the safety of the man requires, I think, that the remainder of the bone should be divided by the saw, for if this be not done, the long continued excitement may wear out the powers of the constitution.

Diseased joints used very frequently to lead to the performance of amputation, in the young as well as in old age; but amputation is much less frequently performed at the present day, in consequence of such disease, than some years back; even diseases of the joints of the upper extremities of children give rise to amputation much less frequently than in the days of our forefathers, but in chronic diseases of the ankle and knee, amputation is still very commonly performed; there is, however, a marked distinction in these chronic enlargements; one variety may be called congenital, which exists from the birth of the child, and the other is from some debilitating cause which produces the complaint after birth; as the constitution therefore was radically weak or vitiated in the former of these affections, you cannot expect that such permanent benefit will result in that case, as in the latter, where the constitution became affected from some accidental circumstance.

With respect to diseases of the ankle and knee joints, amputation for such complaints will occasionally be necessary both of the leg and thigh; indeed chronic affections of the ankle and knee give rise to amputation as frequently as diseases in any parts of the body. For compound fractures we seldom amputate directly: they are seldom so severe as to require immediate amputation, and it is not until gangrene or disease of the bone has taken place, that it is deemed necessary to amputate; compound fractures, however, from the superior manner in which they are now treated, do much better than formerly, and very severe injuries of this description will often terminate most favourably, therefore upon the whole, amputations are much less frequently performed at the present epoch than in the days of our ancestors.

¹ Excerpt from report of lecture delivered by Sir Astley Cooper in the theatre of St. Thomas's Hospital.