refreshed, and took a small quantity of nourishment.—At 10.30 P.M. a severe fit came on, very similar to the one he had in the morning, but of greater intensity; the body suddenly raised; muscles of back of neck, trunk, and lower extremities powerfully affected; body regularly formed into a bow (opisthotonos); an expression of great mental distress and anguish in the countenance; jaws firmly fixed; the whole face greatly distorted and disfigured; breathing hurried and short; pulse weak and irregular; surface of body covered with profuse sweat; muscles of chest and abdomen felt like hard pieces of board. Patient fell back perfectly exhausted and gasping for breath. He was offered a variety of things in the shape of nutritious fluids; he could swallow nothing; a horrible dread if he took anything that another spasm would come on. Injections of morphia were given with no benefit, the spasms coming on with greater frequency and severity. Between the fits the body was hot and damp; temperature 104.6°; pulse rapid but weak; the slightest noise or movement brought on fresh convulsions.

At 6 A.M. of the morning of the 21st the patient was attacked with a final spasm; gave a sudden jerk, and was completely thrown off the bed; took two or three men to hold him during the fit; features assumed a horrible appearance, lips becoming blue; body fearfully convulsed; muscles rigid and tense. Death closed the awful scene.

The man died on the ninth day of the accident, and some fifty or sixty hours after the first signs of tetanus set in. The mind was clear from the beginning to the end, and there was no want of control over the bladder or bowels. Death appeared to take place from suffocation combined with extreme exhaustion.

(To be continued.)

NOTES ON A CASE OF TUMOUR OF THE CEREBELLUM.

By RICHARD CATON, M.D.,

LECTURER ON PHYSIOLOGY, LIVERPOOL SCHOOL OF MEDICINE; ASSISTANT-PHYSICIAN TO THE INFIRMARY FOR CHILDREN.

ON the 15th January, 1875, I was asked to see A. B—, a man somewhat under the middle height, of healthy appearance, aged twenty-eight, the manager of a large farm.

History.—About fifteen years ago patient had an attack of rheumatic fever, which was believed to have left behind it some cardiac affection, which, however, of late years had produced no symptoms of importance. Patient described himself as enjoying excellent health, and was apparently capable of active muscular exercise. About four months ago, when spending too much time in reading after his day's work was over, he suffered for a day or two from occipital headache. On discontinuing the reading this ceased.

Fourteen days before I saw him, having been somewhat short of labourers on his farm, he had helped the men to carry some distance, and to lift up a ladder, a large number of sacks, each weighing about 1½ cwt., the exertion being extremely severe. On that evening he was attacked by a violent pulsating headache in the occipital region. This had remained almost constant up to the time of my seeing him, and was so severe as to prevent sleep and render exertion impossible. He had had two or three attacks of vomiting.

Symptoms when first seen.—Severe pulsating headache in the occipital region complained of. Patient, who is a highly intelligent man, is quite clear intellectually, but is unable to bear much conversation, is very sensitive to noises, and suffers much from tinnitus. He exhibits some degree of difficulty and unsteadiness in walking; has no paralysis of motion or sensation; sight unaffected. He cannot recline on the back without increase of pain. Tongue furred; bowels constipated; no appetite; respiratory sounds normal; a remarkably loud, double aortic murmur; cardiac dulness perhaps slightly increased; pulse about 120, markedly aortic; temperature rather above the normal range; urine normal. He has always been strictly temperate, and has never had syphilis.

The *diagnosis* presented some little difficulty. I was in doubt between two theories—firstly, that of the existence of structural lesion of the brain; secondly, cerebral anæmia or some disturbance of brain circulation, due perhaps to mechanical injury of a previously diseased valve, sustained during the violent exertion on the day from which the symptoms dated. I rather inclined to the latter theory.

The *treatment* consisted in rest, the employment of counter-irritation over the nape of the neck, gentle aperients, and bromide of potassium, and subsequently iodide of potassium.

Progress of the case.—Coincidentally with the treatment referred to, improvement took place in some of the symptoms. The headache, instead of being constant, became intermittent, the remissions became longer, and, finally, in about ten days the headache had become only occasional; any excitement or over-exertion would bring it on. The appetite returned. Notwithstanding improvement in these respects, great and increasing muscular weakness was observed; the muscles were largely developed, but the patient, who a fortnight ago had been capable of muscular exertion such as few of his farm labourers could surpass, now could not ascend two pairs of stairs without assistance. His gait began to be more unsteady; he could not turn round with the eyes shut without falling. Eyesight perfectly good, but a remarkable slowness in accommodation manifested itself. In adjusting the focus of the eye from an object one foot distant to another at fifty feet, a delay of three seconds took place. I measured this carefully on several occasions. The movements of the eyeball were normal; spectra in the form of blue rings were described.

These symptoms again excited fears of the existence of some structural lesion (perhaps of the cerebellum), and I decided to have a careful ophthalmoscopic examination made. Mr. Edgar Browne, surgeon to the Eye and Ear Infirmary, reported the presence of commencing optic neuritis about both discs, with dilated and tortuous vessels and one or two small hæmorrhages. This evidence was considered conclusive as to the existence of coarse disease, tumour, apoplexy, or abscess of the cerebellum being suggested. A very unfavourable prognosis was given. The eyesight as yet was quite good, but, notwithstanding appropriate treatment, the clouding of the retina was seen by us to spread steadily day by day, until at length, within about ten days from its first discovery, it produced loss of sight. About the same time the general symptoms became more severe, pulse and temperature rose, and occasional delirium occurred. Rheumatic symptoms were complained of, and pain in the loins drew attention again to the urine, which a few weeks back had been quite normal, but now was found loaded with albumen and containing numbers of granular tube-casts. Severe night-perspirations occurred; temperature often rose to 103°; pulse ranged from 110 to 130. The patient's hearing, which had formerly been morbidly acute, was now noticed to be defective on the left side. Several weeks now elapsed, during which no important change took place. Pain in the cardiac region and left hypochondrium were occasionally complained of. He suffered from a sense of oppression in the head, but only at intervals from headache, and was usually quite clear intellectually.

Towards the end of April, cedema of the legs appeared, and some fluid collected in the peritoneum. Occasional squint and signs of partial paralysis of the facial muscles of the right side became manifest. About the 1st May he began to have difficulty in expressing himself, on the 3rd became gradually hemiplegic on the right side, and died on the 4th.

Autopsy.—Cerebral hemispheres normal externally. A tumour, the size of a small walnut, was found on the under surface of the left hemisphere of the cerebellum, adjacent to the pons, and beneath the auditory nerve, which was stretched over it. There was a small embolus in the left middle cerebral artery, beyond its first anterior branch. The ventricles contained two or three ounces of fluid.

Remarks.—The sudden onset of the symptoms immediately after violent muscular exertion, valvular disease being present, was very misleading, and caused at first an error in diagnosis. Subsequently the diminution of muscular power without paralysis, the ataxia, together with the occipital headache and optic neuritis, formed a group of symptoms on the whole characteristic of tumour of the cerebellum. The affection of hearing gave, later on, a clue

as to the probable locality of the tumour. The slowness of accommodation is a symptom which I have not seen described before. It was very marked. If the patient glanced up from the book he was reading towards the figure of any person entering the room, he described himself as only being able for two or three seconds to see a dim haze, then he saw quite distinctly. Changes in focal adjustment were obviously made slowly and with difficulty. Prof. Ferrier has shown how largely the cerebellum is concerned in the co-ordination of the muscles of the eyeball; has it a like control over the ciliary muscle and the mechanism of accommodation?

Liverpool.

EMPYEMA, FOLLOWING SCARLET FEVER: THORACENTESIS; RECOVERY.

By BENJAMIN WALKER, M.R.C.S., L.R.C.P., &c.

SARAH JANE W—, aged two years and a half, the child of healthy parents, was seized with convulsions on the 5th of April. The attack lasted about half an hour, and the left half of the body was chiefly affected, the left hand being very blue and the face much distorted. The left half of the body was quite powerless. Speech was not affected. On the following day the characteristic rash of scarlatina appeared. The nearest medical man was called in, and he attended her till the 16th, when she came under my care. This was the twelfth day of the illness. There was then considerable though not complete loss of power of the left half of the body; tongue dry and glazed; pulse 130, feeble and compressible; skin hot and dry. The temperature could not be taken at any time through the illness on account of the restlessness and fear of the child when approached. Throat affection slight. There was great thirst. The bowels had been freely opened by my predecessor. Bark and ammonia were now ordered, with beef-tea, egg, and brandy mixture at regular intervals, and, to quench the thirst, milk and lime-water *ad lib*.

May 1st. — Condition was one of hectic, with frequent flushings and rigors, and the child was greatly emaciated, the support being taken in a very desultory manner.

8th. — An abscess of considerable size under the left pectoral muscle was opened, and gave exit to about three ounces of fairly formed pus; there was a slight drain for a few days and then the discharge ceased. In the course of two weeks the left side of the chest began to increase in size, so that the measurement exceeded the right by about an inch; the heart was displaced to the right, so that the apex beat was to the right of the sternum, a little below and to the inner side of the right nipple. There had been a troublesome cough for three weeks past, latterly increasing in severity, but without much expectoration, till the 17th, when there was a copious discharge of pus tinged with blood, denoting the occurrence of bronchial fistula. Some relief followed this, and purulent matter was from time to time expectorated with the cough, in greater or less quantity; the chest measurement, however, was little affected, and the child continued in much the same state for the next six weeks, when the dyspnoea became daily more marked, and it was evident that, unless relief were obtained by other than nature's method, there would soon be a fatal termination, for the little patient was emaciated to an extreme degree and rapidly sinking from hectic.

July 15th. — With the assistance of my friend Mr. Foulds, who saw her in consultation, a medium-sized needle of Maw's bottle aspirator was inserted, between the sixth and seventh ribs near their angles, and 20 oz. of odourless pus of the consistence and exact appearance of pea-soup drawn off. A little blood commencing to flow warned us to withdraw the needle. The lung gradually expanded whilst the fluid was being withdrawn, as indicated by the cough, and afterwards verified by auscultation and percussion. No anæsthetic was used, and the operation was the simplest in performance, only a sharp twinge of pain being felt at the moment of inserting the needle. Immediate relief followed the operation; the breathing was easier, the cough less

troublesome, and purulent expectoration ceased from this date. Next day the child was better, and there was no irritation or discharge at the seat of puncture, which was seen as a mere spot, and soon entirely disappeared. The relief, however, was but temporary; for in three days (the 18th) the distension of the pleura was as great as before, and the aspirator was again employed, and a little more than a pint of odourless pus similar to the former drawn off, with relief to all the urgent symptoms.

The exhaustive and able "Clinical Remarks on Tapping and Draining the Pleura," by Mr. Berkeley Hill, appeared in THE LANCET just at this time (July 17th), and I derived practical benefit from the excellent rules there laid down.

The purulent effusion again distended the pleura, displacing the heart as much as before, but more slowly. The condition of the child was one of extreme emaciation, the limbs being apparently little more than bones covered with skin. The two aspirations having yielded perfectly inodorous pus made me unwilling to insert a drainage-tube; but, as she was evidently sinking, with the assistance of my former colleague, a trocar and canula was introduced, which gave exit to about a pint of similar pus. The opening was then enlarged, and a No. 9 Holt's winged catheter introduced. Through this a pint more, at a moderate computation, gradually drained in the next twenty-four hours. The end of the tube was surrounded by a loose pad of tenax to absorb the discharge, and this changed daily. The steady improvement inaugurated by this operation showed it to be the correct one. The child lost all its former hectic and febrile symptoms, and began to take nutriment in almost any shape in which it was offered. In the course of a week or two the appetite was ravenous, and chops and steaks were disposed of in a manner which would certainly have taxed the digestive powers of a healthy adult. Sleep became sound and refreshing, the child awaking once or twice in the night, only to ask for food. In addition to solids, large quantities of milk, and of egg and port mixture, were disposed of in the twenty-four hours. The cough and expectoration both disappeared, and in a month's time she was as plump and firm-fleshed as ever in her life. The increase in the girth of her limbs could almost be graduated daily.

Aug. 28th. — Began to take cod-liver oil emulsion, and, in addition to bark, Parrish's syrup.

At present (Oct. 20th) the discharge from the tube varies, some days being very slight, others more profuse, the weather influencing this in some measure; besides which, the house in which the patient lives is very damp. She runs about as in health, and on fine days goes into the air.

Remarks. — The advantage of giving free and permanent opening in empyema, at whatever age occurring, is clearly exemplified by the result of this case. The milder operation of removing the fluid by aspiration is less repugnant, especially in so young a patient; but the reaccumulation in so short a time as three days, with its accompanying hectic, shows that it is but tentative; and on a similar case recurring I should advise drainage after a first aspiration, as giving a better chance of recovery than a repetition of the less severe operation, regardless of the state of exhaustion of the patient.

The subject of antiseptic surgery is again coming to the front, as appears from a leader in THE LANCET of the 16th inst.; where there is also recorded (at p. 577) a case of Mr. Pemberton's, at Birmingham, in which "extremely successful results" have been obtained in a case of empyema, "where the chest was opened antiseptically and a drainage-tube run through it, the man having gained weight in a remarkable manner." I think, without any invidious comparison, this little patient shows that equally satisfactory results may be obtained, with care and cleanliness, without antiseptic surgery.

I would also say a word in favour of the simple and inexpensive bottle aspirator of Messrs. Maw and Son, which acts most efficiently. The winged catheter makes an excellent drainage-tube, though it is well to attach it by a thread secured by a strip of diachylon, as, if the wings alone be trusted to, the tube may escape (as occurred once to me), and give some trouble in its reinsertion, especially in a very young patient.

The parents will be most happy to show the child to any practitioner desirous of seeing it.

Spondon, Derby.