purse suture that most reliance requires to be placed, and it is therefore very important to see when this is introduced that an equal grip is taken of the conjunctiva all round the free edge and also that too wide an interval is not left between the stitches, but the catgut strands to which the muscles are attached afford great additional security. In one of my recent cases in which the suture holding the internal rectus muscle broke, a little opening made its appearance on the seventh day after the operation opposite the attachment of that muscle and the aperture grew gradually larger and larger until the paraffin escaped.

In my series of 22 cases there have been four failures. Thrice the paraffin has come out through inefficiency of the stitching, but in two of these the conjunctiva was simply drawn together by the catgut holding the muscles without any continuous suture. The purse suture was introduced in my fourth case and since then the only occasion on which the stitches have given way was that referred to above where the suture securing the internal rectus muscle broke. On one occasion, after the enucleation of a suppurating eyeball, pus appeared in the socket and set up so much inflammatory reaction that the stitches were removed and the moulded paraffin was allowed to escape. The paraffin employed is a sterilised preparation melting at 40° C., specially prepared by Rogers of Oxford-street, London. It seems simply to lie in the tissues and gives rise to no irritation, but whether it will at length become absorbed or not time alone can show. Up to the present none of the cases have given any sign of either shrinkage of the stump or impairment of its mobility.

Glasgow.

SUNSTROKE.

By J. GRAY DUNCANSON, M.B., C.M. GLASG.

THE following brief notes of a fatal case of sunstroke present a clinical and pathological picture which might almost be accepted as typical of an accident, I fear, rather more common in this country than we are led to suppose by the text-books.

The patient was a thick-set muscular man, aged 44 years, 5 feet 5 inches in height and in weight about 12 stones. He had been a heavy drinker for over 24 years, mainly of fourpenny ale and spirits. He had never previously had any severe illness and his family history was good. On June 30th, 1902, after he had been drinking more than usual, he felt ill and on July 2nd he was found to be suffering from a sharp pain just outside the left nipple. His tongue was coated and his skin was moist. On the 4th, as this was evidently due to pleurodynia, a mustard leaf was applied and he was directed to continue taking a mixture composed of sodium salicylate, solution of bismuth, and ammonia. On the 6th he left home at 11.30 A.M. for a walk; he had previously drunk a pint of fourpenny ale and half a quartern of whisky. The day was close and muggy and the sun was shining brightly; the temperature in the shade was 82° F., probably about 115° in the sun. He was observed to be lying under a tree and was spoken to by an acquaintance at 12.30 P.M.; at 1.30 he was found quite unconscious and was taken home in a cab. I saw him at 2.30 when his face was flushed and dusky, his breathing was rapid (45 to the minute), and irregular in rhythm and depth (Cheyne-Stokes). He was unconscious but was restless, continually moving his arms and legs in an aimless manner. The knee reflexes were in abeyance, the teeth were clenched, and the eyes were closed, the pupils being contracted, and the conjunctivæ sensitive to touch. The pulse was 125 to the minute, full, bounding, regular and fairly compressible. As regards the heart, the first sound over the mitral area was rather prolonged and muffled, the aortic second sound being sharp and loud with a snap; there was no murmur. The skin was intensely hot and dry, there being absolutely no sweat either on the face or body; the temperature was not taken. There was a small swelling over the occipital bone on the left side. At 10 P.M. the patient's temperature was 105.5° F., the respirations being 36 and the pulse 96. He had vomited once, bringing up a greenish-yellow fluid, and he had voided urine several times unconsciously. When called upon loudly by name he opened his eyes. A soft murmur, systolic in rhythm, was now heard over the apex of the heart (mitral), and the respirations were shallow but more regular. He was

able to swallow when fluid was poured into his mouth. The respiratory murmur on the left side was harsh with moist râles, this condition being less pronounced on the right side. The abdomen was distended. On the 7th, at 10 A.M., the temperature was 105° F., the respirations being 36 and the pulse 96. He had passed a very restless night, the unconsciousness being deeper, the face more dusky, and the right side was observed to be partially paralysed, the right eye discharging muco-purulent fluid. The left arm and left leg were now seldom moved and there was some difficulty in swallowing. The pulse was weaker and softer; it was regular in rhythm but not in quality. The respiration was markedly Cheyne-Stokes in character and there was phlegm in the bronchi and in the trachea. Nourishment was taken fairly well. At 7 P.M. the temperature was 104°, the respirations being 44 and the pulse 100. There was no improvement in the patient's general condition. Respiration had now become rapid and shallow, but was not so irregular. The skin felt cooler, the knees were cold, but the feet were hot. The man was picking at the bedclothes with the right hand. On the 8th the temperature was 103°, the respirations were 44, and the pulse was 106. At 7 A.M. on this day he regained partial consciousness, recognised his wife, and answered "yes" and "no" to questions put to him, but he was not capable of carrying on any intelligent conversation. His breathing was now almost regular and there was less phlegm. The pupils were not contracted and he was able to move the left arm and leg as well as the right. At 2 A.M. he had struggled violently and had tried for the first time to get up. Urine was voided unconsciously and there was an action of the bowels. The evening temperature was 103°, the respirations were 44, and the pulse was 100, weaker and very compressible, the vessel filling and emptying suddenly. The râles over the front of the chest and at the bases were copious, with dulness to percussion. The heart was dilated. The general condition of the patient was worse. On the 9th the patient's temperature was 104°, the respirations were 50, and the pulse was 120. The patient had had a fairly quiet night. His face was dusky, his skin was hot and dry, and his breathing was laboured and rapid. His pulse was regular, soft, and weak; his abdomen was tympanitic, and the dulness at the back part of the lungs had increased in extent. At the left side in front the breathing was tubular, and there were râles all over the right side. The tongue was coated but was not very dry. There were no convulsions, but the cough was troublesome. The patient had all through taken milk, eggs, beef-tea, and ice. He died at 8.30 P.M.

A post-mortem examination was held on the 11th. The body was that of a well-nourished man; rigor mortis had been but transitory, the face, the upper parts of the trunk, and the limbs were already decomposing. There was slight superficial swelling over the occipital bone on the left side. The pericranium was rather adherent and the cranium was hard. The dura mater was tense and it bulged on the removal of the skull cap. The blood-vessels were distended and the sinuses, especially the lateral sinus, were full of dark quite fluid blood; the arteries were somewhat rigid. The sulci of the convolutions about the upper end of the fissure of Rolando were covered with yellowish-coloured, irregular masses of semi-purulent material. These were closely associated with the venous branches and were situated in the subarachnoid space; they were not easily removed but clung tenaciously to the vessels and membranes. This condition was more marked on the right side. The venous stems throughout the brain substance, which was slightly softer than usual, were distinctly distended, and when cut fluid dark blood exuded. No distinct hæmorrhage was discovered, though all other parts of the brain were examined. There was little cerebro-spinal fluid. The lungs were engorged with dark fluid blood and at their bases and in the posterior parts were in the condition of splenisation, dark and non-crepitant. No pleuritic adhesions were found; there was blood in the pleural cavity. The pericardial space contained little fluid. The mediastinum was filled with masses of fat. The heart was laden with fat, but the left side was fairly well supplied with muscular fibre. The right ventricle was dilated and contained fluid blood; the left ventricle was empty and dilated. The whole organ was much enlarged and flabby. The mitral valve was incompetent and the aortic valve held water; the aorta immediately above the cusps of the latter valve was the seat of atheromatous change—there was a calcareous plate at one spot one-third of an inch by one-quarter of an inch in size. The stomach, the great omentum, the mesentery,

and the abdominal walls were laden with fatty masses. The liver was smooth, engorged with blood, and oily, and the gall-bladder was full of greenish-yellow bile. The spleen was dark, enlarged, and friable. The kidneys when stripped of their capsules both displayed several deep cup-shaped depressions, evidently due to old embolic infarctions, the general picture upon section being distorted. The intestines were distended and of a pale colour; the bladder was hypertrophied; and the muscles were pale and exuded little blood.

It is recognised in all hot countries that the back of the neck immediately below the occiput is the important spot to be protected, both in man and beast, from the direct rays of the sun. These solar rays, I feel sure, act in the first place directly, and they do so by causing a disturbance in those vaso-motor centres which are situated in the medulla, affecting the proper discharge of their functions, and specially interfering with the action of the sympathetic system. The blood-vessels of the brain are enormously dilated, also those of the cord and the thoracic and abdominal organs; but those throughout the voluntary muscular system do not appear to be affected. The contracted pupils, absence of sweat, Cheyne-Stokes respiration, and hyperpyrexia, all primary symptoms, point to this conclusion. As suggested by Sambrook, it is not impossible that a microbic influence is at work and the early appearance of lepto-meningitis would tend to give support to this theory.

The essentials of treatment are ice, enemata, ice-baths, &c., and if a hose be handy it should be turned on to the patient for a time, as is done on board ship, in order to lower the surface temperature. A small dose of calomel should be given at the outset; then, every hour, and afterwards less frequently, a draught such as this: 20 minims of solution of acetate of ammonium, 15 grains of bromide of potassium, and 20 minims of spirit of nitrous ether in water. I might be inclined in another severe case to try a hypodermic injection of pilocarpine to induce perspiration if I considered the heart would stand it. Then, to improve the flagging circulation, stimulants must be employed and, when the time comes, digitalis—not strychnine.

Shooter's Hill, S.E.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

MEASLES AND CHICKEN-POX OCCURRING SIMULTANEOUSLY.

BY EDWARD J. BLACKETT, M.R.C.S. ENG., L.R.C.P. LOND.,
SURGEON TO THE HOSPITAL OF SS. JOHN AND ELIZABETH,
GROVE-END-ROAD, LONDON, N.W.

THE following brief notes of two cases which were under my care in the early autumn of last year are, I think, of some interest owing to their clinical rarity, for as far as I have been able to gather the combination of chicken-pox with measles is certainly uncommon.

I received a message asking me to go to see a boy and I must own that I went with a biased opinion, as the wording of the message was: "Will you come and see my boy who has measles?" Accordingly I fully expected to find a typical case of measles when I got to the house. Before I saw the patient I elicited the fact that he had been home from school about three days and that there had been an outbreak of measles there before he left. I saw the patient, a boy, aged 13 years, in bed. His face was a little flushed and the eyes were slightly—in fact, very slightly—congested; there was no coryza. The temperature was 100° F. There was no sign of a rash on the face, the chest, or the arms. On examining the back I saw three or four typical chicken-pox vesicles but no sign of a measles rash. I asked the patient if there had been any other illness at the school besides measles and he told me that there had been a few cases of chicken-pox and by further questioning I found that he had seen one of the boys with it about ten days before and had also about the same time come in contact with a patient suffering from measles. So I came to the conclusion that the boy

had chicken-pox at present, whatever else he might be going to have in the future. On the next day his temperature was still about 100° and there were no fresh vesicles, but a morbilliform rash was making its appearance on his face, chest, and back, being most marked on the back. His younger brother, who had been exposed to the same sources of infection, I saw for the first time. He was listless and complained of headache. His eyes were more congested than were his brother's, but there was no coryza or cough. I could find no signs of measles about him, but, as in the case of his brother, a few chicken-pox vesicles were seen. On the next day the elder brother exhibited a characteristic measles rash and the younger one was showing faint signs of the rash, and on the following day the measles rash was well marked on the younger as well as on the elder brother. Nothing further of interest occurred during the progress of the two cases and both boys were convalescent in a fortnight.

The main point of clinical interest in these cases was, I think, the fact that when the rash of measles appeared it completely masked the chicken-pox vesicles and if I had not seen the cases so early I would have looked upon them as simple cases of measles, but the few vesicles I saw first were undoubtedly those of chicken-pox and the history of the cases made my diagnosis, I think, a correct one.

Bryanston-street, W.

A CASE IN WHICH HOUR-GLASS CONTRACTION OF THE UTERUS IN A CASE OF TWIN BIRTH ARRESTED THE EXPULSION OF THE SECOND FŒTUS.

BY STANLEY J. KERFOOT, M.R.C.S. ENG., L.R.C.P. LOND.

ON Jan. 4th I was sent for to attend a married woman, aged 29 years, a multipara, who thought herself to be from five to six months pregnant. Labour was well advanced, the pains occurring about every ten minutes. On vaginal examination the os uteri was found to be fully dilated and the membranes were presenting, which I ruptured. A hand presented and it was necessary to turn to deliver. External examination then indicated a second foetus and on vaginal examination a second bag of membranes presented, which I ruptured. After waiting for some time, during which the woman had four pains, and being unable to discover any presentation, I followed up the cord of the first foetus into an apparently empty uterine cavity. The funis was then found to ascend through a very small constricted opening, into which with difficulty only one finger could be inserted. Gradually, this resistance was overcome and eventually I was able to insert my hand, grasp a foot, and speedily deliver the second foetus. Both placentæ were then removed without the contraction occurring again. I have never had a similar case in my own practice nor can I find any reference to such a case by any of the authorities.

Bedminster, Bristol.

RÖTHELN AND MEASLES

BY J. REID, M.A., M.D., C.M. ABERD.

THE following brief notes of four cases of possible double infection may be interesting.

CASE 1.—On Dec. 26th, 1902, I was called to see a girl, aged 12 years, on whom a rash had been observed by the mother on the 25th. There were slight bronchial catarrh, inflammation of the tonsils and pillars of the fauces, with the typical rash of measles present everywhere except on the face and head. There was no coryza or inflammation of the eye. The disease ran a mild course, the eruption disappeared on the sixth day, and the lungs remained clear. The throat alone gave some little trouble. Koplik and Filatow spots were absent. On Jan. 4th the skin of the fingers peeled off in large pieces. On the 6th ulcerative stomatitis and cardiac bruit were present. On the 10th there were chorea and pericardial friction sound at the base, with irregular heart and sleeplessness. On the 11th she had improved in every respect and desquamation was complete.

CASE 2.—On Dec. 29th a sister of the above patient, aged 13 years, was brought under my notice for a rash which appeared on the previous day. This was a typical case of measles with Koplik and Filatow spots, coryza, rash on the face, &c. There was as sequela rheumatic pain in one foot,