a satisfactory one or otherwise. The asphalt should be one with a high percentage of so-called petroleum as compared with asphaltene; the oily or bituminous liquid above referred to should be one that will make a satisfactory and durable blending with the asphalt, and the proportions of tempering liquid and asphalt which go to make up the cement should be chosen with reference to climate and the character of the traffic that is likely to pass over the street.

The analyses quoted in this lecture will show how asphalts may differ in the first respect. With regard to the nature of the tempering liquid, I may say that coal tar was at one time used, but has been almost universally rejected, as not making a satisfactory or durable blending material, the mixture having been found to disintegrate under atmospheric influences. Petroleum residuums are now almost universally used in the Eastern States, while malthas, or natural "liquid asphalts," so-called, are used on the Pacific Coast and in Western States. I am satisfied, on both theoretical and experimental grounds,* that the last-named tempering materials are to be preferred. The time allowed for an illustrated lecture on a broad, general subject of the nature of this will not allow me to go into a discussion of this question here. Thanking my audience for their kind attention, I will therefore bring my account of asphalts to an end.

RAINFALL AND TYPHOID FEVER.

BY WM. P. MASON.

The first, or at least one of the first, to call attention to the relation between water and typhoid fever was Dr. Michel, of Chaumont, France. In 1855 he observed that typhoid, which was epidemic in the above place, varied in number of cases and in intensity inversely as the quantity of water in the public wells.

Pettenkoffer, of Munich, about the same time, undertook extended observations upon variations in the height of

* *Paving and Municipal Engineering*, September, 1894, p. 118.

ground water, and, a little later, relationship was shown between these variations and the occurrence of typhoid fever.

Those who hold with Pettenkoffer, claim that the elements of the disease readily multiply in the soil, and are driven therefrom along with the ground air, upon the rising of the water level at the time of the autumnal rains.

Latham, in speaking upon this point, says:

"No great variation in the vertical rise and fall of sub-soil water is the healthier condition. The ground always contains air, and, as the ground water sinks, air is drawn in to supply its place. After long dry weather, the air of the soil is thus laden with products of decomposition. A rain now occurring, the ground air is displaced, and since said rain is liable to seal the surface, the tendency of the air is to escape laterally, *i. e.*, into cellars. Dry summers invariably mark unhealthy years. Typhoid fever occurs after the autumn rains.

"All the great epidemics of typhoid have occurred in years when the ground water was especially low, and after a slight rise in the same."

Pettenkoffer's "ground-air theory" is not gaining the majority of supporters, a more reasonable view being, that as the water surface lowers in a well, the base of the cone of drainage, whose apex is at that surface, is extended, and consequently, more widely situated points of pollution are embraced within its influence.

Perhaps the most exhaustive examination of the relation of the height of ground water to the prevalence of typhoid that has been made in America, is to be found in the work of the State Board of Health of Michigan.

Observations have been made by that Board during a period of many years, and the results, graphically shown herewith, indicate in a very marked manner that increase of typhoid and lowness of water in wells move in practically the same curve of variation.

So convinced were the Michigan authorities of the truth of this proposition, that they issued, during the past autumn, a circular of warning, which is here quoted in part:

"Beware! Unusual Danger now from Typhoid Fever, because of Drought.

"The water in the representative well near the center of the State, last September, was three inches more than the average of previous years; this year it is four inches less than the average.

"For the second week in September, typhoid fever is reported from thirteen places more this year than last year, etc., etc."

The Michigan statistics go, further on, to show that in October, 1894, the water in the standard well stood eleven inches lower than in October, 1893, and seven inches lower than the October average for the eight years, 1886-1893.

For September, 1894, typhoid fever was reported from 121 places in the State, an increase of forty-six places over the report for September, 1893.

For October, 1894, typhoid was present at 165 places, as against 109 for the same month of 1893.

For October, 1894, the prevalence of the disease is forty-four per cent. above the October average for the eight years, 1886-1893.

During the three months of September, October and November, 1894, the ground water of Michigan grew constantly lower. It is difficult to see just how these data could be made to fit the "ground-air" theory of Pettenkoffer or Latham as a cause of typhoid fever; for such theory calls for sudden rise in ground-water level.

The precipitation data for Michigan are:

	1893.	1894	Normal.
January	2'55	1'88	2'18
February	2'65	1'81	2'67
March	2'39	2'16	2'32
April	4'43	2'28	2'44
May	2'79	5'79	3'52
June	3'26	2'82	3'91
July	2'74	1'40	3'09
August	1'19	0'49	3'04
September	2'34	3'42	3'00
October	3'67	2'86	3'05
November	2'90	1'76	3'02
December	3'64	1'33	2'51

It has been the continued experience in Michigan that typhoid is coincident with low ground water, as is illustrated graphically herewith, and is not dependent upon sudden rise in the same. An interesting exception to this rule has been noted, occurring during the season of heavy frost, when surface pollution is prevented from reaching the subsoil.

Dr. Henry B. Baker, Secretary of the Michigan State Board of Health, to whom I am indebted for much information, also furnished me with this table :

EXHIBITING THE AVERAGE TOTAL ANNUAL RAINFALL AT STATIONS IN MICHIGAN THE SAME FOR LANSING, THE INCHES OF EARTH ABOVE THE GROUND WATER AT LANSING, THE INCHES OF WATER IN AN UNDISTURBED WELL AT LANSING, AND THE REPORTED SICKNESS, FROM TYPHOID FEVER IN MICHIGAN, AS INDICATED BY THE PER CENT. OF ALL THE WEEKLY CARD-REPORTS WHICH STATED THE PRESENCE OF TYPHOID FEVER DURING THE SEVEN YEARS AND EACH OF THE SEVEN YEARS, 1885-91 :

YEAR, AND PERIOD OF YEARS.	Average Total Annual Rainfall at Stations in Michigan, in Inches.	Total Annual Rainfall at Lan- sing, in Inches.	Inches of Earth above the Ground Water at Lansing.	Inches of Water in an Unused Well at Lansing.	Ground Water Higher (+) or Lower (-) than the Seven Years' Average in Inches.	Average Per Cent. of all Weekly Card-reports Stating the Pres- ence of Typhoid Fever.	More (+) or Less (-) Sickness from Typhoid Fever than the Seven Years' Average.
Av. 7 years, 1885-91 . .	31'06	29'45	293	31	=	9	=
1885	35'82	34'51	284	40	+ 9	8	- 1
1886	32'16	29'52	281	42	+ 11	8	- 1
1887	29'82	30'08	290	34	+ 3	10	+ 1
1888	29'55	25'76	294	29	- 2	10	+ 1
1889	28'18	23'28	304	19	- 12	10	+ 1
1890	30'20	33'95	300	28	- 3	8	- 1
1891	31'66	29'05	301	23	- 8	11	+ 2

We do not possess in New York so complete records as to the condition of the ground water as they have in Michigan ; but the rainfall, upon which ground water depends, is on record, and the reports show that more than the average amount of rain fell in New York during the autumn past, following, as it did, an exceedingly dry summer.

If typhoid fever bears relation to sudden rise in level of ground water, as has been held, rather than to the prolonged low state of such level, as is taught in Michigan, then surely the autumn of 1894 was a favorable time for a marked outbreak of the disease in the State of New York, but no such condition is reported by the sanitary authorities.

Just how the year of 1894 compared, in the matters of typhoid and rainfall, with 1893 and 1891, may be seen from the accompanying charts.

The above years were chosen for comparison because the summer of 1893 was very wet, and because the entire year of 1891 was especially noted for prevalence of typhoid fever.

These curves are very irregular, and suggest in places relation between low rainfall and prevalence of the disease; but the remarkable concordance exhibited in the Michigan chart is here sought in vain.

New York does not stand alone in its failure to accord with the Michigan rule, as is seen from the following Connecticut statistics :

	1894.		1893.		NORMAL.	
	Rainfall in Inches.	Deaths Typhoid.	Rainfall in Inches.	Deaths Typhoid.	Rainfall in Inches.	Deaths Typhoid.
July	2'40	15	1'89	18	4'99	14'8
August	1'70	38	4'86	14	5'17	32'8
September	4'63	38	2'24	37	3'76	41'4
October	6'11	32	4'75	49	3'90	42'6
November	4'23	37	2'56	35	3'90	40'2

As elsewhere, so in Connecticut, the summer of 1894 was very dry, and very heavy rains fell in the autumn, yet the autumn death rate was below the normal. If any weight is to be attached to the *sudden rise* in ground water, surely here was an opportunity for its exhibition; but the rise in water level was followed by no increase in typhoid.

Minnesota was also an apparent exception to the Michigan rule, as we see from the following :

	1894.		NORMAL.	
	Rainfall in Inches.	Deaths Typhoid.	Rainfall in Inches.	Deaths Typhoid.
August	1'22	39	2'80	43
September	2'04	42	2'04	67
October	3'37	47	1'55	87
November	0'54	30	0'69	62

This State presents a refutation of the "ground air" theory, for there was certainly a "sudden rise" in level of the ground water, without corresponding increase of typhoid; but there is no real exception here to the "Michigan rule," for it will be remembered that the said rule calls for marked lowness of ground water, and we notice that the autumn rainfall was above the normal in this State.

Among the other States heard from, the majority unquestionably fall under the Michigan rule. Definite information regarding mortuary statistics, was, in the cases of many States, impossible to secure, and only such expressions as "considerable typhoid," "largely increased typhoid" were obtainable. No statistics whatever are kept in certain States, and from them no results could be recorded.

The health department of Iowa writes, under date of December 17, 1894:

"There is greatly increased typhoid fever in this State. Scarcely a town or township is exempt."

The Iowa rainfall was:

	1894.	Normal.
August	1'58	3'60
September	3'57	3'70
October	2'67	2'85

Here again is noticed the probable low condition of ground water, and the application of the Michigan rule.

The Board of Health of Ohio writes: "We have noticed a very decided increase of typhoid fever in our State during the past autumn. It has also been noticeable that the increase has been almost wholly in our small villages, where wells are used for water supply."

The Ohio rainfall was:

	1894.	Normal.
August	1'67	3'04
September	3'31	2'94
October	2'01	2'57
November	2'17	2'99

This was also an instance of prolonged low water, with results following the rule.

In Pennsylvania, the records show large amount of typhoid present during the past autumn, with rainfall as follows:

	1894.	Normal.
August	1'84	4'90
September	6'30	3'72
October	4'26	3'48
November	2'50	3'33

A word is necessary here regarding the great precipitation for September. It will be remembered that, on the eighth of that month, an exceedingly violent storm swept over the eastern coast of the United States, with very heavy rainfall.

A great portion of this rain found its way directly to the streams and water-ways, and the ground water received but little reinforcement.

In Maryland "there has been a decided increase in the number of typhoid fever cases during the past six months (*i. e.* June to December) in all parts of the State. The rate of mortality has been low."

The rainfall was:

	1894.	Normal.
August	1'55	4'76
September	2'45	3'87
October	3'17	3'76
November	3'65	2'78

Low condition of ground water is here very apparent, and the State falls under the rule.

In Wisconsin "there is thought to have been a slight increase of typhoid in the State during the past autumn but not very marked."

The Wisconsin rainfall was :

	1894.	1893.
August	0'78	2'03
September	3'58	2'32
October	3'04	2'49
November	1'93	1'33

Comparison with the normal for this State is not possible from the data in my possession. It will be noticed, however, that the autumn was probably wet, and the typhoid was practically normal.

I do not possess the normal values for Massachusetts, but a comparison of 1894 with the previous year stands as follows :

	1894.		1893.	
	<i>Typhoid.</i>	<i>Rain.</i>	<i>Typhoid.</i>	<i>Rain.</i>
August	27	1'75	40	5'22
September	83	3'46	60	2'38
October	60	4'96	116	4'01
November	58	3'36	104	2'17

Evidently, Massachusetts is not to be rated as in accord with the Michigan rule.

In Indiana, so far as the Board of Health has been advised, "there has been about the usual amount of typhoid fever, following dry weather, more than there is after a season of plentiful rainfall."

The Western States, so far as heard from, are thus seen to follow, as a class, what has been styled the "Michigan rule," with the apparent exception of Minnesota; but it has been shown that this is really no exception at all in view of the more than normal rainfall. Now, what reasonable explanation can be given for the failure of New York, Connecticut and Massachusetts to accord with the rule?

While not wishing to dogmatize upon manifestly scanty data, the suggestion is offered that, so far as these three States are concerned, larger shares of their populations derive their drinking water from more or less carefully selected sources of public supply, and are consequently less exposed to danger from the local contamination of private wells.

Whether the exhaustive study of facts does, or does not,

FORTNIGHT
ENDING

LOW WATER IN WELLS AND TYPHOID FEVER—MICHIGAN.

VARIATIONS FROM NORMAL—STATE OF NEW YORK.

support the view that the relation of typhoid fever and rainfall, so far as ground water is concerned, deals with the question of low ground water, rather than with fluctuations in its vertical height, it admits of ready illustration that marked relationship certainly exists between this disease and the sudden influx of storm waters, which flood the polluted channels of some smaller rivers.

A recent publication of the London Local Government Board deals with this question most thoroughly, in an inquiry concerning a typhoid fever outbreak occurring in the valley of the river Tees.

The Tees is a small stream of northern England, about seventy miles long, and navigable for about four miles from its mouth.

Many important towns upon its banks pump from this river, and a goodly number of the inhabitants of the sanitary district use its waters for domestic purposes.

In many places, especially in the towns, it receives all sorts of polluting additions, which are carried on by the current to intakes below. During dry weather, the stream recedes considerably, leaving uncovered its rocky foreshores (shown in the accompanying views), which receive and accumulate filth of every variety, and retain the same until, by reason of heavy rain, the river suddenly rises and sweeps the refuse downward towards the towns nearer the sea.

The result produced upon the thoughtless public, of such an extra and concentrated dose of sewage material added to their water supply, is best shown graphically by the accompanying chart, where it will be observed that increase of rainfall is followed by increase in cases of typhoid fever, among the 219,435 persons using the Tees water, after an interval corresponding to the incubation period of the disease, while no appreciable result is noticed among 284,181 people of the same district, using other sources of supply.

The "typhoid rates" given in the chart are "cases," not "deaths."