these methods may be considered out of date, but until then something of the sort spoken of seems to be the only directly effective means at our command.

FAT EMBOLISM.¹

BY W. F. WHITNEY, M.D.

THE following case of comminuted fracture of the femur, with death on the fourth day, for which no evident cause was found, aside from a fat embolism of the lungs, leads to a consideration of how far we are justified in regarding this as an adequate cause of a fatal termination.

A male, sixty-two years old, entered the Massachusetts General Hospital in the service of Dr. Maurice H. Richardson, with a fracture of the right femur, from an elevator falling upon him two hours before.

During the first two days he was reported as comfortable, but having had no movement of the bowels was given a saline cathartic on the second day, without any effect.

Up to noon of the third day he had no defecation, and at that time his bowels began to swell, and increased till four p. m., with diminution of the area of liver dulness. An enema caused a defecation of the contents of the rectum. At one p. m., vomiting commenced and continued interruptedly till eight p. m. Vomitus was very dark brown, but without faecal odor. No gas passed either way. Urine normal and passed without difficulty. On the fourth day there were the same general symptoms. Vomitus of a black color. Abdomen aspirated and a clear straw-colored serum obtained.

Conscious, but restless. Suffered no pain, his only complaint was of difficulty in breathing. Pulse went up during the day while the temperature fell. Died suddenly at quarter-past five p. m.

	Aug. 31.	Sept. 1.	Sept. 2.	Sept. 3.	Sept. 4.
Temperature . . .	99°	101.5°	100°	101°	98°
Pulse	60	70	70	90	130

Autopsy by Dr. W. F. Whitney, eighteen hours after death. Body of a large man. Rigor mortis present, and some lividity of the dependent portions of the body. The right foot was everted, the leg shortened, and there were numerous ecchymoses over the ankle and middle of the thigh. Upon opening the latter a comminuted fracture of the femur in its middle third with considerable laceration of the muscles was found.

Head not opened.

Thorax, pericardium and heart were normal.

Both pleural cavities contained a little clear fluid with some old adhesions of the surface of the right lung.

The lungs were partly retracted, the upper lobes dry, the lower lobes of a dark bluish color, and considerable dark thin fluid escaped from the cut surface. There was no evidence of consolidation anywhere. Microscopic examinations showed the small vessels extensively plugged with a highly refracting fluid substance (fat).

Abdomen. The peritoneal cavity contained a small amount of fluid. The peritoneum was injected, but had everywhere its normal lustre, nor was there any-

where evidence of a deposit or exudation of any sort upon the surface. The stomach was filled with a dark, foully-smelling fluid, and this could be traced to the end of the jejunum. Beyond this point the intestines contained a little yellowish faeces. The inner surface of the stomach was marked here and there by a network of injected capillaries and a few superficial losses of substance with a hæmorrhagic infiltrated edge.

The intestines from the stomach to the rectum were extremely distended with gas.

The spleen was of normal size and pale. The kidneys and liver were pale, slightly opaque, and upon microscopic examination the cells were found to be markedly granular.

Before considering this case further it will be well to recall, briefly, what has been brought out from the study of the subject.

Magendie, in 1827, was the first to prove experimentally that oil injected into the veins of an animal would produce death. But it was not till 1862 that Zenker called attention to the presence of fat in the vessels of the lungs in some cases of death after fracture of the leg; and it then came to be regarded as a sufficient cause. But gradually, as the infectious process with which it was often associated were better understood, the conviction grew that fat embolism of the lungs alone was not sufficient to produce death, and that its action on centres more remote must be considered. By far the most thorough and exhaustive study was made by Scriba,² and published in 1879. And since then but little has been added to our knowledge. His work consisted in experiments on animals and the analysis of cases by the light of the results thus obtained.

The problems which he proposed were:

(1) How much fat is required to kill? (2) In what way does it act? (3) How is it eliminated?

In answer to the first he found that in order to kill an animal by fat embolism a quantity is necessary varying with the force with which it is injected, and with the distance of the point of injection from the heart. But in no case was it less than three times the amount of fat contained in the entire femur of the animal.

His conclusion as to the mode of action is: that death after aseptic fat emboli occurs only from large numbers in the brain and spinal cord, and in consequence of the arterial anæmia and the resulting disturbance of nutrition.

Of course, any deleterious action on the heart and lungs had first to be eliminated. In regard to the former it was done by direct observation on the living animal, and a comparison of the length of time that the heart of an animal killed by fat injection reacted to stimulus when compared with the normal. From these experiments he found that the disturbances which are caused by the entrance of oil into the heart are only to be found in the increase of the quantity of fluid of which the heart has to take care. And the speedy death is caused neither wholly nor in part by paralysis of the heart, and that aseptic fat has no permanent influence on its action.

As for the lungs, experiments have shown that five-sixths of the arteries must be tied before an animal is killed, and that at least three-quarters of the entire pulmonary system shut out before the arterial tension in the body is the least altered. Therefore he reasons that in man there is never such a degree of embolism

¹ Read before the Boston Society for Medical Observation, December 7, 1891.

² *Deutsch Zeitsft. f. Chirurgie*, Bd. 12, s. 118.

of the lung capillaries that death must result; and he was never able to produce this in animals even by direct intravenous injection.

On the other hand, it is the fat which has passed the lungs and reached the central nervous system that is the cause of death.

In answer to the third question, his experiments and clinical evidence (to be mentioned later) point to the kidneys as the organs by which the greater part is eliminated. A smaller part is undoubtedly burned up in the system.

Of the symptoms noted, one of the most striking was the constant fall in temperature. When large quantities of oil were used this fell 9.5°C . in nine hours in one case, and in another 7°C . in one and a half hours. In those cases in which death did not result, the return to normal was accomplished in about the same time as the fall.

Another symptom was dyspnoea, due to the loss of oxydation of the blood. This often was only transitory; but may be persistent and very urgent.

Slight hæmoptysis, due to infarction, has also been noticed. Edema of the lungs was never seen in any of his experiments although its occasional occurrence may not be denied.

The most important clinical symptom of fat embolism is the presence of oil in the urine. He examined this from a number of fracture patients, and was able in nearly all cases to demonstrate it to a greater or less degree. The oil did not appear in but a few cases as a continuous sheet on the surface, but as a whitish, slimy cloud in the upper part, at times three centimetres thick. It would not be recognized as fat by the eye alone; but with the microscope there were seen numerous very small, round drops, which were highly refracting, and had a fine dark border. At first, they might be regarded as bacteria, but the reaction with osmic acid and recrystallization after extracting with ether proved their true nature. A curious fact was ascertained, namely, that the elimination is not continuous, but interrupted. Fat was first noticed on the second, third and fourth days after the accident, then again from the tenth to the fourteenth day, and rarely once or twice again with intervals of from six to ten days between.

A marked pallor of the skin and mucous membranes was noticed in all the animals experimented upon.

In the fatal cases there was great irregularity of respiration, marked at the end by superficial respiration alternating with deep and spasmodic inspiratory movements.

These again point to anæmia of the brain, and death followed from paralysis of the respiratory centres, while the heart still continued to beat.

He last of all analyzed one hundred and seventy-seven reported cases, and found only fourteen in which death could be attributed to this cause alone, according to his standard.

The clinical picture which he draws, is as follows:

- (1) Certain proof of periodic occurrence of fat in the urine.
- (2) Transitory attacks of dyspnoea.
- (3) At times occurrence of slight hæmoptysis without fever and usually without dulness or râles.
- (4) Diminution of temperature.
- (5) Irregular action of the heart.
- (6) Collapse, with extreme pallor of the skin and mucous; at first, shallow respiration, at times broken

by a deep sighing inspiration, later, Cheyne-Stokes' phenomena.

(7) Spasms of different kinds or paralyses, which in the experiments were usually bilateral.

(8) Diminution of reflex irritability.

If the history of the case given at the beginning is now reviewed, it is hardly just with data given to ascribe the death to fat embolism alone.

The death within the first week, the dyspnoea, and the rapid fall in temperature in the last few hours are the clinical features which can be best brought in accordance with picture as drawn above. On the other hand, the fever during the first three days, the paralysis of the intestines with extreme gastric disturbance (although these two latter might be of central nervous origin) would rather point to a septic infection. This, however, had failed to localize itself either in the spleen or serous surfaces. The only positive evidence were the fat emboli in the lungs. It is now seen that the failure to examine the brain was a grave omission.

With the data at hand, the case is best regarded as one of septicæmia with fat embolism, in which the fatal result cannot be attributed to either alone but to both as factors.

The two important points brought out by the above consideration, are:

(1) On the clinical side, that the urine of patients with fracture should be carefully watched from the second to the fourth day for the presence of free or emulsified fat.

(2) On the pathological side, that unless found in the brain the cause of death cannot surely be ascribed to fat embolism.

OBSERVATIONS ON ALBUMINURIA IN SCARLATINA.

BY JAMES W. DUDLEY, M.D.,
Medical House-Officer, Boston City Hospital.

(Concluded from No. 6, page 140.)

TAKING the above standard as to the presence of nephritis there were found 18 cases out of the whole 100 which developed nephritis; that is, in 18% of the urines were albumen and casts found. Of these, eight were of the more severe type, and 10 the variety spoken of as catarrh.² One case only began as a catarrh and later developed into a severe nephritis. The catarrh was first noticed the third day of the disease, and on the eighth day, as desquamation was getting well-established, the albumen increased in amount, and epithelial and blood casts and free blood appeared. In this case the high temperature of the eruptive stage was prolonged evidently by the nephritis. This case is interesting as showing how the catarrh may terminate, a method of termination, however, which seems to be rare. The case is classified in the following statistics as one of severe nephritis.

Scarlatinal nephritis is said to occur most frequently between the ages of five and nine.³ The age of the patients having nephritis in this series of cases is shown

² As has been said it seems impossible to draw a definite line between mild and severe nephritis from the urinary examination, but for practical purposes it has seemed best to speak of such cases as presented a normal or high-colored urine with a trace of albumen and a few hyaline and granular casts only, as renal catarrh, and those which in any way gave evidence of more serious renal changes, such as blood or fatty elements, as the more severe form of nephritis. Only in one or two cases was there difficulty in deciding which variety the case should be called.

³ See Guy's Hospital Reports, vol. xxxi, 1889.