the source of vapour, and (2) at a height of three to four feet and three feet from the solution. Some threads were exposed freely to the vapour, and some were placed between folds of filter paper. Samples were removed at the end of two hours and again at the end of four hours. The results were not decisive, though showing decided effects of the vapour. There is no reason why a much larger quantity of antiseptic should not be used. The opening of the room at the end of two hours also considerably diminished the amount of vapour present.

Air examination.—The air examined before the action of the vapour yielded 429 organisms per ten litres, and after disinfection 71.

Micro-organisms.—The organisms employed were staphylococcus pyogenes aureus, bacillus typhosus, bacillus coli communis, bacillus pyocyaneus, and the spores of bacillus subtilis. All the organisms were affected—i.e., they produced more scanty cultures, or cultures of slower development than control threads. Those threads exposed on the floor were slightly more affected than those at a height of three or four feet, and there was a slight difference in favour of those protected by the filter paper. Of the individual organisms (1) staphylococcus pyogenes aureus was in no case sterilised, though the growths were less than the control; (2) bacillus typhosus: of 10 threads exposed 5 were sterile; (3) bacillus coli communis: 2 out of 10 threads were sterile; (4) bacillus pyocyaneus: delayed growths, but none were sterile; and (5) bacillus subtilis spores were in no case destroyed. A few experiments were made on the possibility of sterilising the skin, but in view of the difficulties noted by Lockwood they are too few to warrant any conclusions. The experiments were made on the cadaver, and the test used was the inoculation of a portion of the treated skin. Skin which was well washed with soap and water, then with 1 per cent. formaldehyde and left covered for five minutes with a cloth soaked in the solution, proved, after washing with ether to remove the antiseptic, to be in many cases sterile. From this point of view it has the practical advantage that it has little or no effect in retarding the formation of a lather with soap, comparing favourably with corrosive sublimate in this respect.

Formaldehyde in solution and as a vapour possesses decided antiseptic and disinfectant properties, and its non-poisonous character, easy vaporisation, and freedom from corrosive or other damaging action on fabrics will render it useful for many purposes of practical disinfection.

Northwick-terrace, St. John's Wood, N. W.

A Mirror

OF

HOSPITAL PRACTICE, BRITISH AND FOREIGN.

Nulla autem est alia pro certo noscendi via, nisi quamplurimas et morborum et dissectionum historias, tum aliorum tum proprias collectas habere, et inter se comparare.—MORGAGNI *De Sed. et Caus. Morb.*, lib. iv. Proemium.

GUY'S HOSPITAL.

SARCOMA IN THE MESO-SIGMOID; RESECTION OF SIGMOID WITH GROWTH; USE OF MURPHY'S BUTTON; RECOVERY.

(Under the care of Mr. W. ARBUTHNOT LANE.)

THE presence of a tumour in the meso-sigmoid as an isolated growth not of secondary character is an occurrence of great rarity. There are few instances published of removal of growths thus situated, involving as it does excision of bowel also. Dr. Duret¹ removed a large fibro-lipoma of the descending mesocolon. There was also marked prolapse of the uterus, and operation was advised, the diagnosis being a solid tumour of the adenoid variety, probably originating from the left ovary. Laparotomy revealed a solid tumour of the mesocolon; twelve centimetres of the large intestine were resected in removal of the growth. Death followed from shock. The methods of resection of the intestine before the

profession are numerous, and it is likely to be some time before surgeons are agreed as to the best of them. The fact that new methods are being constantly suggested implies that the profession is not satisfied with those at present in use. Another point to be noticed in the case, besides the method of resection, is the employment of the gauze as an additional security against the spread of septic fluid into the peritoneum, which might occur in consequence of a possible leakage.

A woman forty-three years of age was admitted into Guy's Hospital under the care of Mr. Lane on Jan. 13th, 1894, having been sent to the hospital by Dr. Gibbon of Wisbech, under whose observation she had recently come. About four years previously the patient noticed for the first time that she had a lump on the left side of the lower part of the abdomen. Since that time it had increased steadily in size till it was as large as two closed fists. It caused her great pain by its very free mobility. During the last six months she had lost flesh rapidly and had been obliged to take a large amount of purgative medicine to keep her bowels opened. Unless this was done she suffered much from distension. On admission she was found to have a well-defined tumour, which was globular in shape and about six or seven inches in diameter. When she occupied the supine posture the tumour lay in the upper part of the left iliac fossa, but could readily be displaced in any direction for six or eight inches. The abdomen was full and flaccid, and it was obvious that she had at one time been very stout. The symptoms she complained of were the obstinate constipation and the pain, tenderness and dragging produced by the tumour. It was apparent, therefore, that the mass was a growth which had recently begun to encroach on the lumen of the bowel, and, probably from its position and mobility, the part involved was the sigmoid flexure about its centre. On Jan. 15th, an incision being made over the growth, which was then readily exposed, it was found to be a smooth tumour, globular in form and but slightly irregular, and situated in the meso-sigmoid. It extended outwards to the bowel, which was stretched over it and apparently involved by it for a length of ten or twelve inches. With some difficulty the tumour was removed by ligaturing the mesentery about it, and it, together with the portion of sigmoid involved, was cut away. Messrs. Down Bros. had made for Mr. Lane some buttons after the patterns designed by Dr. Murphy of Chicago, and as Mr. Lane was anxious to give them a trial he proceeded to use them in this case. The particular difficulty which he experienced in the use of the buttons was that the sigmoid was loaded with fat and the appendices epiploicæ were very large and bulky. He had, therefore, some difficulty in puckering up the peritoneal coat, especially of the proximal segment of the bowel, around the tube of the button. If the button had been much larger this difficulty could have been met much better, but it would also have afforded an obstacle to the perfect coaptation of the divided ends by any form of suture or appliance. Feeling that the presence of such an abundant quantity of fat exposed the patient to the risk of imperfect union, he wrapped the buttoned intestine in gauze and replaced it in the peritoneal cavity.

On Jan. 20th, five days after the operation, the gauze-tampon was found to be moistened with pus, which had a faecal odour. The gauze was removed and fresh material introduced. On the 27th the button was passed with a natural motion. Some fluid faeces were present in the wound. The wound gradually closed up, and, being homesick, the patient insisted on leaving the hospital on Feb. 28th, with a small fistulous tract which discharged a little pus. There had been no evidence of any faecal matter in the discharge for a considerable time, so that the sinus will probably close very shortly. She gained flesh and strength very rapidly.

Remarks by Mr. LANE.—I was most pleased with the buttons, and hope that the next bowel on which I shall have an opportunity of trying them will contain less fat than this one did. Whether in all operations for establishing continuity of bowel it is wise to secure the patient from the possible yielding at the line of approximation, and consequent death from peritonitis, by the use of the gauze tampon is a matter of opinion; personally I use it freely in these and similar cases. At the same time it seems to me that the too early removal of the tampon may, by suddenly relieving the recently united bowel from external pressure, result in its yielding to the tension exerted by its contents. Apart from this, is it possible that matting of intestines around the tamponed area may, at a subsequent period, produce risks of intestinal obstruction? In other words, does this form of treatment,

¹ Journal des Sciences Médicales de Lille, 1892. Sajous, vol. iii, 1893, c. 60.

which secures the patient from immediate risk, expose him to another at a later period? On this point further information is requisite.

HOSPITAL FOR WOMEN, NOTTINGHAM.

A SUCCESSFUL CASE OF PORRO'S OPERATION.

(Under the care of Mr. H. R. HATHERLY.)

THE following very successful case of the advantages of the method of operation known as Porro's will be read with interest. By this method, as we need scarcely remind our readers, the child is delivered by the Cæsarean section, and then the uterus and its appendages are removed, the cervical portion only being left, as described below. Dr. R. P. Harris¹ gives in a paper some statistics of this operation which are of importance, and show its increasing success in skilled hands. The records for four years are as follow: 1885—24 cases, 7 deaths; 1886—23 cases, 4 deaths; 1887—18 cases, 5 deaths, 1 suicide; 1888—28 cases, 4 deaths. Of the first 50 women out of the 264 cases recorded, 30 died as the result of the operation, and of the last 50 but 10—a reduction of the death-rate from 60 per cent. down to 20. Of the whole 264 women, 117 died from the operation, divided as follows: 74 out of the first half—that is, 132—and 43 out of the second 132. Thus it will be seen, under several forms of calculation, that Porro's operation of to-day, under proper management, is far less fatal than that of five years ago—in 1884, for example, when 17 women died out of 29, or nearly 58 per cent.

A woman twenty-three years of age was admitted into the Hospital for Women, Nottingham, on Jan. 14th of this year, being in labour at the full term of pregnancy. She was the wife of a dwarf, and had been exhibited in various towns with her husband, the two styling themselves "The Midgets." The husband, twenty-eight years of age, was 3 ft. 10 in. in height and weighed 7½ st. He married on Oct. 21st, 1891. His wife was 3 ft. 8 in. in height and weighed before her pregnancy 6½ st. She had suffered from rickets in early life, and when seventeen years old had been admitted into the Leeds Infirmary, where (judging from her statement) osteotomy was performed. She appeared to have made a good recovery from the operation. There was nothing unusual in her menstrual history, and she had no recollection of ever having had any illness, or of having required medical attendance except when at the Leeds Infirmary. Her parents are stated to have been healthy people of average height and weight. On admission she was found to have rickety curves of the femora and spine, and a much deformed pelvis, the antero-posterior diameter of which was less than an inch and a half. The cervix was slightly dilated, and the vertex could be distinguished. As pains were infrequent and slight it was decided to defer any operative measures until the following morning so as to obtain the advantage of daylight.

Jan. 15th.—At intervals during the night the patient had had pains. The position of the head is unaltered. A consultation was held by the staff, assisted by Dr. Roberts and Mr. Paul, and the conclusion arrived at was that nothing could be gained by delay, and that Porro's operation was preferable to any attempt at craniotomy, the difficulties of which would have been enormous, or to Cæsarean section pure and simple. Chloroform was administered by Mr. E. W. Paul, the patient proving a good subject, and no anxiety was experienced throughout the operation on account of the anæsthetic. A median incision was made about seven inches in length; the abdomen was so distended that the skin appeared to be stretched tensely over the uterus and its contents. It was thought better under these circumstances to make a free incision, relying upon the contraction which would follow. At this stage of the operation hæmorrhage was very trifling and no ligature was required. On incising the uterus there was an alarming gush of blood, the placenta lying immediately under the incision. The child was removed with all speed and the placenta extracted, after which the immediate and rapid contraction of the uterus controlled the hæmorrhage almost entirely. The uterus being fairly contracted was withdrawn from the pelvis, transfixed and ligatured with stout silk in two parts immediately above the cervix. A clamp was then applied, and the body of the uterus, together with the appendages, was cut off as close to the clamp as practicable. The clamp being carefully secured and fixed externally the

abdominal edges were brought together by interrupted sutures, the stump was freely dusted with iodoform, the wound was simply covered with iodoform wool, a firm bandage was applied, and the patient was lifted from the operating table on to her bed.

On the evening of the operation the temperature rose to 101.6° F. and the pulse to 110; she had some sleep and a little pain, but not of a severe character. The wound healed rapidly by first intention, except in the immediate neighbourhood of the clamp, where a little suppuration occurred. Recovery, with some unimportant fluctuations, was uninterrupted. A peculiar vesicular eruption occurred about a week after the operation all over the lower part of the abdomen, the vulva, and the inner sides of the thighs. This eruption speedily disappeared after the substitution of dry boracic wool for the iodoform dressing. The clamp did not separate until the sixteenth day after the operation, when the stump was found to be firmly incorporated with the surrounding tissues. The child, which is still living, was 19 in. long at birth and weighed 6¼ lb. It was a female child, the head and trunk being well formed, but the legs and arms seemed to be short in proportion. The mother did not suckle her child, and, although there were a little fulness of the breasts and a slight secretion of milk, no trouble of any kind ensued. The patient remained in the hospital altogether six weeks, although she was able to be up and dressed after a month. She is now in good health, able to walk about with ease and comfort and to carry her own baby.

YORK COUNTY HOSPITAL.

LYMPHO-SARCOMA OF THE SMALL INTESTINE.—SARCOMA OF THE FEMUR.

(Under the care of Mr. JALLAND.)

LYMPHO-SARCOMA of the small intestine as a primary condition is extremely rare, and we must regard the growth in this instance as probably secondary to growth elsewhere. The second case is also an example of sarcomatous tumour. It is interesting as one of "acute traumatic malignancy," which was not submitted to operation. It was of extremely rapid course. A fortnight after a slight injury to the knee of a child in whose family there was a history of "internal cancer" a round-celled periosteal sarcoma developed and caused death through secondary affection of the lungs in less than three months. Operation in such cases may, perhaps, prolong life; it is not often that it cures the patient.

CASE 1. *Lympho-sarcoma of the small intestine.*—A boy aged six years was admitted to the York County Hospital on March 16th, 1893. He was supposed to have been kicked six weeks before admission and one week before the commencement of his illness. On admission the abdomen was found to be very enlarged, swollen, and tense, but not very tender to the touch. Down in the right iliac region a harder swelling was felt. The lungs were in a condition of broncho-pneumonia. On March 17th three separate lumps could be felt in both iliac regions. The urine contained a copious deposit of urates. The temperature ranged from normal to 100° F. On the 18th the child wandered considerably during the night. Perspiration was profuse, and he had offensive diarrhœa. On the 20th the lower half of the chest, one inch below the nipple on the right side, was dull on percussion. The breath sounds were almost absent, and the side did not expand on inspiration. The upper half of the abdomen was marked with large veins descending from the tissues over the lower half of the thorax. The lower half of the abdomen was singularly free from any trace of bloodvessels. The subcutaneous tissue was infiltrated with inflammatory material. The urine was scanty in amount, with a copious deposit of urates, and he had a good deal of pain on micturition. It was doubtful whether the tumours felt were inflammatory swellings or growths having some connexion with the bladder.

Mr. Jalland decided upon an exploratory operation. On opening the abdomen the peritoneum was found to be remarkably thickened. Serum, tending to be purulent, was allowed to escape and measured several ounces. In the iliac regions nodules could be felt, and as there was little doubt that the disease was malignant nothing more was attempted.

Post-mortem examination.—The omentum was immensely thickened, not with fat, but it was apparently hypertrophied. The appendices epiploicæ were universally enlarged. A large

¹ Sajous, vol. ii., 1891.