necessary to a determination of the mean position is still more scarce, and still more data must be obtained to form a mean value, although the pack-ice girdle thus obtained can only have an ideal importance. If we follow the boundaries of this pack-ice girdle, we notice to the south of New Zealand a bend towards the south analogous to that of the drift-ice, only shifted in accordance with the general direction of the current; also on the west side of Graham Land this tendency of the ice to bend towards the south is noticed, for it is always influenced by the direction of the land. To the south of Kerguelen Land the pack will only be to the west of Termination Land, and, indeed, stretching in a direction which makes us suspect another bend towards the south.

We must yet mention a sort of product of the drift, of considerable importance in ascertaining the condition of the drift and the ocean-currents, viz., the seaweed. This is loosened from its position especially by drifting icebergs, and turned adrift to the play of the waves. This conglomeration of plants, which reaches a little above the surface, though the long fibres stretch deep down into the water, shows the currents very correctly, for it is as little influenced by the winds as a drifting bottle. The curve that limits the place of growth is confined, in the Atlantic and Indian Oceans, to about the 60th meridian of E. long., near to that of the drift-ice. In some places this also holds good for regions to the south of Australia and the south-east of New Zealand. At all events, it can thus be understood how those places which are free from drift-ice are also free from drift-weed, a point of some importance when we consider the fact that drift-weed is the attendant of icebergs. According to the Dutch voyagers, on the west of the meridian of Paul and Amsterdam are found weeds different from those on the east thereof, which points to a different current.

(To be continued.)

TYNDALL'S RESEARCHES ON RADIANT HEAT

Contributions to Molecular Physics in the Domain of Radiant Heat. By John Tyndall, LL.D., F.R.S. (Longmans, 1872.)

FORTY years ago our knowledge concerning the radiation and absorption of heat was very meagre. We believe that the earliest experiment in this direction is to be found in that wonderful book containing the "Essays" of the Florentine Academicians. There we meet with the fact that the heat of the sun, converged by a mirror, can ignite a pastile placed within a Torricellian vacuum. A little later, in 1682, Mariotte communicated to the Paris Academy of Sciences that the heat of a common fire, made very sensible in the focus of a burning mirror, was entirely cut off by the interposition of a sheet of glass. The mirror in this case must have been of polished metal. These experiments were subsequently repeated and extended by Lambert, who, assigning the true cause, pointed out the necessity of employing metal, and not glass mirrors, in the reflection of heat from terrestrial sources. Lambert further showed that if the radiation from a clear fire were converged by a large glass lens, no heat was felt where the brilliant focus was seen,

whilst Hoffmann first collected the obscure heat of a stove to a focus by a metal mirror.

About a century ago Franklin made his experiments with bits of coloured cloth on snow, and found that, whereas colour strongly influenced the amount of solar heat absorbed, it made little or no difference on the emission from a lamp or candle. Beyond the foregoing information, we believe little more was known in this subject till, in 1777, Scheele published his famous treatise on "Air and Fire." But a great deal of ignorant talk, clothing itself beneath a barbarous jargon, was prevalent at that time. Putting much of this aside, Scheele asks himself the question, which he was unaware Lambert had already answered, whether it was really the heat, or only the light of a fire, that was reflected from a metal mirror; but first he seeks to know whether there are different kinds of heat. "There will be many," he remarks, "who will not hesitate to give an answer to my queries; for I am well acquainted with the vague phrases according to which everything is called Fire that hath a distant similarity to it. But I am of opinion," he naively adds, "it is best not to begin to read before one knows the elements of the alphabet, and to withhold an answer till one has reflected on the following experiments." Then comes a series of admirably simple experiments, pregnant with important results. As they deserve to be more widely known than they seem to be, the reader will pardon our quoting Scheele's own words at length. They are to be found in section 57 of his work already alluded to:—"From these experiments"—made in winter, before an open stove, and with only a candle, a concave mirror, and sheet of glass and metal, for apparatus—"it appears," says Scheele, "that the heat, mounting with the air in the stove, and passing through the chimney, is materially different from the heat passing through the door of the stove into the room. That the latter heat departs from the centre, where it is generated in straight lines, and is reflected from polished metals under an angle which is equal to that of incidence. That it unites not with air, nor can be diverted by a current of air, into a direction different from that which it originally had received. For that very reason, the vapours of the breath are visible in this great heat; for since air and heat are really united during summer, and warm air can dissolve more water than cold air, it likewise hence appears that this kind of heat is not united with air, nor has this kind of heat probably rarefied the air; and of consequence it becomes evident why it causes no tremulous motion in sunshine. You may, by means of a glass mirror, separate the light from this heat, when the heat remains in the glass, and the radiant light yields no heat. Hence it follows that the heat, passing through the door of the stove, coincides, in some points, with the light, but is not yet quite become light, since it is not reflected in the same manner from a glass surface as from a metallic one—a remarkable circumstance. This heat is soon changed into the usual heat, whenever it unites with a body, which may be observed in the glass, and in the metallic concave mirror blackened by soot, and in more instances. Represent to yourself," he concludes, "a little hillock of burning coals; in this case the heat darting from this hillock all around is that which may be reflected by a metallic polished plate; that,

on the contrary, which rises upwards, and may be driven by winds to and fro, unites with air. I will call the first kind, for distinction sake, *Radiant Heat*." Thus arose the term we still employ. The whole passage reads as if it were written almost at the present day; and the lucid style of the last sentence cannot fail to strike the reader. This is the more remarkable if we contrast it with the current ideas of the time, or even with Scheele's own description of the heat of contact; for a little further on he states, "This heat is a peculiar acid, which has admitted a certain quantity of phlogiston in its mixture."

Soon after this Pictet made his well-known experiments on the reflection of heat. In these he confirms the fact of the reflection and convergence of obscure rays, and discovers that the velocity of radiant heat is beyond the reach of experiment. To him is also due our first knowledge of the apparent reflection of cold, a fact explained some ten years later (1792) by Prevost, according to his famous theory of exchanges.

The experiments on radiation published in England at the close of the last and the early part of the present century, will be familiar to most of our readers. It will be remembered that Sir William Herschel established the refraction of heat, and the difference in the quality of solar and terrestrial heat; that he confirmed Leslie's experiments on the heating power of different parts of the solar spectrum, and first discovered that the maximum heat was beyond the visible red (experiments subsequently verified by Sir H. Englefield); that he also determined the transalency of various kinds and colours of glass, both to white light and to the light of the spectrum.

Both Rumford's and Leslie's inquiries into the Nature and Propagation of Heat quickly added to this knowledge; to Leslie belonging the capital discovery of the reciprocity of radiation and absorption. The accounts in our modern text-books render a further allusion to these experiments unnecessary.

A quarter of a century now lapsed; the attention of the scientific world being diverted by the electro-chemical discoveries of that period. One of the products of the new activity thus aroused was the discovery of thermo-electricity by Prof. Seebeck in 1822. Some ten years afterwards, Nobili constructed the well-known thermo-electric pile. Associating this instrument with a galvanometer, Melloni at once turned Seebeck's discovery into a thermoscope of surpassing delicacy. The fruit of one man's work thus soon became the seed of new and more vigorous investigation. And so prolific was this seed in Melloni's hands, that the blackened face of a thermo-pile is at present considered the indispensable pre-requisite in every exploration in "the domain of radiant heat." For six years Melloni pushed on with his researches; determining the amount of heat transmitted through innumerable solids and liquids—their relative *diathermancy*, as he expressed it—and using these determinations to investigate the quality of heat emitted from various sources. But the discovery with which his name will always be associated is that each material possesses a selective absorption, a veritable heat-tint, to which he gave the name of *thermochrosis*; thus confirming and explaining a similar fact previously noticed by De La Roche. Hence it was that Melloni called the volume "*La Thermochrose*," in which he grouped together the investigations that he

had published, in the "*Annales de Chimie*," and the *Comptes Rendus*, between the years 1833 and 1839. The appearance of these researches was characterised by M. Biot as "un nouveau champ de découvertes, que M. Melloni a exploité avec un sagacité une adresse et une patience inimaginables;" the subsequent verdict of physicists has not lessened this high opinion.

The interest awakened by Melloni's inquiries was no doubt the main cause of the rapid additions to our knowledge of the phenomena of radiation and absorption, that followed. Among others, Forbes, Dulong and Petit, De la Provostaye and Desains, Knoblauch, Jamin, Masson and Courtépée, Müller and Balfour Stewart, signalled themselves by the value of their investigations in this department of natural knowledge. But the whole of these inquiries were directed to the behaviour of solids or liquids, or the analysis of radiation itself. The influence of gases and vapours on radiant heat was not entertained. Melloni, indeed, thought such attenuated bodies could not come within the reach of experiment; for he had ascertained that a column of air some 20 ft. long exerted practically no absorption on the radiation from his source. Pouillet and Forbes, however, showed that the heat of the solar rays are largely absorbed by our atmosphere; and Franz believed (though erroneously) that he discovered a considerable absorption of heat by the air contained in a tube only 3 ft. long.

Briefly speaking, this was the state of our knowledge in this branch of Physics when Dr. Tyndall approached the subject in 1859. After having wrought for twelve years, Prof. Tyndall has now gathered into the volume before us the important results his unremitting labour has won. A summary of these results must be left to another article.

W. F. BARRETT

LETTERS TO THE EDITOR

[The Editor does not hold himself responsible for opinions expressed by his correspondents. No notice is taken of anonymous communications.]

On the supposed new Marine Animal from Barraud's Inlet

As some interest seems to be excited as to the nature of the animal that forms the long calcareous axis that has been received from Barraud's Inlet, I forward you a copy of a short notice I read on the subject before the California Academy of Sciences, July 17, 1871. I also enclose a piece of the stem with some of the soft parts still adhering to it, as it is possible its examination by competent observers will determine if I correctly referred it to the sponges.

"An examination of the specimens received from Barraud's Inlet enables me to refer them to the Protozoa class, Spongiæ or sponges. Although apparently nothing but the calcareous stem has reached us, yet on some specimens I found one end of the stem covered with a horny substance, which, when moistened and examined under the microscope, presents the character of a true sponge, being formed of a tough sarcode arranged in the form of irregularly reticulated tubes, the sides of which are studded with minute pores. The arrangement of the sarcode round the axis is not circular, but has somewhat the appearance of a Maltese cross. The central axis is formed by calcified layers of a tough chitinous substance. In the specimens we have received the greatest thickness is about $\frac{1}{4}$ of an inch, but the longitudinal fissures found in many of the stems would indicate that they had shrunk. They contain about 80 per cent. of carbonate of lime. At each end the stem is tapered off. The top terminates in a fine hair-like prolongation of uncalcified chitinous substance. The lower part of the stem, which in our specimens is the only part covered with sarcode,