digestive organs, as various other visceral disorders, are very prejudicial to efficient vocalization.

It is most important, therefore, in order that one shall get the best and fullest service out of his voice and be less liable to develop strain even from ordinary use, that all the bodily functions be maintained as nearly possible at par, and that all things which tend to depress the general health be religiously abstained from.

A robust constitution and a generally vigorous state of health are essential factors in the success of the habitual voice user.

Briefly summarized the general principles of vocal strain prevention are:

1. To avoid excessive use, to implicitly observe the warnings of fatigue, and to give the necessary opportunity for repose and restoration of the functions.
2. To avoid wrong methods of using the voice, which can be generally recognized by the fact that they require cramped, unnatural positions of the throat and are quickly followed by fatigue.
3. To avoid using the voice when the throat is inflamed from a cold, and to have removed without delay any local troubles in the nose or throat which hinder effective vocalization.
4. To observe all general hygienic laws, and to seek to maintain a healthy condition of the organism.

5. To strive by properly directed exercise to strengthen the neuro-muscular mechanism of the voice, and to acquire by learning and by practice the art of getting the maximal effect with the least expenditure of energy.

"The Rochambeau," 815 Conn. Ave.

AUTHOR'S ABSTRACTS.

Eye, Ear, Nose and Throat.

Something New in Tonsil Surgery. By W. F. Cole, Waco, Texas. Southwest Journal of Medicine and Surgery, May, 1915, pp. 144-148.

We have a history of tonsil surgery for more than two thousand years, and we find that radical operations have been repeatedly abandoned because found too dangerous to life. I know of no operation in surgery where the technique varied so little in all this time. I have always been conservative in my views, preferring safety, rather than to adopt the radical surgery advocated by so many. A school has arisen which contends that every vestige of the tonsil, with its so-called capsule, must be removed; and in order to accomplish this they have resorted to general anaesthesia and the use of knives and snares, resulting in many deaths. The operation, as done now by many surgeons, is more dangerous than appendectomy, but in the case of the tonsils it cannot be said, as in appendectomy, that it is necessary to save life, for hypertrophied tonsils rarely, if ever, cause death, even if left alone. I am as much in favor as any for the removal of hypertrophied or diseased tonsils, but I do not believe that we are justified in risking life that every bit of the tonsil and capsule shall be removed. I contend that a remnant of the tonsil or capsule left does no great harm. The complete extirpation of the tonsils has become a fad with many physicians, and the difficulties and dangers of the operation have taken it out of the hands of the general practitioner and given it into the sole care of the expert, with increased cost.