upon Neuroptera and Diptera. Among insects, the Lepidoptera were studied with especial care, the type selected being *Thyridopteryx ephemeriformis*. The careful account of this embryology, together with the numerous excellent figures, entirely substantiate the author's claim that the study of this type, "if it has brought to light nothing new, has, in the opinion of the writer at least, settled some important points connected with the embryology of this group of insects." In the account of segmentation and the formation of the blastoderm, the author completely confirms Bobretzky's descriptions. The development of *Thyridopteryx* occupies twelve quarto pages: for the details the original must be consulted. The account of the embryology of Orthoptera, represented by *Mantis* and the grasshopper, and of the embryology of spiders, is also very complete.

At the end of the paper many interesting and suggestive conclusions are appended. Among these it is significant that a writer who has done so much work upon the early stages of Limulus should unhesitatingly regard this latter form as an Arachnid. The Trilobites he considers as "possibly the ancestral form of Limulus."

Only a short account of this excellent paper has been given here. All those interested in embryology, and the light shed by it upon morphological science, will, of course, make a careful study of this work. E. B. P.

LETTERS TO THE EDITOR.

[The Editor does not hold himself responsible for opinions expressed by his correspondents. Neither can he undertake to return, or to correspond with the writers of, rejected manuscripts intended for this or any other part of NATURE. No notice is taken of anonymous communications.]

The Satellite of Procyon.

It is well-known that Procyon, like Sirius, does not travel through space in a straight line, its motion undergoing changes similar to those which would result from the disturbing action of a large satellite. This explanation was advanced by Bessel in 1844, and though the satellite has not yet been seen, its existence has been placed beyond reasonable doubt by Dr. Auwers's investigations on the subject.

Does it not seem probable that this interesting body may be revealed by the aid of photography? It is now possible to photograph stars and nebulae which are beyond the visual range of the most powerful telescopes; and if the companion of Procyon, like that of Sirius, be self-luminous, there would seem to be a good prospect of obtaining its image on the sensitive plate.

As the companion is doubtless in pretty close proximity to its brilliant primary, it will be necessary, for photographic purposes, to intercept the image of the latter by means of a suitable screen. Since the *direction* of the satellite at any time can be found from Dr. Auwers's elements,¹ there would be no uncertainty as to the position in which this eclipsing disk (or wire) ought to be placed, though its proper adjustment would be a somewhat delicate operation. Should the satellite be photographed, its position will become known from its configuration with respect to other stars recorded on the negative.

If a very large telescope were employed, the images of both components, as distinct and separate dots, might be obtained on the plate. For Procyon, a very short exposure would be requisite, and this could be secured by the use of a movable stop or screen, similar to that devised by Prof. Pritchard, and used for parallax work at the Oxford University Observatory.

It is scarcely necessary to dilate upon the interest which would attach to a photograph showing Procyon's companion. As the parallax of Procyon has been satisfactorily determined by Dr. Elkin and others—being 0".266 according to the Yale College observations—we could ascertain the *actual* as well as the *relative* masses of the two components. And the brightness, or more strictly speaking the photographic magnitude, of the satellite might also be determined with some precision.

¹ At present the position-angle of the satellite is about 233°. Its distance probably amounts to but a few seconds of arc, and may be within 2".

It may not be too late to obtain such results during the present season, but exposures of four or five hours, under good conditions, will not be practicable before next winter.

J. M. BARR.

St. Catharine's, Ontario, Canada, March 4.

"Les Tremblements de Terre."

THE issue of NATURE for February 7 (p. 337) contains a review of the little work on earthquakes published for me by Messrs. J. B. Baillière. The anonymous author of the article makes several criticisms on my book to which I desire to reply.

Your critic thinks it a grave fault not to have entered into a detailed description of the seismographs and seismometers at present in use. He reproaches me in particular for having but just mentioned Prof. Ewing's duplex pendulum seismograph; for having omitted to speak of the same inventor's horizontal pendulum seismograph; and especially for seeming to ignore the experiments made with these instruments by Prof. Ewing in Japan. I confess that I had not been struck by the excellence of the instruments in question, and that it was not through an oversight that I omitted to describe Prof. Ewing's observations in Japan, while I quoted in detail those of his *confrères*, Messrs. Milne and Gray, in the same country.

Your critic defends with some acerbity a certain class of seismographs, and wrongly accuses me of failing to appreciate the principles on which their construction is based. The objections which he makes to my treatment of M. Cavalleri's pendulums of unequal length are entirely refuted from a theoretical point of view by the learned note due to M. Poincaré, which is inserted on p. 46 of my book. I need not insist further on this point.

Your critic thinks I have not done sufficient justice to the work of the Italian savants: he forgets the limits necessarily imposed on a book destined especially to give to the general public an idea of the present state of an important question.

The writer of the article regards the seismographs of to-day as perfectly sufficient for all scientific needs. I am far from being the only person engaged in the study of earthquakes who does not share this opinion. Finally, he describes, and not very clearly, the experiments which I made with M. Michel Lévy to measure the rate of propagation of disturbances through the soil, and the registering apparatus designed for this purpose. According to the writer, these experiments constitute the only advances we have made in the study of earthquakes. If he had rendered justice to our work on the subject, we should have been content, and I should have raised no objections to his article. But your critic reproaches us with having given results which are masked by inevitable causes of error. A more careful study of the book, and especially of the extracts from our original memoir, published in the *Comptes rendus* for 1885 and 1886, would have preserved him from so inexact an assertion. In fact, he unjustifiably mixes up the *preliminary* experiments, made at Le Creusot by means of an apparatus exactly similar to that used by Mallet and Abbott,¹ with what really constitutes the basis of our work—I mean the determinations made with the aid of photographic registration and explosives. What your critic calls the "personal equation" of the instrument is here nearly negligible²; and the merit of our method lies precisely in this point, which distinguishes us from our predecessors.

We would beg the readers of NATURE to verify for themselves the truth of our statement. This verification will enable them to judge of the value of the article laid before them.

F. FOUQUE.

Finding Factors.

IT may add interest to Mr. Busk's ingenious method of distinguishing between prime and composite numbers to state the algebraic basis on which it rests.

Let N be any number, and n^2 the next higher square number, and let $N = n^2 - r_0 = (n+1)^2 - r_1 = (n+2)^2 - r_2 = \&c.$ r_1, r_2 are formed successively from r_0 by the successive additions of $2n+1, 2n+3, \dots$ the increments being in arithmetic progression, so that $r_m = r_0 + 2mn + m^2$. As soon as r_m becomes a square, N is expressed as the difference of two squares, and its factors are found.

¹ See p. 219 *et seq.*

² The only subsisting cause of error is that due to the inertia of the mercury, which we have estimated and introduced into all our calculations. (See p. 246).

If N is prime, it is expressible as the difference of two squares in only one way, viz. $\frac{1}{2}(N+1)^2 - \frac{1}{2}(N-1)^2$. To prove that N is prime by this method, the number of additions required is $\frac{1}{2}(N+1) - n$, which is $\frac{1}{2}(n-1)^2 - r_0$.

It may be noticed that when $n+m$ and r_m have a common measure, it must be a factor of N , and the additions need be continued no further.

$$\begin{array}{rcl} \text{For example,} & N = 8131, & n^2 = 8281. \\ & n = 91 & r_0 = 150 \\ & & 183 \\ & n + 1 = 92 & r_1 = 133 \\ & & 185 \\ & n + 2 = 93 & r_2 = 518 \\ & & 187 \\ & n + 3 = 94 & r_3 = 705 \end{array}$$

94 and 705 have a common measure, 47; therefore 8131 is divisible by 47, and the other factor is then found to be 173.

Mr. Busk's method of shortening, exemplified on p. 414 by his proof that $73 = 37^2 - 36^2$, depends upon the following:

Let $r_0 + 2mn + m^2 = (k+m)^2$, then $m = \frac{1}{2}(k^2 - r_0)/(n-k)$: since $k^2 - r_0$ is even, k is even or odd according as r_0 is even or odd; it is necessary only to try values of k descending by differences of 2; the greatest possible number of operations is $\frac{1}{2}(n-1-k_0)$, when k_0 is the value of k , with which we begin.

The process may conveniently be arranged as in the following example:

$$\begin{array}{ccccccc} \text{Let } N = 6667, & n^2 = 6724 = 82^2, & r_0 = 57. & & & & \\ \begin{array}{c} k \\ 15 \\ \\ 17 \\ \\ 19 \\ \\ 21 \\ \\ 23 \end{array} & \begin{array}{c} \dots \\ \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{array} & \begin{array}{c} 82-k \\ 67 \\ 65 \\ 63 \\ 61 \\ 59 \end{array} & \begin{array}{c} \dots \\ \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{array} & \begin{array}{c} \frac{1}{2}(k^2 - 57) \\ 84 \\ 116 \\ 152 \\ 192 \\ 236 \end{array} & \begin{array}{c} \dots \\ \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{array} & \begin{array}{c} \text{Quotient.} \\ \text{a fraction} \\ \\ \text{''} \\ \\ \text{''} \\ \\ \text{''} \\ \\ 4 \end{array} \end{array}$$

therefore $6667 = (82+4)^2 - (23+4)^2 = 113 \times 59$.

If N is composite, this method is not always shorter than the former. It will be shorter whenever $2m > k - k_0$, but it is not easy to see how to determine *a priori* whether this is the case.

The method by decreasing squares is not one of general application. For instance, the factors of 323,171 cannot so be found. It is the difference of two squares, each more than ten times as large as the first square used.

W. H. H. HUDSON.

King's College, London, March 15.

Dolomedes fimbriatus, Clerck, at Killarney.

It may interest some of your readers to know that this rare and fine aquatic spider occurs on Cromaglaun Mountain, near Killarney Lakes. I first found it when collecting the little shell, *Limnaeus involutus*, and though I had it two or three times in my hand, it was so active that it escaped, and I, not knowing its powers of diving, never thought of looking for it under water. The following year I again visited the little lake, which is called Crincaum, with some friends, and this time we fairly captured the spider, which I then easily identified as *Dolomedes fimbriatus*. There is a good account of it in Blackwall's "British Spiders," and also in Andrew Murray's "Economic Entomology—Aptera," but I am not aware that it had been observed in Ireland before I found it.

A. G. MORE.

March 18.

BEECH-WOOD.

IT is so characteristic of the science of to-day to find specialists narrowing their field of research, and confining their investigations to a deep narrow channel, that no surprise can be felt that two able men should devote

their energies for two years to the examination of the biology and chemistry of the wood of a single tree. It is not so easy to avoid astonishment at the results of the two years' work, however, appearing as they do in the form of a large book¹ of 238 pages of close description and argument, interspersed with long tables of figures, abounding in interesting information when properly read.

The authors have divided their work very fairly, the botanist having set himself the task of elucidating in detail the histology of the wood, the distribution of water, starch, and other contents, the formation of annual rings, and the growth in thickness of the trunk, and a number of other problems throwing light on the growth of the beech in the forest; while the chemist has confined himself to the task of analyzing the timber, so as to discover (1) the quantities of total ash, water, nitrogen, &c., in different parts of the tree; (2) the percentage composition of the ash, and the manner of distribution of the individual constituents; (3) the absolute quantities of each ash-constituent in 1000 parts, and other chosen quantities of dry substance of the wood; (4) the annual in-take and out-put of these constituents on a hectare of beech forest; and (5) similar particulars for the nitrogenous constituents.

The authors have by no means spared their trees. It is enough to make one envious to read of the trees cut down at all ages from 15 to 150 years, and of the specimens selected at all heights from each; how the research was extended to good, bad, and indifferent soils, and how trees in shade and in the open, trees entire and trees pruned, &c., were all laid under contribution as required. More than 100 stems of all ages were thus employed.

The manner of utilizing this enormous mass of material is worth noticing, for every kind of determination was made that would yield practical information.

The height of the trees was found, as the best indication of the value of the situation; the number of stems on a given area, their surface, contents, &c., were also determined; the age of the trees, their physiological condition, &c., were all considered in due course. The selected stems were then cut up as follows: transverse disks were cut at the successive heights of 1'3, 5'5, 10'7, 15'9, 21'1, and 26'3 metres, and separate determinations made of the specific gravity, histological peculiarities, analysis, &c., and these not only for wood and cortex separately, but also for each 30 annual rings of the stem. The thickness, density, &c., of the annual rings were also tabulated, and attention paid to north, south, east, and west sides of the stem.

Not only are all these data given in detail in the tables, but other tables are provided showing the mean densities, cubic contents, &c., &c., of whole trees, or of the trees on given areas; and the patient compilation and ingenious methods here displayed reflect the greatest credit on the authors. It is, in fact, especially in the application of their measurements, &c., to the forest as a whole that the tables will find their greatest practical value. There is also much of more abstract scientific interest to be learnt from the results.

On examining the histology of the wood, several new facts were discovered. The curious dipping in of the annual rings where they cross the broader medullary rays, and the deposits of grains of calcium carbonate on the septa of the vessels, may be mentioned by the way; but the most important results are those relating to the length of the elements, the lumina of the vessels, and the relative numbers and distribution of the latter on a square millimetre of transverse section.

The wood of the beech consists of the usual elements—vessels, tracheides, libriform fibres, and wood parenchyma, with transitional elements difficult to classify under any one of these heads. As was long ago pointed out by

¹ "Das Holz der Rothbuche," by Profs. R. Hartig and R. Weber (Berlin: Springer, 1888.)