Stopford, J. S. B., and E. D. Telford. COMPRESSION OF LOWER TRUNK OF BRACHIAL PLEXUS BY A FIRST DORSAL RIB. [British Journal of Surgery, Vol. VII, No. 26, 1919.]

A few cases of brachial compression neuritis by a normal first dorsal rib have been recorded previously, but an account is given here of ten cases observed within a period of two years proving the condition not so rare as formerly believed. The writer emphasises the fact that, even if an X-ray examination fails to reveal a supernumerary rib as cause of the compression, it must be remembered that a first dorsal rib can produce precisely similar manifestations. Probably a large proportion of obscure cases of atrophy of the hand are due to compression of the lower trunk by a first dorsal rib. All the patients suffered from neuralgic pain along the ulnar border of the forearm, which was induced or accentuated by anything producing depression of the shoulder girdle. Paresis and atrophy of the intrinsic muscles of the hand were noticeable about the same time as the pain, and, in some, the weakness affected the flexors and extensors of the wrist also. In 9 out of the 10 cases objective sensory disturbances were present, and in all nine the loss of protopathic sensibility was greater than the epicritic loss—a dissociation which the writer suggests as characteristic of nerve-compression. Trophic and vasomotor phenomena were very constantly present. In two patients the causation of the onset of the symptoms was clearly trauma, which had caused prolapse of the shoulder and stretching of the plexus over the rib. In other patients the onset occurs most frequently about puberty or early adult life the etiology probably being the same as in cervical rib. Three chief factors are suggested as contributing to the onset of clinical manifestations, and in most patients there is probably a combination of two or more:

1. A predisposing cause seems to be the anatomical relation of the nerve trunk and the rib; as when the bone is "bevelled" by a trunk which receives a large contribution from the upper two dorsal roots.
2. Excessive descent of the shoulder after birth.
3. Weakness or loss of tone in muscles which support the pectoral girdle; therefore all debilitating conditions may be the exciting cause.

In the majority, treatment is surgical, consisting of excision of that portion of the rib which is causing the compression. For the traumatic cases and a proportion of the early non-traumatic, development of the trapezius by faradic stimulation, exercises and massage are best. Patients were cured by this procedure. It is further recommended that similar methods should be employed as a routine after excision of the rib.

Where any underlying systemic condition is the cause of muscular loss of tone it is of fundamental importance to treat this. Early diagnosis is essential if a perfect recovery is to be expected. [Authors' abstract.]