CASE 23.—Gilford and Davis.²⁰ A man, aged 62, had a left-sided tumor of 6 months growth, the last two weeks' growth being very rapid. His appetite was poor and the man was wasting and weakening. There was no glandular involvement. He was operated on in 1904. The three carotid arteries were ligated. The vagus and descendens noni and the internal jugular vein were resected. Soon after recovery from the anesthetic, hemiplegia and double pneumonia ensued. Death occurred on the seventh day. No postmortem was allowed. The tumor surrounded the internal jugular vein and occluded it. The carotids were pressed on, but not occluded by the growth.

CASE 24.—Gilford and Davis.²⁰ A man, aged 52, had a left-sided tumor for 3 months, increasing slowly in size. There were no other symptoms; no enlarged glands. He was operated on in 1904. The tumor was dissected from the carotid arteries and 3 cm. of the internal jugular were removed. There was local recurrence six weeks later, which within two months had extended from the mastoid nearly to the clavicle. The patient died five months after the operation.

CASE 25.—Gilford and Davis.²⁰ A man, aged 74, had a left-sided tumor for over a year. The tumor was nodular and of stony hardness (Fig. 21). There were no enlarged glands, but the liver was enlarged to 3 cm. below the margin of the ribs. No operation was done, the patient died 18 months after the first appearance of the growth. At postmortem metastases were found in the liver.

CASE 26.—DaCosta.[†] A man, aged 52, had had a right-sided tumor the size of a pigeon's egg for 20 years. During the last six months it had begun to grow rapidly and was the size of a hen's egg when it was seen. It was firm but not hard. The anterior margin was lifted with each beat of the artery; the rest of the tumor was not lifted. The man was operated on in 1906. Very great hemorrhage occurred and about 50 ligatures were used. The three carotids and the internal jugular vein were tied. The nerves were uninjured. The tumor reached almost to the base of the skull. Twelve hours after operation, hemiplegia of the left side occurred which rapidly cleared up in a few days. A week later a second hemiplegia (embolic in internal capsule) occurred which is slowly clearing up (June, 1906). A few days after operation atelectasis of the left lung developed from unknown cause; over one-half of the lung collapsed. It slowly but finally expanded. Recovery ensued.

CASE 27.—Case reported in this paper by Keen, 1906.

CASES DISCOVERED POSTMORTEM.

In addition to the above-mentioned cases, the following were found at postmortems:

CASE 28.—Hedinger,²¹ in 1904, accidentally discovered a tumor on the right side in a woman of 62. Size of specimen, 3x1.5 x1.5 cm.

CASE 29.—Oberndorfer,²² in 1905, discovered a tumor on the right side in a woman of 58 who had suffered from severe progressive osteomalacia. The vessels were not involved in the tumor except by lying in grooves on its surface. Microscopic examination showed the tumor to be one of the carotid gland.

(To be concluded.)

CONGENITAL CYSTIC KIDNEY.*

Medical Association, at the Fifty-seventh Annual Session, June, 1906.

F. B. LUND, M.D.
BOSTON.

Congenital cystic disease of the kidney is a rare affection and one which so far appears to have interested the pathologists far more than clinical observers. The pathology of the disease presents an abundance of controversial literature, for the reason, perhaps, that the condition has been hard to explain and has afforded abundant

opportunity for discussion and theorizing, but we, as practitioners of surgery, must not forget that the disease is of eminent practical and surgical importance, and that, although we may not meet with it on the operating table many times in a long surgical career, it is well worth while, when those times come, to be prepared by a previous consideration of its clinical history, prognosis and surgical treatment.

It is a disease which frequently runs its course without a diagnosis being made during life, death occurring from uremia or cerebral hemorrhage, due to kidney insufficiency. The real cause of the insufficiency is first discovered at the autopsy, when the enlarged cystic kidneys, not felt during life, are found by the pathologist. In fact, the course of the disease may be so latent that a patient in whom pathologic changes have not been suspected may be stricken with uremia or cerebral hemorrhage.

Strange to say, these cases come more often during life under the observation of the surgeon and obstetrician than of the medical man, for the reason that the symptoms which attract notice are more frequently those due to the size of the tumor than to the renal insufficiency. As is well known, cystic kidneys in the newborn are frequently so enlarged that the abdomen of the child is enormously distended. Normal parturition becomes impossible, version is generally attempted, the limbs are pulled off and it is often necessary to open the abdomen of the fetus and remove the enormous kidneys before the body and head of the child can be delivered. With this class of cases, except to allude to their important bearing on the etiology and pathology of the disease, I do not intend to deal.

It is the cases in which gradual enlargement of a polycystic character takes place later in life, and which present themselves to the surgeon on account of dragging pain and the presence of a tumor in the loin, or possibly on account of mobility of the enlarged kidney, that I wish to consider. Such tumors may suggest malignant disease of the kidney or hydronephrosis, and it is frequently only when they are cut down on that their true nature is revealed, and the importance of a knowledge of their nature and treatment becomes evident. Such kidneys have not infrequently been removed, with the result that the patients have died of uremia, and the autopsy has revealed the presence of cystic disease of the other kidney, usually less advanced, so that a tumor was not felt before operation. A knowledge of the fact that this disease is usually bilateral might have saved the patient's life and the surgeon's discomfiture. Again, the superficial resemblance of polycystic kidneys to sarcoma has been an additional reason for the performance of nephrectomy, which has sometimes been fatal and sometimes not, the issue depending more on good fortune than surgical skill. A slight familiarity with the pathologic condition should ordinarily suffice to prevent such a mistake, and a brief consideration of the pathology and etiology becomes important.

ETIOLOGY AND PATHOLOGY.

The term congenital cystic kidney does not include hydronephrosis, which is a dilatation primarily of the pelvis and sinus of the kidney from retention due to obstruction in the pelvis itself or the ureter, and does not concern the urinary tubules at all. It does not include cystic dilatation of the tubules due to the contraction of the interstitial tissue, which sometimes accompanies a chronic interstitial nephritis. Here the kidney cortex becomes studded with little cysts, seldom larger than a

† To be reported in the *Annals of Surgery*, September, 1906.

20. Gilford and Davis: *The Practitioner*, December, 1904, p. 729.

21. Hedinger: *Corr-bl. f. Schweizer Aerzte*, No. 8, 1905.

22. Oberndorfer: *Centbl. f. allg. Path. u. path. Anat.*, 1905, vol. xvi, 1905.

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.

pea, and not, as in the true congenital cystic kidney, greatly enlarged and varying in size. The cyst in interstitial nephritis play only a subordinate rôle and are secondary to scar contraction. Neither does the subject under discussion include the single or occasionally multiple cortical cysts which sometimes result from the absorption of embolic infarcts. In the true cystic kidney the kidneys may be enormously enlarged, e. g., the cases in which they present an obstacle to parturition, or in adults occupy half or two-thirds of the abdominal cavity. The whole organ, both cortex and medulla, is studded with cysts of various sizes, from that of a pinhead to that of a mandarin orange (Fig. 1). The surface is

Fig. 1.—Cystic kidney removed at autopsy from a patient of Dr. H. T. Baldwin. The man died at 42 years, of uremia. Both kidneys were equally involved and caused a large abdominal tumor. The patient's mother suffered from the same condition.

knobby with cysts of different size and color, brown, red, green, translucent, like windows bossed with vari-colored glass, making a pathologic specimen of truly remarkable and imposing appearance. The walls of the cysts are thin, vascular, and the larger ones show folds on their inner surface, which are the remains of the walls of the smaller cysts, which have coalesced to form them. The fluid consists of urine, since in it are found urinary constituents, such as uric acid and hippuric acid, calcic oxalate, cystin, leucin and tyrosin, but in the larger cysts this fluid is considerably diluted (probably by serum which has transuded through the walls), albu-

minous, and contains blood, pigment, fat and cholesterol in varying quantities, so that the color of the cysts may be red, brown or yellow, and the fluid in them more or less cloudy. Between the cysts are often found more or less extensive areas of normal renal parenchyma; other cysts are separated merely by fibrous tissue. In some cases, with large amounts of parenchymal tissue, we can not distinguish between cortex and medulla.

The efferent passages, which protrude into the calices or pelvis, frequently show no changes, but in some specimens are shrunken, narrowed, and may be almost obliterated. Thin-walled cysts may sometimes be seen projecting into the pelvis, just as they do on the outer surface of the kidney. Microscopically, each cyst is lined with a single layer of flat epithelial cells, which in the smaller cysts correspond closely to a tunica propria. In the larger cysts this is not found, either because they have no proper wall or because the cysts appear as dilatations in the interstitial tissue. The partitions between the cysts are so thin as to make it probable that the larger cysts have been formed by the coalescence of smaller ones. In some of the smaller cysts glomeruli have been found attached to the walls, flattened, but still permeable to injected fluid (Figs. 2 and 3). It is thus evident that all portions of the tubular tract, from the glomeruli to the terminations in the papillæ, are subject to cystic dilatation. The suggestion of certain investigators that they might arise by cystic degeneration of the interstitial tissue is not supported by recorded observations. That certain of the cysts show no epithelial lining is due to the fact that the epithelium is desquamated and disappears. In the smaller or younger cysts the epithelium is always found.

Cystic kidneys in adults and in the new-born are usually bilateral. If, as is rarely the case, the disease is unilateral, the other kidney is hypertrophied. To the obstetric importance of this disease we have alluded above. Even earlier than this the degeneration of the kidney may have an unfavorable influence on the development of the fetus, so that birth is premature. The child may die within a few moments of birth from respiratory failure, due to the interference with the movements of the diaphragm consequent on the tumor, or may live a few days, only to succumb to the same cause. Death may result early in these cases from renal insufficiency, the amount of normal functioning renal tissue between the cysts being insufficient for excretory purposes. On the other hand, the amount of normal renal tissue may be sufficient to prolong life for indefinite periods, perhaps for decades, before renal insufficiency results in death. The most favorable conditions will naturally be presented by unilateral cases with compensatory hypertrophy of the opposite kidney, which are unfortunately rare.

In considering the etiology of this disease we enter a field which has been fruitful of much wordy discussion. The theory of Neuwerck, Hufschmidt and von Kahlden that these tumors are cyst-adenomata is negated, it seems to me, by the fact that the cysts may often be demonstrated to be dilatations of the renal tubules, and contain urine, often diluted with serum, it is true, but showing distinct urinary constituents. The epithelial proliferations and connective tissue growths, which Brigid and Severi had relied on as evidence of tumor growth or adenoma, have been shown by von Mutach to be tissues characteristic of embryonal kidneys. The presence of these proliferations, etc., therefore, is an argument in favor of the developmental rather than the neoplastic origin of the tumors. In order to explain these

cases we have to assume, with Orth and others, the existence of some obstruction to the escape of urine from the dilated and cystic canals. The oldest theory, that of Virchow, that the urinary concretions, "uric acid infarcts," so common in the new-born, have plugged the canals, is untenable, since these are rare during intra-uterine life, and after birth so common that cystic kidneys, if due to them, ought to be much more frequent than they are. There seems to be no doubt, and that Virchow himself acknowledged later, that we must assume an organic obstruction and actual obliteration of the tubules. Orth agrees with Virchow's later view that the condition results from a fetal inflammation, involving the papillæ, with resulting obliteration. He says that the papillary atresia can not be attributed to a congenital narrowing, because certain embryologic investigations have shown that the tubules are developed from the Wolffian ducts and do not develop separately from them and afterward unite with them. The fact that we find cysts in all parts of the kidney, including the papillæ, does not, however, make it necessary to assume that the obstruction must be located in the papillæ. The cysts of the upper tubules might extend into the papillary tract and displace the straight tubules and project into the pelvis.

In favor of the theory that the condition is a developmental malformation is the fact that other congenital malformations frequently coexist. The presence of congenital cystic liver in association with cystic kidney, which occurs in 19 per cent. of the cases (Moscowitz), is also significant of the embryonal origin of the condition. The cysts of the liver are ascribed by Moscowitz¹ as due to the obstruction of afferent intra-acinar bile ducts, and are evidence also of congenital origin. Other anomalies which have been found associated with cystic kidney are hypospadias, atresia and absence of ureter, vagina, bladder, etc., also hare-lip, spina bifida or absence of extremities. There is now pretty general agreement among investigators that the condition is due to defect in development, which is explained something as follows: The tubular structure of the kidney, including the glomeruli, the convoluted tubules, the loops of Henle and the straight tubules, to a considerable extent, originate by a development of tubules in the mesoderm of the embryo. The ureter, kidney, pelvis and an uncertain amount of the straight tubules in the pyramids develop from a tubular process which originates from the Wolffian duct. The straight tubules grow out and meet the primitive vesicle, which becomes the glomerulus, and then later from this glomerulus the convoluted tubules and Henle's loops are developed (Brödel). In case this tubule growing from the Wolffian duct fails to connect with the primitive glomerular vesicle, an obstruction occurs, resulting in cystic dilatation of the entire tubular system above the straight tubules. The extent of the process will depend on the relative number of tubules that fail to meet. If a large proportion fail to connect, then the cystic formation will be rapid and complete, and, no kidney tissue being left to eliminate urine, the patient will die. If, on the other hand, there be left comparatively large areas of healthy kidney tissue, enough for the elimination of uric acid, the patient may live for years and die from other causes. The presence of the increasing cystic tissue may finally so squeeze the intervening healthy tissue as to cause insufficiency, and the patient dies from this. This theory seems adequate to explain the facts as observed, and to be in every way more probable than that fetal nephritis involves the papillæ.

1. *Am. Jour. Med. Sci.*, April 1, 1906.

Bunting,² in a recent article based on a careful microscopic study of specimens from two cases, found that in these cases the urinary path could be opened from the glomeruli through to the pelvis of the kidney, and, therefore, added no evidence in support of the obstruction theory or a failure of two sets of tubules to join with each other. He believes the primary cause of the trouble is the epithelial hyperplasia, which, so to speak, forces out the walls of the tubules. This theory does not satisfactorily explain, however, why we get cystic forms instead of epithelial growths, and needs further confirmation. Whatever theory of cystic kidney be adopted, we can not help concluding that it is a developmental condition, a malformation rather than a new growth. A fact that speaks strongly for the malformation origin is the occasional occurrence in several generations of the same family, showing a tendency to inheritance, as in the case illustrated by Fig. 1.

Cystic kidney has oftenest been observed in the middle decades of life, but cases have been reported in the first and second decades as well. In the adult the pathology corresponds in every way with that of the new-

Fig. 2.—Case 3. Section of cystic kidney removed at operation, showing normal kidney tissue between the dilated or cystic tubules. In the upper part of the picture a flattened glomerulus can be seen.

born, except that the individual cysts are larger, as are also the entire kidneys. It becomes evident, therefore, that if our explanation is correct we must assume that in case of cystic kidney in the adult the condition has existed since birth, but has not until later years given rise to symptoms which attract attention.

SYMPTOMS AND DIAGNOSIS.

If our views of the etiology and pathology of the disease as given above are correct, they will furnish us an adequate explanation of the somewhat remarkable symptomatology and course of the disease. The most thorough and careful description of the course and symptoms which I have found is that of Milward of Birmingham, who divides the clinical history of the disease into three stages:

1. The stage of progressive enlargement of one or both

2. *Jour. of Exp. Med.*, March 26, 1906.

kidneys without subjective symptoms. The renal enlargement is discovered, if at all, by accident. This stage may last from a few months to several years.

2. The stage of subjective symptoms and objective signs. This stage lasts from a few months to six or eight years, or even longer. The signs and symptoms are dependent on the size and weight of the tumors.

3. The stage of decreasing elimination of urine. In this stage appear the symptoms of uremia, or cerebral complications. It lasts two or three weeks or a few months.

Males and females are affected in about equal numbers. The disease usually terminates fatally in the fourth, fifth or sixth decade. Renal enlargement is rarely discovered before adolescence, and when the disease develops in infancy the child hardly ever lives till the tenth year, so that the disease has been rarely reported between the tenth and thirtieth years.

The symptoms noted during the second stage are a dull, aching pain in the region of the kidneys during both night and day, but usually worse in the daytime, especially if the kidney is movable. A large part of the

usually slightly increased, but may be diminished. There is usually only a slight trace of albumin, and often albumin is entirely absent. In the second stage it is never present in large quantities, except when there is a great deal of blood. Casts are practically always absent; leucocytes are common and often occur in large numbers. Pus may be present in acid urine. Blood is usually present in traces, and when there is hemorrhage into the cysts may be poured out in such large quantities as to constitute attacks of hematuria. Bleeding may occur every few months or once a year, or may never be sufficient to tinge the urine. Some blood, however, is always present, and its presence is significant (Milward).

The elimination of urea may amount to from 250 to 350 grains until late in the second stage, and it is not infrequent when death occurs suddenly, as it often does in the third stage, for the physician after death to find, to his surprise, the large cystic kidneys.

The symptoms of the third stage are those of uremia or some allied complication, such as cerebral hemorrhage, due to arteriosclerosis, and consist of headache, dyspnea, drowsiness, convulsions, paralysis and coma.

Duration of the Disease.—Ritchie,³ reporting on eighty-eight cases, found that in seventy-five the average age was 45.3.

Duration of Symptoms.—In the majority of cases the symptoms last from a few months to five or six years. Twenty-two and nineteen years respectively was the duration of symptoms in the longest cases noted.

Of seventy-eight cases in which the symptoms were analyzed, thirty had renal symptoms apart from albuminuria, twenty having pains in the loins and abdomen. Sixteen had hematuria at some time. Seven had no kidney trouble and no evidence of illness, but cystic kidneys were found postmortem. Eleven suffered or had suffered from edema; in ten the absence of edema was stated. In thirty-three the urine contained albumin, in nineteen a trace and in eleven a considerable quantity. Vomiting or intestinal symptoms, constipation or diarrhea were present in nineteen cases. Twenty showed anemia, cachexia or loss of flesh. Thirteen had dyspnea or some pulmonary affection. In twenty-one a tumor was discovered during life in one or both flanks.

A brief review of the literature will show us that these cases which come under the surgeon's care present themselves during the second stage for pain in the side, due either to the weight or mobility of the tumor; the surgeon, finding the tumor, operates, and if it is large and malignant looking not infrequently removes it. The patient lives or dies, according to the presence or absence in the other kidney of sufficient healthy tissue to attend to the elimination of urea.

TREATMENT.

Numerous operators have removed congenital cystic kidneys successfully; among them may be mentioned the following:

Bevan⁴ performed a successful nephrectomy for hematuria and secondary anemia in a case of cystic kidney, with recovery.

Syms,⁴ operating for floating kidney, found a cystic kidney, which he removed and the patient recovered.

Krönlein⁵ removed a large cystic kidney in a patient three months pregnant, with recovery.

3. Laboratory Reports of the Colleges of Physicians of Edinburgh, 1892.

4. Annals of Surg., Phila., vol. xxxix, 467.

5. Beits. sur Klin. Chir., Tub., 1903-04, vol. xli, p. 170.

Fig. 3.—Cross section of the same specimen as Figure 2.

pain, especially in the later stages, seems to be due to mobility of the kidney. There is tenderness on palpation or pressure, and occasionally the pain may be paroxysmal and very severe, with resulting prostration. Since many of the kidneys examined postmortem or removed at operation present cysts filled with blood under high pressure, it is reasonable to suppose that some of the pain may be due to hemorrhages into the cysts. Occasionally suppuration has taken place in one or more of the cysts and has required operation. Flatulence, headache, dyspepsia, vomiting, anorexia and constipation are frequently concomitant symptoms. It is an important and somewhat interesting fact that the examination of the urine often throws very little light on the diagnosis, a circumstance which contributes greatly to the difficulty of the latter.

During the second stage, and until the deformation is far advanced, the kidneys are capable of continuing their functions almost perfectly. The specific gravity is usually low, below 110, and the amount passed is

Stander performed nephrectomy on a cystic kidney for pain and hemorrhage. Infection and abscess ensued, requiring removal of the kidney, with recovery of the patient.

Schmid, Niemeyer, Bardeleben, Hayems and others, however, have performed nephrectomy for polycystic kidneys, but the patients have died of uremia, and at autopsy the other kidney has been found the seat of rather extensive disease.

The conclusion becomes evident that while from hemorrhage or suppuration in a cystic kidney, or from symptoms due to the size of the tumor, nephrectomy may be indicated, it should never be performed until, by exploratory incision or ureteral catheterization, the other kidney has been found to be healthy. Especially important is it in operating on obscure kidney tumors to have in mind the possibility of congenital cystic kidney and be prepared to deal with the condition if found.

In certain rare instances not over 2 to 3 per cent. (Bevan) congenital cystic kidney is unilateral, and occasionally, as in one of the cases found at autopsy at the hospital, compensatory hypertrophy of the other kidney is found. In such a case as this, if the symptoms justified it, nephrectomy would be obviously proper.

The question now arises, What are we to do in case the mobility or enlargement of a kidney is making life unendurable and the condition of the other kidney precludes nephrectomy?

Under these conditions Curtis and Kammerer⁶ have practiced incision and breaking down of the walls of the larger cysts sufficiently to reduce the size of the kidney, and suture of the depleted organ to the lumbar muscles. In the two cases in which this was practiced the symptoms were relieved and the patients enabled to return to work. Curtis' patient was in good health and had gained in weight at the end of three years. The method of these surgeons consists in free incision of the diseased kidney and systematic opening of the larger cysts, which are packed with gauze to secure obliteration. By this⁶ the size of the organ and tension of the capsule are markedly diminished, relieving the compression of the more or less healthy tubules. The kidney is thoroughly anchored in the loin and symptoms due to mobility disappear. Permanent fistula does not ensue unless there is some stricture in the pelvis. Such a procedure may be applied first to one kidney and then to the other as it increases in size, thus preserving as long as possible the functions of the diseased organs and prolonging the life of the patient. This procedure is far safer than nephrectomy, and accomplishes the desired result of diminishing the size and mobility of the tumor and relieving pressure on the remaining tubules in the simplest and safest way. It seems to me the only rational surgical treatment and is therefore advised.

Without knowledge of Curtis' and Kammerer's work, a similar procedure was adopted in Case 1 of this paper, a report of which follows.

REPORT OF CASES.

In Cases 2 and 3, which were kindly placed at my disposal by Drs. J. B. Blake and G. H. Monks, the only other cases operated on which I have been able to find in the records of the City Hospital, nephrectomy was performed, as the size of the tumor was so great and the symptoms so severe that this procedure alone seemed to promise relief. Death resulted apparently from uremia.

CASE 1.—M. H., 26 years of age, single, a nurse, was admitted to the City Hospital in August, 1904. Two and one-half years ago she had felt a small tumor in the right upper abdomen, which had gradually grown larger and descended. At various intervals she had felt pain in the region of the tumor. The symptoms were much worse in the last eight months, and the tumor had fallen as low as pubes and been felt to go back on recumbency. She has suffered from headaches. There have been no urinary symptoms.

Examination.—Urine pale; specific gravity, 1018. Sediment normal, and abnormal blood, slight in amount. Numerous casts, hyaline and granular, with an occasional cell adherent. A few leucocytes. A few round cells. A mass the size of two fists, freely movable, was felt below the right costal margin. The left kidney was just palpable. There was no tenderness or spasm. A right lumbar incision showed a multilocular cystic kidney. The cysts were of various colors and the size from a hickory nut to an English walnut.

Operation.—The kidney was so greatly enlarged and there was so little renal tissue that a nephrectomy was considered, but before proceeding to do it it was thought wiser to ascertain the condition of the other kidney, which was accordingly done by a left lumbar incision. That kidney was also found to be somewhat enlarged and cystic, though not nearly so large as the right. After suture of the wound on the left, a number of the larger cysts in the right kidney were opened and the kidney sutured as well as might be to the lumbar muscles. During anesthesia and during the afternoon after the operation the patient was attacked with cyanosis and collapse, the symptoms requiring artificial respiration. These attacks occurred after the administration of one-sixth of a grain of morphin, and were apparently due to a particular susceptibility to that drug. Otherwise the convalescence was normal. The patient left the hospital in about six weeks, and up to January, 1905, was well, save for some gastric disturbance, and able to take care of a case. Seven weeks after the operation she says she was "really feeling well" and "fell the whole length of the schoolhouse stairs without pain," thus showing that the kidney was apparently fixed in place.

Subsequent History.—Nearly six months after the operation she was taken ill with pain in the side and tenderness over the kidney and a temperature rising to 101 in the latter part of the day. She had headaches, loss of appetite and bloody urine. This condition lasted for three weeks and then gradually wore away. She was well until early in October, fourteen months after the operation, when she had a week of weariness, with considerable sideache and headache. At this time she had a chill. A diarrhea accompanied this illness and the patient thinks it may not have been due to any trouble in the kidney. Ever since then up to the present time (May, 1906) she has been in good health, or, as she expresses it, "comfortably well." Walking more than a mile at a time always produces an ache in the side. Standing still for only a few moments will cause this ache. Standing hurts more than anything else. She has occasional dull headaches, which gradually wear away in the middle of the day, and every few weeks she has a severe headache. Before the operation she had complained of swelling of the ankles. This has not appeared since the operation. An examination of the urine, taken about three months after the operation, showed a trace of albumin, and another examination, about six months after the operation, showed also a trace of albumin, normal red-blood corpuscles and squamous cells, but no casts. During this week (May, 1906) a test for albumin, made by Dr. Thomas, showed no albumin at all.

About fifteen months after the operation the patient began to work steadily at her profession and has worked since not more than half the time, but doing any kind of work that offered. At no time has work seemed to cause any trouble. Her weight has been about ten to twelve pounds less than before the operation. The kidney seemed to the patient to feel smaller after the illness in January, 1905.

It will thus be seen that the operation, though I felt very doubtful of it at the time, succeeded in diminishing the size of the kidney and fixing it in position, so that the patient has since lived a useful and fairly active life. In another similar case I should incise a much larger

6. *Annals of Surgery*, 1901, vol. xxxiv, 419.

number of the cysts and perhaps pack them with gauze in order to secure obliteration, but at the time this was done I was unfamiliar with this procedure.

CASE 2.—J. S., an Italian blacksmith, 45 years old, was admitted to the City Hospital on Sept. 11, 1903. No history was obtained.

Examination.—He was well developed and nourished. Abdomen full, somewhat obese. The left side of the abdomen contained a large mass, which was hard, rounded and apparently more than five inches in diameter. It was dull on percussion, did not move with respiration, and the fingers seemed to get between it and the spleen. It was not especially evident by palpation in the flanks. There were no symptoms except slight abdominal pain. The urine was pale, acid, specific gravity 1007, albumin slight trace, quantity not stated.

Operation.—A diagnosis was made of tumor of the left kidney, and on September 17 he was operated on by Dr. J. B. Blake. A five-inch incision was made outside of the left rectus. A mass made up of multiple cysts, the shape of the kidney but four times the size, occupied the position of the left kidney. Transperitoneal nephrectomy was performed. The right kidney was examined by palpation through the operation wound. Palpation showed it to be somewhat cystic and slightly, if any, larger than normal.

Subsequent History.—September 21, after the operation and recovery from the ether, the patient passed his own urine for twenty-four hours, followed by a period of suppression. After twenty-four hours a sudden rise in pulse and respiration took place. He was catheterized. The urine was pale, specific gravity 1010. Chest examination showed congestion at the right base, with rales and high-pitched respiration. The pulse continued to rise, and in spite of stimulation and salt infusions he died three days after the operation.

CASE 3.—M. J., a woman, 33 years of age, single, was admitted to the City Hospital in the gynecologic service of Dr. C. M. Greene, May 9, 1906. A diagnosis was made of pregnancy, with eclampsia. For the past few months menstruation had been irregular and profuse, and for the last five months she had not menstruated. There had been noticed a painless enlargement of the abdomen, which had aroused suspicion of pregnancy. The possibility of this the patient denied. The symptoms before she entered the hospital were enlargement of the abdomen and dull, general abdominal pain. When she entered the hospital she was unconscious, but could be partially roused.

Examination.—She was well developed but poorly nourished, skin pasty and yellow, respiration slow and labored, heart negative. Urine, May 10, was acid, pale, 1010. Albumin, $\frac{1}{8}$ per cent. Much normal blood. Small amount of pus and cholesterol crystals. Two days later a few fine granular casts and many squamous cells were found. Blood count showed reds 2,200,000, whites 14,800. Hemoglobin 35 per cent. Differential count of 500 whites showed polynuclears 87 per cent., large mononuclears 10 per cent., small mononuclears 3 per cent. High color index. Marked diminution in number of blood platelets. Five monoblasts and four megaloblasts are seen in two smears.

Local examination showed the abdomen distended. A hard nodular, non-fluctuating, movable tumor could be felt in the right iliac fossa, hypochondrium and right lumbar region, extending over the mid-line to the left lower quadrant. The tumor was dull, not tender, and showed apparently a sulcus at the level of the anterior superior spine, running toward the gall-bladder region. A vaginal examination showed the os hard and the uterus not freely movable. The fundus could not be made out. A probable diagnosis was made of fibroma of the uterus, obstructing the ureter and causing a right hydronephrosis.

Operation.—Dr. George H. Monks operated by a median incision below the umbilicus. He found a mass which filled the entire abdomen, arising from the right kidney and covered with multiple, vari-colored cysts of varying size. The ureter was tied off and the tumor removed. The left kidney was also markedly enlarged and cystic. The uterus was normal. The patient was in a critical condition at the end of the operation and did not regain consciousness. Pulse 80 and soon became

weaker. The patient passed a fair amount of urine, but in spite of stimulation failed and died at 3:50 o'clock of the morning of the day following the operation.

Examination of the tumor showed congenital cystic kidney.

This case is of interest as showing that enlargement of the kidney in the female, associated with amenorrhea, may cause suspicion of pregnancy. A review of the symptoms indicates that the partial unconsciousness at entrance was undoubtedly due to uremia. The necessity of relieving the patient of the weight and dragging, together with the possibility that the tumor was a sarcoma, led to the performance of nephrectomy. The fact that nephrectomy was performed did not, apparently, affect the ultimate result, as the patient was markedly uremic and in very poor condition at the time of the operation.

The following cases, taken from the autopsy records of the pathologic department, are appended as illustrating the fact that the condition frequently escapes attention until revealed at autopsy. The first case is also interesting as showing cystic disease of the liver, which is not infrequently associated with cystic kidney. Its bearing on the etiology of the disease is interesting, and, as mentioned above, is confirmatory of congenital origin:

CASE A.—A woman, 47 years old, was admitted to the City Hospital Feb. 3, 1896. She died of purulent bronchitis and bronchopneumonia. "The liver extends 5 cm. below the border of the ribs. It is filled everywhere with cysts, which project somewhat above the surface of the liver in numerous places. They are lined with a smooth membrane, sometimes showing numerous projecting septa, which are filled with a perfectly clear, straw-colored fluid. The largest cysts are in the upper portion of the right lobe. They vary from 5 cm. in diameter to microscopic size. The consistency of the liver, apart from the cysts, appears normal.

"Both kidneys greatly enlarged. Weight of both, 700 grams. Capsule stripped off easily. Filled with innumerable number of cysts, ranging in size from pinhead to 3 cm. in diameter. They are in all parts of the kidney. Contain in part a perfectly clear fluid and in part a brownish, chocolate-colored fluid. On stripping off the capsule of the kidney numerous cysts are found. The fluid in them appears to be under considerable tension. None of these cysts communicate with the pelvis of the kidney. They are probably more numerous in the cortex than elsewhere. Both kidneys are equally affected. Both of same size. The renal tissue between the cysts is pale, reddish, somewhat granular and opaque. Markings can not be made out.

"Urine, 1026. Albumin, $\frac{1}{4}$ per cent. Much blood. Hyalin and fine granular and epithelial casts, with fat adherent. Compound granule cells covered with fat and epithelium. Large and small round cells. Casts of small diameter. Few pus cells.

"Clinical History.—Thirteen weeks of vomiting and diarrhea, swelling of legs and abdomen. Liver noted as enlarged, hard and irregular. Kidneys not noted as enlarged."

CASE B.—R. M., a man, 20 years old, was admitted to the City Hospital Nov. 25, 1898. Clinical diagnosis, malignant endocarditis. Anatomic diagnosis, endocarditis and acute lobar pneumonia. Kidneys.—Weight, 635 grams. Large. Surface shows everywhere cysts, varying in size from 1 mm. to 5 cm. in diameter. On section kidney appears to be made up of a series of cysts, which are found in both the medullary and cortical portions. In some places the cortex is entirely replaced by cysts. Cortex measures 7 mm., except over some cysts, where it measures as low as 4 mm. In other places it has entirely disappeared. In the cortex the glomeruli are seen as rather prominent grayish specks. Microscopically the kidneys showed very slight increase of connective tissue here and there in the cortex. Usually shows infiltration, with plasma cells. Some exudation of lymphoid and plasma cells in the intermediate zone.

Urine turbid, 1018. Albumin, slightest possible trace. Few hyalin and fine granular casts, some renal cells adherent and some of very large diameter. Numerous small round cells and considerable pus free and in clumps. Illness three weeks. No history of renal involvement.

CASE C.—M. L., a woman, 36 years old, was admitted Aug. 26, 1900. Glossitis. Purulent bronchitis with edema of lungs. Diaphragmatic hernia.

Kidneys encroach greatly on peritoneal cavity. Weight of right kidney, 1,160 grams; left, 780 grams. Right kidney, 27x10.5x8 cm. Surface markedly irregular and distorted, due to multiple cysts, which have apparently, except as mentioned below, entirely replaced normal kidney substance. Cysts vary in size from 1 mm. to 4 cm., thin-walled for the most part, so that the cyst contents are visible in many as a straw-colored fluid. Other rounded masses scattered at irregular intervals over the surface and surrounded by cysts of a dark grayish-brown color and of considerable firmness, suggesting the remains of kidney substance. Left kidney 20.5x10.5x7.5 cm. On posterior surface a firm grayish mass, evidently kidney substance, measures 4x2.5 cm. No abdominal tumor or symptoms on account of kidneys were noted during life.

Urine, 1012. Albumin, very slight trace. Sediment very slight, consisting of pavement epithelium. Cells from neck of bladder, coffin-lid crystals, bacteria, a few normal blood globules and leucocytes.

CASE D.—F. B., a man, 70 years old, was admitted to the hospital April 5, 1901. Death from double pneumonia.

Kidneys.—Right, 250 grams; left, 220 grams. From anterior surface of right kidney projects a thin-walled cyst, 4 cm. in diameter. At lower pole of same kidney a similar cyst, 6 cm. in diameter. Left kidney presents a similar cyst and several smaller cysts. Contain clear straw-colored fluid. Specific gravity, 1010. Slight fatty degeneration. Glomeruli stand out as bright-red points. Microscopically the glomeruli are large and injected. Slight focal increase of connective tissue, with atrophy of adjacent tubules. In places are dilated tubules (?), in which is granular material. Certain tubules contain granular casts.

No symptoms due to cystic kidney.

CASE E.—M. K., a woman was admitted to the hospital May 8, 1901. Ischiorectal abscess. Bronchopneumonia. Edema of lungs. Congenital cystic degeneration of right kidney, with compensatory hypertrophy of left. Age 60 years. Right kidney, weight 40 grams, measures 9x2.5x2.5 cm. Made up almost entirely of cysts from 1 to 2 cm. in diameter and filled with clear fluid. These are separated by only a thin wall of kidney substance. At the upper pole, kidney tissue reaches its maximum thickness, being there 1 cm. Hilum of kidney dilated, but ureter patent, showing no evidence of obstruction. Right renal cortex about one-third size of left. Left kidney, weight 235 grams, 14x6.5x4.5 cm. Smooth. Capsule strips easily. Pale on section. Glomeruli and markings of pyramids indistinct. Cortex 8 mm. thick. Many fat droplets in tubular epithelium.

No symptoms recorded in relation to kidney.

Urine, much pus. (Ischiorectal abscess.)

The case is evidently one of unilateral cystic kidney, with compensatory hypertrophy of the other.

Several other cases of cystic kidney in the autopsy records are not included, as a careful study showed that the cysts were secondary to interstitial nephritis, and the above cases are the only ones of undoubted congenital cystic kidney taken from a very large number of autopsy records. It is interesting to note that in none of them was a diagnosis of the condition made during life.

CONCLUSIONS.

From the above discussion we may draw the following conclusions:

1. If there be sufficient healthy tissue between the cysts to perform the normal renal functions the disease may never attract attention, and the patient may live a normal life and die of disease independent of the kidneys.

2. Under the same conditions as to renal tissue, while the renal excretion is normal, the cysts may cause pain and discomfort from their size and weight without dan-

gerous symptoms for years. Mobility of the kidney, with its accompanying troubles, may result from this enlargement.

3. The growth and enlargement of the cysts may so strangulate and compress the normal kidney tissue between them as to throw it out of commission, so that stasis and uremia may result.

4. The normal tissue between the cysts may become affected by the various forms of nephritis, just as healthy kidney may, and this nephritis may be fatal in itself or by reason of the cardiac and arterial changes which accompany it.

5. These same cardiac and renal changes may and do accompany the compression of the intervening tissue by the cysts, and are not an infrequent cause of death in these cases.

6. The surgical aspects of the disease relate to the pain and discomfort due to the size of the tumors or mobility resulting from them, and in operating for these conditions it is all important to remember the pathology of the disease and more especially the fact that it is bilateral in 98 per cent. of the cases. A cystic kidney, while causing pain and disability by its enlargement, may still be performing so large a part in urinary excretion that its removal would be dangerous to life from renal insufficiency.

7. It, therefore, becomes important if possible to take some means to diminish the size of the offending organ and anchor it in place without interfering with its secreting functions. Such a procedure, which will be vastly safer than nephrectomy, consists of the free incision and puncture of the larger cysts, thus diminishing notably the size of the kidney, and suturing of the organ in position. If for any reason a nephrectomy seems desirable it should not be performed unless an exploratory incision or ureteral catheterization has demonstrated that the other kidney is healthy.

529 Beacon Street.

OPHTHALMIC PRACTICE OF THE PRESENT TIME.*

FRANCIS VALK, M.D.

NEW YORK.

An extremely interesting paper by Dr. George M. Gould¹ on "A Study of Failures in Ophthalmic Practice" has called my attention to the subject of the work of the ophthalmologist of the present, and we may well ask the questions: What is it? What shall it be? Dr. Gould's candid acknowledgment of his failures and the reasons therefor will be readily accepted by most ophthalmologists, who will agree with the premises so well set forth in his paper. But I do not think, however, that the final word has been said as to the causes of our failures, nor do I believe that the present method of examination can not be improved. If we do have failures, such conditions must always exist, more or less, with all medical men, be they specialists or general practitioners. No one is infallible, even the most expert may be misled or fail by reason of conditions which the physician can not control. This subject can not be considered from any one standpoint; the work of all the investigators now working along these special lines in our profession must be included.

We may object to the statement: "By failures to

* Read in the Section on Ophthalmology of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.
1. Med. Rec., Sept. 30, 1905.