The Journal of the American Medical Association

VOL. XVIII.

CHICAGO, MARCH 12, 1892.

No. 11.

ORIGINAL ARTICLES.

THE TREATMENT OF CORNEAL ULCERS.

Read before the Milwaukee Medical Society, February 9, 1892.

BY J. A. BACH, M.D.,
OF MILWAUKEE, WIS.

One of the most formidable pathological processes of the eye the physician has to deal with is what is known as suppurative keratitis, or ulceration of the cornea. It is safe to state that more eyes are partially or totally destroyed through corneal ulcerations than through any other ocular disease, including injuries and operations.

It is with great anxiety that, when any of the purulent ophthalmies present themselves to the physician for treatment, he carefully opens the eyelids to inspect whether or not the cornea shows any evidence of ulceration. A sense of relief is felt to find the negative true. But not only in the suppurative ophthalmies is this so, ulcerations under many other circumstances may become equally unpleasant and formidable. The many staphylomatous and white eyes, and often no eyes at all, one sees in the various institutions for the blind, as well as on the streets, are a living evidence to this fact. It is not my object to speak of the pathology of ulcerative processes of the cornea, but to form a basis for a successful treatment. I will mention a few facts relative to corneal infection and ulceration.

Ulcerations of the cornea depend, with but few exceptions, for their immediate or remote cause upon some microorganism or ptomaine which, through some avenue or other, gains admission. It is a well-known fact that under apparently unfavorable circumstances the cornea has great power of resistance to infections; but once infected, it often becomes a very difficult matter to eliminate the infection. This depends upon the peculiar structure of the cornea. As you all well know, the cornea is made up of five distinct layers, an outer or epithelial layer being continuous with the conjunctiva; next comes a thin but very dense membrane, known as Bowman's membrane; then comes the corneal tissue proper, which forms about two-thirds the thickness of the cornea; beyond that there is the membrane of Descemet, another very dense and resisting membrane, and next to the aqueous is an endothelial layer. The outer or epithelial layer has but slight resisting power, but it is due to Bowman's membrane that deep infection becomes difficult. Should, however, the infection penetrate beyond Bowman's membrane it is easy to see how the less dense corneal tissue would become quickly infected, and being protected by the overlying dense membrane of Bowman surrounding the source of infection it is equally easy to understand the difficulty encountered in the effective application of remedies.

Although the cornea absorbs liquids with avidity, yet the more powerful antiseptic solutions, such as the sublimate, are not well absorbed, and consequently their value as active and reliable remedies in ulcerations of the cornea has, in many cases, shown itself to be insufficient. Within recent time the analine dyes, being readily absorbed, have been advocated as of great utility, and in the hands of those that have used them have given good results. There is the one great objection to their use, of leaving opacities. A most rational means of cure, but one which is being much neglected, to the great detriment of many eyes, is the use of the galvano-cautery. As a preliminary step of all the severer and deeper ulcerations of the cornea, nothing can commend itself as the red-hot point. From a surgical standpoint, this is the only rational means to favor a quick and safe conclusion of the disease process. With it we are enabled to, without the least danger, lay open our field of infection, remove the overlying obstructions, and give free access to our remedies to act, and thus insure a speedy termination of the ulceration. In conjunction with the above it is important to stimulate local nutrition and tissue change, which is best done by means of hot fomentations. The instillation of an atropine solution is also much to be desired, as it relieves the irritation of the ciliary nerves and spasm of the iris. The antiseptic and local stimulant of nutrition I have found to give the greatest satisfaction is a 10 per cent. aristol solution in sweet oil. This seems to be readily absorbed, is a comfort to the patient, and in my experience with it I have never found it to fail in doing good. In the lighter and more superficial ulcerations I have often depended upon it, instilling the same into the eye twice or thrice daily, after atropinization and drying of the eye. I have also found it to have a most happy effect upon the purulent process in cases of ophthalmia neonatorum, generally using it immediately after the local application of a 2 per cent. nitrate of silver solution.

Before reporting a few select cases I wish to recapitulate a mode of treatment which may be of value to you. Careful cleanliness in all cases is prerequisite. The use of any mild antiseptic solution such as boracic acid sol. or peroxide of hydrogen will answer the purpose. Atropine is used in nearly all cases, not so much for its curative effect as for the fact that it relieves spasm of the iris and ciliary irritation. It also assists in avoiding a possible prolapse of the iris in cases of perforating ulcer.

In peripheral perforations eserine is safer, though attended with pain at times.

In a case of simple ulceration of the cornea, the use of a 1 per cent. atropine solution combined with the instillation of a 10 per cent. solution of aristol in sweet oil, hot fomentations and bandage, gives, generally speaking, without any further means being

used, a speedy and satisfactory cure. Should the ulceration persist and spread in depth and width, be painful even after the above, it becomes advisable to resort to the galvano-caustic at once. A great mistake is often made by putting off its use from day to day until the ulceration has become quite extensive, necessitating large cauterizations. If used early and before the ulcer has reached the pupillary field, the cautery puts an effective check to its spread without leaving objectionable scars. If the ulcer is quite deep and a perforation seems to be inevitable, it is best to anticipate such in the performance of a paracentesis. We can thus better avoid a prolapse of the iris into the wound of the perforation. Paracentesis will not be found necessary often where the galvano-caustic has been used thoroughly at the proper time. Although being an operation that is quite innocent if properly done, it may be resorted to with benefit where the ulceration is deep and the pain great. By relieving the internal pressure it favors a speedy healing. After any of the surgical means used in any given case the use of atropine and aristol is to be continued in conjunction with the hot fomentations and rest in bed as the case may require, depending upon its severity.

Case 1.—Mr. Christian Differt came to consult me about a large *ulcus serpiginosus* covering nearly one-third of the cornea. Mr. Differt had been under treatment for over two weeks, but in spite of all treatment the *ulcus* grew larger and the pain became more unbearable. The history of the ulcer was that it started as a small marginal infection, which by degrees crept back and forth over the cornea until it had gradually infected a larger part of the cornea, and the remedies that were applied seemed to have had no effect whatsoever.

This is one of the most severe and persisting forms of ulceration and demanded a very decided interference. The thorough application of the galvano-caustic with the use of the other means mentioned put an immediate check not only to the further extension of the ulcer but to all pain and discomfort, and the patient could be discharged cured in a very short time.

Case 2.—Mrs. Martha Guember, after having been treated for upwards of three weeks for a deep annular ulcer of the left eye, covering a large portion of the cornea, presented herself for treatment at my office, and after a single application of the galvano-caustic with 10 per cent. aristol and atropine made a speedy and safe recovery without the least extension of the ulcer.

Case 3.—Mr. Coon came to me with a history of gonorrhœal ophthalmia with ulceration of the cornea. It was a very severe and almost hopeless case, yet by careful treatment as above outlined made a quick and very satisfactory recovery.

I do not wish to take up your time any further in reporting these separate cases but would state that in at least twenty cases during last year I pursued this treatment with such uniform satisfaction in even the most severe and hopeless cases, that I can highly recommend this course of treatment. The amount of cicatricial tissue produced by the galvano-caustic is not any more than the ulcer itself would leave, and if operated before the pupillary field is reached no harm can result from it.

DISCUSSION.

Dr. Hayes: Would you apply the cautery where the pupillary field of the cornea is affected?

Dr. Farnham: With reference to the peroxide of hydrogen treatment, can that be pressed and followed by the aristol, so that a large per cent. of cases are cured with it? Cannot you get the peroxide to penetrate and thus cure the corneal ulcer? I refer, of course, to the persistent use of the remedy. And cannot the wound be cleansed with it?

Dr. Würdemann: This subject is one on which we are all apt to temporize. I have seen ulcerations going weeks and weeks, and finally having to come to the galvano-cautery. In most all cases one application will heal a severe ulcer such as comes after gonorrhœal ophthalmia. The use of atropine in corneal ulcerations gives relief to pain, but probably has no effect on iris prolapse, because if the anterior chamber is cut open, or if there is a leaking of the aqueous humor, the atropine will have no effect upon the iris; but the pupil will remain contracted or in a state of partial contraction. Some place much reliance upon the use of eserine; it is claimed that it has a specific action in the healing of the cornea; I have been in the habit of using it sometimes in connection with atropine, using the latter purely on account of its analgesic effect.

Dr. Frank: Has not resorcin been used in similar ulcerations of the cornea?

Dr. Farnham: In the application of the cautery, do you cover the whole surface of the ulcer?

Dr. Würdemann: There are no bad results in the use of the peroxide of hydrogen with small corneal ulcerations; I have used it in one case where it raised the whole epithelial tissue, but the next day I found that no harm had been done; the gas had been absorbed and the tissue had gone back to place.

Dr. Bach: In reply to Dr. Hayes' question I would state that in case of the involution of the pupillary field and when the ulcer has perforated Bowman's membrane and entered the tissue, it is best to apply the cautery there, and check the spread of the disease and allow opportunity for iridectomy later on; in that way we may gain a very good vision, but otherwise we may destroy the field for iridectomy. The method is being used more and more with continued and increasing satisfaction.

With regard to Dr. Farnham's remarks about the peroxide of hydrogen treatment, I would say that while peroxide of hydrogen acts very nicely in some cases, nevertheless the activity is not great enough to destroy some of the severer forms of ulceration, and the spreading of the disease is a great source of danger; you can use the peroxide for cleansing the wound, but you cannot destroy the germ in the severer forms of the disease by the employment of it.

With regard to the employment of resorcin, the latest reports have not been as favorable as others; resorcin is not perfectly satisfactory in these diseases.

This is an ulceration which goes on under a dense, well protected membrane. The ulceration does not progress, as such, in the epithelial layer at all; the ulceration creeps under Bowman's membrane and the epithelial tissue proper, and the latter, of course, becomes destroyed.

The galvano-cautery should be applied only to the borders of the ulcer.

In reference to Dr. Würdemann's remarks, I should say that eserine seems to exercise a very favorable effect in some cases, but it is sometimes attended with considerable pain on account of the spasm it produces to the membrane.

With regard to raising the epithelial tissue by the use of peroxide of hydrogen, I would say that that happens quite generally. When the peroxide is used, it enters quite a distance under the epithelial layer; but it will disappear in a few hours.

Concerning drugs generally I would say that they are hardly sufficient and that the only rational means are surgical. For that reason I have dwelt considerably on the subject of the cautery, and since there is no objection to its use, I do not see why we should depend upon various drugs except for the purpose of cleanliness. The cautery is a simple, quick and effective means and saves a great deal of time and pain; it does not produce inflammation, and is applied painlessly by the use of cocaine, and in every possible way it is an advantage, and really it is the only rational method; which is not the case with any drug. They can only be auxiliary to the operation.

IMMIGRATION AND TYPHUS.—A resolution has been introduced in the national House of Representatives in concurrence with the Senate providing for an investigation by the Senate and House Committees on Immigration, of the law relative to immigration, and the facts attending the condition of the typhus-stricken Russians at New York.