he, his friends and lawyers instinctively combat any circumstance or statement tending to minimize his suffering. And finally, he is taught and teaches himself

Figure 9.—Case of general bruising—thrown from buggy by collision. Heavy line indicates tenderness along spine. Light line indicates same when examined from below upward; dotted line when examined from above downward. Light and dotted lines are to be thought of as occupying the median line.

Figure 10.—Case of railway accident. Black spot indicates location of excessive tenderness, circle shows place to which it shifted within a few minutes.

to look upon the future as one long valley of the shadow of death, ending only in death itself, for on the blackness of the future depends the brightness of the verdict.

How important, then, is the prophylactic treatment of these poor people. They are to be protected from ignorant doctors, foolish friends, designing lawyers and their susceptible selves. It is our duty—since it is generally within our power—to give them this protection; protection from ignorance by being not ignorant; protection from friends, first by perfectly understanding the case, then by clear and positive statements to the patient, and, last, by preventing or combating the officious suggestions of meddlesome well-meaners; protection from litigation by promoting prompt and definite settlement; protection from the patient himself by getting him out and at work as soon as possible, by begetting a wholesome and cheerful frame of mind and by getting the accident and all that pertained to it completely out of his life.

Because of the foregoing, in every case in which the question of “damages” arises or the possibility of litigation is present, the physician should advise both sides alike to make an immediate and final settlement.

A consideration of the treatment of the fully developed neuroses would carry me quite beyond a reasonable time limit. The proper methods to be employed may readily be inferred from the foregoing remarks and from our knowledge of the best means of handling the fundamental neuroses, but the successful employment of these methods requires the utmost tact, patience and energy—in short, the finest of medical technique.

34 Washington Street.

NOTES ON SMALLPOX.

S. L. JEPSON, M.D.

EX-HEALTH OFFICER.
WHEELING, W. VA.

I can not believe that there are many physicians who any longer entertain a doubt as to the genuineness of the mild smallpox that has prevailed to so general an extent during the past two years. The many valuable papers on this subject recently printed in *THE JOURNAL* should convince even Dr. Happel, whom we may honor as the leader of the doubters, and whose papers have no doubt led many inexperienced physicians into error.

Having served for eleven years as Health Officer of Wheeling, and observed several hundred cases of smallpox of every possible variety, and through a period of nearly thirty years, I wish to give my opinion, in the most emphatic way possible, that our recent smallpox, of which this city has had about eighty cases, was no new disease. In every outbreak of smallpox extremely mild cases occur, cases that are atypical and calculated to deceive physicians of limited experience. In the recent epidemic there occurred very many such cases, very few typical ones, and some unusual varieties, which defied positive diagnosis by the most experienced, unless watched closely for days. In my experience these most resembled varicella; but whereas the vesicles of varicella dry up in four or five days at most, that of even these mildest cases of smallpox ran a course at least twice as long.

Last November smallpox was brought to this city by colored convalescents from an infected camp of men working on a new railroad near Clarksburg. Within a few days fourteen colored men were taken down with it, all of whom contracted it from the same origin, in a saloon and gambling room conducted by a colored man, and patronized almost exclusively by colored men. All

were treated by me in the Municipal Hospital. Not one had ever been vaccinated, and yet there were all grades of severity exhibited. One man was in the hospital but six days, and was dismissed within ten days of the first appearance of the eruption, not having more than twenty vesicles on his whole person. Had this been an isolated case no physician could possibly have made a positive diagnosis, unless the history should have determined it. Yet in this same group of cases there was a confluent case, with a temperature of 108.5° , observed by myself at 4 p. m., and by the nurse at 8 p. m., taken in the mouth with a thermometer whose correctness I have verified. This patient recovered. Later I had a hemorrhagic case, which recovered. Were these last two cases variola vera? Even Dr. Happel would say, "Yes." So, therefore, were all the others, which represented all degrees of severity, although nearly all were very mild.

How was it in other localities? Called 70 miles away to determine the nature of an eruptive disease in an infant, I found a well-marked case of smallpox concerning which two physicians had differed; one called it chicken-pox, which it in no way resembled. A week later it proved fatal, and both parents contracted smallpox.

Dr. A. Wilson and myself were called to a mining town in Ohio, to determine a dispute between physicians and advise concerning an outbreak of an eruptive disease. We were taken to the house of one miner lying dead, after an illness of but three or four days. The appearance of the body and the history of the case left no doubt that the man had died of malignant smallpox. On the street we met another miner, just from his work in the mines, with an abundant variolous eruption, at about the tenth or eleventh day, and yet never ill enough to stop work for a single day. We found that fifteen cases had occurred, a number of the patients adults, and all had passed as varicella until a death aroused suspicions. Did the dead man have variola vera and all the remainder have varicella? A number of these cases were typical in appearance, although quite mild. In over thirty years' practice I have seen but one case of chicken-pox in an adult. I therefore hold that any acute eruptive disease in the adult that in any way resembles either chicken-pox or smallpox, should receive the same sanitary care as the latter disease, until its true nature is determined by close observation and time.

My observations here lead me to the conclusion that but little infection attends these very mild cases of smallpox. The primary fever is mild, sometimes passing unobserved; the secondary fever is often absent, and in many cases the eruption is very sparse, and it dries up without passing through the usual stages. As a consequence, a patient is less apt than in severe cases to convey the disease to exposed persons. Otherwise this city must have had hundreds of cases, for a number of patients strolled through our streets, frequenting saloons and other crowded resorts, for two to ten days after the eruption appeared. One man slept in a family of five, three of whom were unvaccinated children, for eleven days after the appearance of the eruption, and spent every evening at a social club-house. Not a single case resulted. One colored barber shaved his patrons for two days, and another for five days after the eruption appeared, and no case resulted. It is scarcely possible that all persons thus exposed had been vaccinated.

Only one of my patients made any claim to having been vaccinated, and I found no evidence of it in this case, one of my worst, which was in the hospital four weeks.

But one death in 76 cases occurred, and this patient was an infant. Why were the cases so mild? Dr. Hyde, in his paper in *Popular Science Monthly*, expresses the belief that the mildness of the cases in our recent epidemic was due to an immunity acquired by our people by inheritance, several past generations having been vaccinated. His own language is: "When it (the smallpox) gathered to itself the added power by which it was enabled to spread from community to community, its extension was not through a population virgin of protection by previous vaccination, or the children of the vaccinated." That a degree of immunity may be conferred by inheritance I am not prepared to deny, although this theory would be difficult to prove. The facts in reference to our recent epidemic can not be correctly held to support it in any way. After this smallpox entered the country from Cuba, it extended rapidly through the former slave-holding states, attacking chiefly the colored race, many thousands of whom were its subjects. These colored people in this city and state, and I think all through the South, are emphatically "virgin of protection by previous vaccination." Neither the present nor past generations of them have been vaccinated at all generally, and hence there was no modifying influence by inheritance. And yet, so far as my personal observation extended, the cases were quite as mild in the colored race as in the white. My very mildest case was in an unvaccinated black man.

Again, if there has recently been any such modifying influence by inheritance, as Dr. Hyde claims, it should have shown itself just as distinctly five or more years ago. But such was not the case. In 1895 and 1896 nearly 300 cases of smallpox occurred in this city and vicinity, but the cases were by no means of such mild type as those recently observed, and quite a number of deaths occurred.

No one has yet satisfactorily accounted for the mildness of our recent cases of variola. Has any explanation been presented of the recent mildness of typhoid fever, diphtheria and scarlatina? I am quite sure that none of these diseases occurs in this locality with the same virulence as in former years. No inherited modifying influence can account for this fact, and I think we will have to search further before we can explain why our late smallpox has been so mild.

As a sanitary officer, I have encountered much opposition to vaccination, not a few parents withdrawing their children from the public schools rather than have them vaccinated. In view of this fact, which is true of every community, and considering the very little suffering and very low mortality from smallpox at present, I have very seriously doubted the wisdom of communities going to great trouble and expense in the sanitary care of this disease, virtually to protect a limited number of ignorant or bigoted persons. The mortality has been nowhere greater than 2 per cent., which is much less than that of whooping cough, the suffering from which also is so great. We can do little, and we actually do nothing, to limit the spread of the latter. Why interfere with the former, against which each family can with certainty protect itself by vaccination?

Permanganate Treatment of Noma.—A. A. Kissel reports most encouraging success with local treatment of noma, supplemented by exceptionally nourishing food and strict hygienic measures. The lesion is curetted and then irrigated with a hot solution of potassium permanganate, concluding by dusting with iodoform. His statistics published in *Khirurgia*, x, 56, confirm the benefits of this treatment.