extracts than with salicylic acid. Incidentally, it may be said that Sadtler's procedure is more complicated than is necessary and correspondingly less reliable. The use of alcohol can advantageously be omitted. If ur-a-sol be shaken one minute with benzol and a little of the filtered benzol is poured directly into a very dilute ferric chlorid solution and shaken a very decided color reaction is obtained in the aqueous solution. Why diminish the reaction by the addition of alcohol?

(2) The Organic Chemical Mfg. Co. has made all the capital it could out of Professor Sadtler's accusation that Dr. Sollmann had erroneously quoted Fränkel as to the behavior of methylene-di-salicylic acid in the human organism. Fränkel has settled this point in favor of Sollmann.

(3) The question as to formaldehyd liberation when the formalal products are treated with various reagents (water, acids and alkali) has lost its significance since we now know that no salicylic acid is split off when the products pass through the human organism. To quote from the original report of Sadtler & Son "If salicylic acid is found so abundantly in the urine when acetyl-methylene-di-salicylic acid (ur-a-sol) is taken, it is evident that a corresponding amount of formaldehyd was liberated." That no formaldehyd is liberated is now equally evident from the demonstrated absence of salicylic acid from the urine.

The referee has gone over the tests made by Sollmann and by Sadtler & Son (except the phenyl hydrazin test which it seemed not worth while to take up) as to the liberation of formaldehyd by ur-a-sol, sodiformasal, iodomuth and guaialin. In view of the sensitiveness of these color tests it is entirely losing sight of the essential point at issue to dispute as to which test is more sensitive or more reliable. If formaldehyd were split off there would be no trouble about getting abundant tests and all results short of most decided positive ones are really negative in so far as any practical point of view is concerned. As a matter of fact, however, none of the products split off any formaldehyd when treated with water, hot or cold, or with dilute acid or alkali. Sollmann's original report is therefore correct and the findings of Sadtler & Son are erroneous.

Sollmann's later tests as reported to the referee give traces or "fair traces" by the Jorissen test when by Sollmann's original test they were entirely negative and by Sadtler's tests distinctly positive. Sollmann's faint reactions as reported to me are due, in the opinion of the referee, to the fact that he reversed the order of adding his reagents. What Sadtler's "distinct" or "strong" tests are due to, the referee does not know. When properly made the Jorissen test gives entirely negative results.

As to the tests used by Sadtler & Son and involving the use of concentrated sulphuric acid the referee entirely agrees with Sollmann who says that such tests are not adapted to prove the formation of formaldehyd under the influence of water or very dilute acids and alkalies. Sadtler's reply to this criticism of Sollmann is rather disingenuous: "How Dr. Sollmann can read into my description of my tests the idea that any of the substances in solid form were put into strong sulphuric acid I do not see, and yet that is what he is apparently talking about."

No, that is not what Sollmann is talking about. Placing "the substance to be investigated in contact with concentrated sulphuric acid" does not imply that the substance must be added in solid form. It also covers the conditions of Sadtler's experiments of adding the solution of a substance to be investigated in contact with concentrated sulphuric acid.

(4) For the sake of completeness the referee ought perhaps to explain the findings of Dr. Latham Clarke, whose report the Organic Chemical Mfg. Co. has quoted so ostentatiously in behalf of their contention that the taking of ur-a-sol gives rise to salicylic acid in the urine. Dr. Clarke was engaged by that company for the purpose of working out a practical method of manufacture of a certain product. On the last day of his engagement in the shop of that company Dr. Clarke was asked to test a "ur-a-sol" urine for salicylic acid. He did so and obtained positive results. The method consisted of passing compressed steam over the dried urine residue and

testing the solution so obtained by means of ferric chlorid. No attempt was made to determine whether methylene-di-salicylic acid would not also go over with steam. *The test was therefore, of course, not conclusive and Dr. Clarke authorizes the referee to say that he is no longer willing to stand for the validity of the test and further that he very much regrets that his name was ever connected with it.*

(5) As to the antiseptic properties of ur-a-sol, Sadtler & Son sent the referee a small sample of ur-a-sol urine which gave a fair ferric chlorid reaction. Twelve hours after this urine was received it began to grow cloudy and in the course of two or three days a considerable deposit had formed. The usual decomposition resulting in the formation of ammonium carbonate and a characteristic odor did not set in. The referee's own ur-a-sol urine behaved in the same manner, while a sample to which he added a little chloroform remained clear. Essentially similar results were obtained by adding ur-a-sol directly to a fresh sample of urine.

(6) Finally it should be said that the letter to the Organic Chemical Mfg. Co. from the secretary of the Council on Pharmacy and Chemistry with regard to the rejection of its products might have been more full in which case it undoubtedly would also have been more accurate. The brevity and the inaccuracy of the letter was, however, more than compensated for by the fact that the company also received the full report of the referee. The pamphlet published by the company indicates that the author of it understood very well the import of the details contained in Sollmann's report, yet chose for advertising purposes to confuse the contents of a mere routine letter with the well considered statements of the referee, Dr. Sollmann.

The referee does not recommend reconsideration of the Council's action on the products of the Organic Chemical Mfg. Co.

Correspondence

Serum Sickness

To the Editor:—The increasing interest in serum poisoning, as shown by the recent report of cases, leads me to relate the following personal experience. I am furthermore induced to do so by the possibility herein shown of the sensitive condition far exceeding the usual time limits.

In 1902, I attended a case of diphtheria. I took an immunizing injection of antitoxin of 1,000 units. No inconvenience or rash followed. A few months later my duties brought me into close relation with bubonic plague and I was injected with Haffkine's prophylactic. This is a vaccine. A very severe reaction followed, temperature rose to 102 F., there was much urticaria and general malaise for two days. I looked on this as a toxic and not a serum reaction. In January of this year I was in very close relation to an unusually severe case of diphtheria, in which I administered 108,000 units of antitoxin to the patient, a child of 7. The latter recovered, slight multiple neuritis followed, but no serum symptoms. The hospital interne and myself were both much exposed to infection, and we thought it advisable to take an injection of 1,000 units each. In the case of the interne no results followed. I remained quite well for seven days, when suddenly much painless swelling, in and around the parotid, appeared. This subsided by the next day, to be followed by the sudden appearance of urticaria, distributed much as in herpes zoster, over the course of an intercostal nerve corresponding to the point of injection of the serum. It subsided in the course of a few hours, to be followed at 10 p. m. by an urticarial eruption over every part of the body. The irritation was unbearable as the night advanced, the temperature rose to 102.5 F. The pulse was 160. Vertigo and partial syncope followed any attempt to rise. I think this may be accounted for by the great engorgement of the skin depleting vessels and brain. Toward morning, much relief was obtained by bathing the surface with ammonia and water (about 1-10). The next day was passed in bed in a state of prostration. There was some recrudescence of the urticaria in the evening, promptly allayed by the ammonia. Good sleep followed, and

the next day I was well except for some pains in the joints, which persisted for nearly a week.

Now the main interest of the case lies in the fact that in 1902 I was not susceptible to the toxic action of the serum; in 1909 I was intensely so. The serum of the latter date was not toxic in itself, as shown by its lack of action in the case of both patient and interne—ergo it would seem as though I had been activated by the first dose and remained so after a period of seven years.

We clearly need the aid of a careful investigation into the question of time limit. It may well be that my experience was not so exceptional as it appears.

H. D'ARCY POWER, M.D., San Francisco.

Delegates for Pharmacopeia Convention

To the Editor:—In accordance with the provisions of Article VIII, Chapter I, of the by-laws of the U. S. Pharmacopeial Convention, the president of the convention hereby invites the several bodies, entitled under the constitution to representation therein, to appoint delegates to the first decennial meeting of the said convention to be held in the City of Washington, May 10, 1910.

The attention of all concerned is invited to the following extract from the constitution from Article II on membership:

SECTION 1. The members of the United States Pharmacopeial Convention, in addition to the incorporators and their associates, shall be delegates elected by the following organizations in the manner they shall respectively provide: incorporated medical colleges and medical schools connected with incorporated colleges and universities; incorporated colleges of pharmacy, and pharmaceutical schools connected with incorporated universities; incorporated state medical associations; incorporated state pharmaceutical associations; the American Medical Association, the American Pharmaceutical Association, and the American Chemical Society; provided that no such organization shall be entitled to representation unless it shall have been incorporated within and shall have been in continuous operation in the United States for at least five years before the time fixed for the decennial meeting of this corporation.

SECTION 2. Delegates appointed by the Surgeon-General of the United States Army, the Surgeon-General of the United States Navy, and the Surgeon-General of the United States Marine-Hospital Service, and by the organizations not hereinbefore named which were admitted to representation in the convention of 1900, shall also be members of the corporation. Each body and each branch of the United States Government above mentioned shall be entitled to send three delegates to the meetings of this corporation. But no such delegates as are provided for in this article shall be members until their credentials shall have been examined and acted on as provided for by the by-laws. Delegates admitted as members at any decennial meeting shall continue to be members of the United States Pharmacopeial Convention until their successors shall have been appointed and admitted as delegates to the ensuing convention, and no longer.

Notification of the appointment of delegates, accompanied by the necessary certification of eligibility as required by Article II, Section 1, of the constitution above quoted, should be forwarded as soon as practicable to the secretary of the Board of Trustees.

HORATIO C. WOOD, M.D., Philadelphia, President.

MURRAY GALT MOTTER, M.D., Secretary of the Board of Trustees, 1841 Summit Place, Washington, D. C.

Miscellany

Care of Weak Infants and Children.—That something more than temporary relief or routine institutional care is necessary to improve the defective vitality of the sick and ailing infants of the poor and combat the influence of their wretched environment is quite evident from the indifferent results obtained from ordinary methods. Dr. Henry Dwight Chapin (*Charities and The Commons*, March 27, 1909) describes the methods adopted by the Speedwell Society in caring for this

class of children. The plan followed is the boarding-out system, and the results have been remarkably good considering that bottle-feeding has been employed exclusively. The children have been placed in carefully selected homes in a healthful location. A physician and a trained nurse have oversight of the cases. The foster-mother is instructed in the feeding and care of the baby and receives \$12 a month as compensation. Many take a great interest in the work and are very successful in nursing feeble infants back to health. Some tire of the work and others are dropped because of inefficiency, but the aim is to retain one same families in the work so far as practicable. Dr. Chapin says: "The boarding-out method, under careful supervision in the country, is the best plan yet devised to relieve and save these infants. The first series of 121 cases would nearly all have died under ordinary methods of handling and yet nearly two-thirds of them were saved. They were poorly nourished, from bad hygienic surroundings, with various degrees of digestive disturbances from faulty feeding on the bottle, and stationary or losing weight. A large proportion were restored to fair and even vigorous vitality, although kept on the bottle. The greater number of those from three to six months to one year was saved, in the first series nearly three-quarters being the proportion. These are brilliant results considering that artificial feeding and was still of necessity exclusively employed."

Medical Statistics for European Russia in 1906.—The *St. Petersburg. med. Wochenschrift*, March 27, published the vital statistics and public health reports just issued for 1906. The total mortality is stated to have been 29.2 per thousand inhabitants, the births 45.9 per thousand. There were 17,096 civilian physicians, 21,670 "feldshers," 11,032 midwives, 4,917 dentists, and 8,828 pharmacists, to a population of 106,114,589. The number of sick treated in hospitals during the year was 2,458,207, with a mortality of 5.2 per cent. There were 133 asylums for the insane, with 30,671 beds and 566 maternities or obstetric wards with 4,755 beds, and a record of 178,877 deliveries, with a mortality of 0.5 per cent. There are 20 Pasteur institutes, and 19,668 persons were given Pasteur treatment during the year, of whom 106 died. There were 178 medical societies and 44 hygienic laboratories. The total outlay for medical purposes was \$38,760,000 (76,000,000 rubles). Of this 7 per cent. was borne by the central and 50 per cent. by the provincial treasuries, 24 per cent. by the cities and 17 per cent. by private individuals and societies.

Appendicitis in Childhood.—In *Hygiea Festband*, Carlson discusses acute appendicitis during the first fifteen years of life on the basis of 47 original cases. He believes that appendicitis is more serious in children than in adults, because it more often passes over into diffuse peritonitis, because changes in the appendix more often are gangrenous and because the septic form is relatively common. The death rate in his cases has been much greater than in the case of adults, the probable reason being that the children have been seen too late for operation. Every child diagnosed to have appendicitis during the first 24 hours of sickness should be operated on before the end of that time. Operation is also indicated during the second 24 hours. During the third, fourth and fifth days operation should be made in every case that does not show definite and persistent signs of improvement; but it is to be remembered that during this later period the operation is technically much more difficult. After the first attack operation should be made in the free interval. The points on which Carlson lays special emphasis are the great danger of appendicitis in children and the necessity of early and correct diagnosis—not an easy task—so that operation may be done as early as possible during the first attack.

Experimental Researches on Some Ferments of the Bile.—G. Bonanno (*Arch. di farm. sper. e sc. aff.*, vii, 466) finds that the bile of the human being, of pigs, dogs, sheep and calves always possesses amyolytic and lipolytic properties, but no proteolytic action. The amyolytic action is stronger in the bile of herbivora than omnivora, while the lipolytic is about the same for all. Bile from either animal after the addition of pieces of duodenal mucosa shows no proteolytic action.