in the same individual not uncommonly. Of these 34 cases 15 had hemorrhage from the stomach, 8 from the nose, 7 from the bowels, 1 from the mouth and gums, 1 from the kidney and 1 from the genitalia. Next to splenic enlargement the anemia was the most striking symptom, being well marked in all of the cases, with a low color index and usually a leucopenia. In only two cases was there an excess of leucocytes. The average percentage of hemoglobin was 47 per cent., average number of red blood corpuscles 3,293,000, and of white corpuscles 5,594. If we exclude the two cases in which there was a leucocytosis, the average number of leucocytes is 4,381. The liver was enlarged in 16 cases, small in one case (Benedict). Ascites was present in 7, swelling of the feet and legs in 9, glandular enlargement in 3, gastrointestinal symptoms in 8, and cardiac murmurs, generally hemic, in 11. In only two cases aside from my own is there any report of albumin in the urine and then "only a trace," while my case has had considerable albumin at times, as well as some pus and casts.

The summary of the treatment and its results is most interesting. Of the 25 cases which were treated medically only 5 are recorded as "recovered," 12 were improved, 1 "not improved," and 7 died, hemorrhage having been the most common cause of death. Operation was performed on 8 cases, of whom 5 recovered and 3 died.

SQUINT.*

ALBERT B. HALE, M.D.
CHICAGO.

Squint ought to be made an annually discussed subject at mothers' meetings, for it is through the mother that it must be detected, if this improper functional condition is to be corrected. It is not merely a question of two good eyes, although physiologically we need all our organs in the best of working order; cosmetically, the individual is nearer the type of perfect man if he not only sees with both eyes at once, but if the intended relationship of one visual axis to the other is constantly maintained; if no squint is observable. The young adult feels the effect of a cosmetic imperfection very keenly, and is grievously disappointed to find, when treatment or cure is sought, that the operation does not always lead to even cosmetic improvement, and that in the great majority of cases the squinting eye, if it does not persist in squinting, is, after all, a rather useless organ, on which no trust can be placed.

The correction of a squint, and if possible the restoration of function, must, then, be begun in early childhood. This is a fact not so much known, nor so usually accepted, even in our own profession—not even among all physicians who make a specialty of the eye—and the family counselor is sometimes guilty of saying that the child will outgrow its squint, or that the time to heal it, perhaps to operate on it, is about or after puberty. No greater error can be made. The time to treat a squint is as soon as it is noticed, which is very early in life, and the causes of the squint must be well understood before this advice is taken to heart and any treatment attempted.

Let me emphasize this statement: that many completely blind eyes do not squint—they jog along with their fellows content to obey the impulses from the brain without protest or discontent. On the other hand, most squinting eyes do see something, or did ear-

lier in life, and at the beginning of the trouble they protested vigorously at the impulses from the brain and tried to act by themselves; tried to see something till they fell into bad habits, and long after they gave up trying to see accurately, they continued these bad habits and can not be reformed.

What, then, is squint? Of course, there is a paralytic squint, due to some acquired interruption to the nerve impulse or to injury to the muscle, but these are always characterized by double vision and usually come on later in life, through disease or trauma after the fusion faculty has been established. This may be called an organic lesion. The type to which I refer is, in contrast, called concomitant squint and has its origin in a functional disturbance, and the disturbed function is that of fusion. It arises early in life before the function is well established, and not being such a long-inherited function or instinct—for we know that binocular single vision is possessed by but few of the highest animals—it is easily disturbed. Generally, there is some exciting cause, such as a fever, a shock or fright, in childhood before the sixth year, while this habit is becoming acquired, and when weakened or lost, it is so delicate a function that neglect may often destroy it altogether, while to re-establish it demands attention, training, or the stimulus to spontaneous exercise of the habit, by placing the eyes in such relationship to each other that it is as easy to exercise as it is to neglect this function.

Herein lies the great error on the part of both physician and parent, the assumption that the child will "grow out" of the habit of squinting. Perhaps he will, but in the great majority of cases only at the expense of the vision of the squinting eye. Once have it well understood that treatment should begin early, as soon as the defect is seen, and there will be fewer squinters in the world, more sound eyes, and less complaint that the operation so seldom accomplishes what is expected of it.

The diagnosis is easy. Any layman can tell when a child or adult squints, but the etiology in the light of modern study shows that neither nerves nor muscles are primarily at fault. It is the brain, where the fusion function is located, and it is the brain we must first educate before we can hope to control the eyes. Usually—not always—one eye will be found to see less distinctly than the other eye, the difference in degree varying with the individual and the duration of the squint. The result of disturbed fusion sense is imperfect innervation, exaggerated accommodative effort in one eye—squint.

The treatment resolves itself into three factors: (1) Correction of refractive errors; (2) training of fusion sense; (3) operation.

Of these, the first is the essential factor, and without it practically no case of squint can be overcome, or if the eyes do later in life grow straight, it will be at the expense of the vision of one eye.

The second factor is of great importance, and implies such methods as the occlusive bandage, perhaps atropin to the good eye, stereoscopic, amblyoscopic and other visual exercises.

The third factor is operation, which is usually necessary, though squint may be entirely overcome without it. As a rule, however, some operation on a rectus muscle is necessary, and my later preference is for the advancement of a rectus muscle.

103 State Street.

* Read before the Chicago Medical Society, 1903.