tests are concerned, I believe the reader of the paper is correct when he states that all are objectionable. A certain percentage of error in the use of von Fleischel's instrument can be reduced. I have come to the conclusion that very frequently, in making our comparison by looking through the entire field of the cylinder of distilled water and diluted blood, the attempt to cover this broad surface in one view causes a certain percentage of error. About a year ago, at a meeting of the Philadelphia Pathological Society, I suggested that a cap be placed on the cylinder, containing a slit measuring one-eighth inch in width, made of any material, which would limit the field of observation, and thus reduce some of the errors of the instrument. Another error to be overcome is the distance at which the eye is held from the instrument; a distance of eight or nine inches is the best. A number of other interesting sources of error regarding von Fleischel's instrument might be mentioned, but time will not permit. I agree with the reader of the paper that we do not usually get 100 per cent. of hemoglobin; the percentage is usually lower.

It is also well to remember that if two pipettes or more be used, the reading will be more accurate in cases where the percentage of red blood-corpuscles is less than 50. Gower's instrument is seldom employed; it is subject to a wide range of error. Occasionally the colored tube of gelatin is decolorized in the course of time.

As regards the method described, I have had no personal knowledge as to its use. The percentage of blood pigment as determined by various methods differs greatly. From a long series of observations extending over a considerable period of time, it is necessary to determine which method is the best. The translation of the hemoglobin percentage is subject to considerable variations taking place in the plasma, but this percentage is not very great; it is exceedingly limited. Although it is true that the specific gravity of the blood is chiefly due to its contained hemoglobin, the plasma also contains substances that vary in disease. It is therefore manifest that the specific gravity of the blood should not be considered as the exact equivalent of the amount of blood pigment present.

Dr. GEORGE DOCK, Ann Arbor, Mich.—I would like to corroborate everything the reader has stated regarding the advantages of the specific gravity method. It is easy to carry out. It must be borne in mind, however, that the specific gravity does not always correspond to the amount of hemoglobin present. A large amount of comparative observations with the specific gravity test and von Fleischel's hemometer have shown that hemoglobin chiefly affects the specific gravity, and there are few cases in which the readings do not correspond. Gower's method is much more faulty than von Fleischel's. In using the specific gravity method great care is necessary. If the estimation is difficult to make, a fresh drop should be used as a control, for in a trial which takes some time the density of the drop becomes altered by various causes. I have long taught that if one is limited to one method of estimating hemoglobin, Hammerschlag's specific gravity method should be chosen rather than Gower's instrument. The cap devised by Dr. Daland has for some years been a part of Meischer's modification of von Fleischel's instrument, a modification which also does away with several of the other sources of error in the old apparatus. The instrument is expensive, but we can hardly give up the estimation of the coloring matter of the blood, and for that purpose it should be preferred to the old apparatus or to that of Gower. The practitioner can, however, get along very well by the aid of Hammerschlag's method, carefully carried out.

Dr. HORACE B. ARNOLD, Boston—I made quite a number of experiments this spring on the effect of chlorosis on the heart, using the color and specific gravity methods in these cases. The results obtained are important, especially in the case of chlorosis, where the hemoglobin has a low percentage; this might have been due to the heart changes or to error in the color tests. In this case there was particular satisfaction in using the specific gravity method. The comparison made between the color test and the specific gravity method by my assistants showed that they ran close. In considering the cases of chlorosis it is important to take count of the blood-corpuscles, because in some cases we find an improvement in the condition with an increase in the amount of hemoglobin, and an increase in the number of red blood-corpuscles out of proportion to the increase in the percentage of hemoglobin. So in this case apparently there is a diminution in the percentage of hemoglobin.

ANGINA PECTORIS.*

BY JAMES DUDLEY MORGAN, M.D.

ATTENDING PHYSICIAN AT GARFIELD AND GEORGETOWN HOSPITALS, ETC.
WASHINGTON, D. C.

Pain is so seldom directly associated with serious diseases of the heart that most specialists and practitioners of medicine are wont to treat lightly a patient's anxious inquiry about the heart, and to dismiss the case with no particular guidance or injunction, as one having an imaginary or neurasthenic origin.

Though it has been considerably over one hundred years—1768—since Herbenden gave the name of angina pectoris to a certain affection of the heart, which he declared must be distinguished from ordinary cardiac dyspnea, and a few years later Edward Jenner described some of the morbid conditions, which may be found in this disease, yet there is to-day no one certain and positively accepted statement of the cause of this sudden taking off of a human being, who may have previously complained but little, or not at all, of feeling badly.

Since Herbenden's time nearly all medical writers and clinicians have had something to say of the etiology of angina pectoris. Desportes located the disturbance in the vagus; Laennec in the sympathetic; Jolly in the intercostal nerves; Brouillard in the phrenic nerve; Kirsch in the spinal cord; Piorry and Cahn placed it in the brachial plexus; Romberg, Peter and Lanaceaux in the cardiac plexus; Leyden, Potain and Charcot express opinions not in harmony with each other; v. Basch declares it to be sclerosis of the coronary arteries; Osler says that the disease is associated more particularly with sclerosis of the root of the aorta and changes in the coronary arteries and in the myocardium. Constantin Paul, of Lariboisiere Hospital, acknowledges the frequency of atheroma in patients who have died of angina pectoris, but when he considers the large numbers of patients at Bicêtre and la Salpêtrière, who suffer from atheroma and are free from angina, he can not convince himself that the two diseases bear an undeniable casual relation to one another.

Kinnear believes the disturbance is due to hyperemia of the spinal sensory centers. Anders in his late "Practice" concerning the nature of angina, says: We possess few if any positive data; and conclusive post-mortem evidence in support of the various theories that have been and are advocated is, as yet, wanting.

Theodor Schott of Bad Nauheim, Germany, in a very recent article on the treatment of cardiac neuroses, speaks of: 1, angina pectoris nervosa; 2, angina pectoris vasomotoria, and this complaint he attributes to spasmodic contraction of the vessels, and mentions that besides the disturbances of a nervous character, there exist also other abnormal changes in the heart; 3, a form of angina pectoris he calls vera, and states it is more frequent than the nervosa or the vasomotoria, a statement with which the writer does not believe most practitioners will agree. Musser says angina pectoris in its typical form and in association with diseases of the heart is not of common occurrence. Schott goes on to speak of the difference of the pulse in true angina pectoris, which was not appreciable to the finger and only to be found in sphygmographic tracings and which often indicated arrhythmia, tachycardia and bradycardia. He believes that the most frequent causes of this complaint are arteriosclerosis, chiefly of the coronary vessels, insufficiency and stenosis of the aortic valve, myocarditis, etc.,

*Read before the Medical and Surgical Society of Washington, D. C., April 6, 1899.

but he must confess that as yet the opinions of medical men concerning the nature of angina pectoris vera do not quite agree, but in general we are nowadays led to suppose that this disease is apt to affect the heart, when the latter has been enfeebled before.

These various affirmations and negations of conditions, which are necessarily found or are wanting in this much-discussed disease, are not brought forth to further mystify opinions, but with the desire that we may recall how little we know of angina pectoris and that a free discussion may engender more positive knowledge.

It could hardly be expected that this paper, in the time which is allotted, can take into consideration the numerous pseudo-anginas of the heart, for their causes are legion and spread over and ramify the body from migraine to podagra. It is freely admitted that cases of pseudo and true angina are sometimes associated, and that pseudo-angina may aggravate and precipitate an impending attack of true angina. The onset of this terrible disease is so sudden and startling, the suffering so great and agonizing, the termination so rapid and fearful, that consternation and fear seize both patient and friends. Balfour described the sensation of angina "as if a mailed hand grasped the chest in the cardiac region and squirted through its fingers flashes of excruciating agony."

A male, aged 42, who had previously shown no cardiac disturbance and was otherwise healthy and attending regularly to business, was seized on preparing to take his usual morning bath, with the most horrible constriction and agonizing pain in the chest. Within a half hour, after suffering untold torture with the agonies of pent-up life and suffocation, he died. At the autopsy, the left ventricle was dilated and the myocardium showed increased growth of connective tissue at the expense of the muscular tissue. The coronary arteries were in no sense occluded and the aorta was not sclerotic.

Another patient was a middle-aged married woman who had informed me that six years before, on a trip to Europe, she had been taken suddenly with the most fearful and terrifying pains over the region of the heart, associated with a sense of great suffocation, the heart feeling as if clamped within a vise; that Dr. Austin Flint, Sr., was aboard the steamer and attended her, and after some ten to fifteen minutes she was able to breathe more freely and was rid of the pain, but felt most profoundly shocked and depressed for the following twenty-four hours. He told her she had angina pectoris and must be careful in the future about her mode of living. She had no perceptible organic lesion. Being hastily summoned one day to her residence, which was some ten minutes' walk, I found she had suddenly been seized with a dagger-like pain in the heart and, exclaiming, "O, my heart," died within a few minutes. Autopsy revealed no morbid condition of the heart nor its vessels.

The writer has had what he considers a few cases of true angina pectoris, in which happily the fatal termination was at least postponed, and will present the notes of one or two, which seem most characteristic.

During the early hours of the morning, being hastily summoned to see a gentleman, aged 57, who was declared to be dying, I found the patient sitting upright in bed, with his hands over his heart, and laboring for breath. His hands and feet were cold, face blanched, pulse small and inclined to irregularity. He had a most distressing appearance, and I expected him to collapse at any moment. I gave him ether internally and nitrite of amyl

to inhale, and applied heat over his heart and to his extremities; later on dry cups were applied over the chest. This patient recovered after a severe struggle, and within less than half an hour pain had ceased and breathing was more quiet; but it was three days before he could resume his journey to Boston. I was informed that the patient had a similar, but much lighter, attack, some eighteen months before. His heart showed no organic murmurs, but the rhythm was feeble. There was no perceptible arteriosclerosis. Examination of urine was negative.

Another patient, living to-day, whom the author attended some ten years ago for frequent attacks of angina pectoris, has some interesting points connected with his case. He was a man about 40 years of age, of no bad habits nor history, inclined somewhat to corpulency, but had no digestive disturbances, nor was he in any way a neurasthenic. For a man of his size and strength, he had what you might call a feeble and somewhat compressible pulse. The heart showed no intrinsic nor extrinsic disturbance. The urine was repeatedly examined with negative results. The gentleman has had some eight or ten characteristic attacks of angina pectoris, extending over several years, and but for speedy help and remedies, which he has been taught to keep on hand, he would have, it appeared, died in one of these attacks. The treatment consisted in a complete change in his mode of living. He was made to work less, and take moderate exercise; was forbidden alcohol in any shape, and less nitrogenous food was eaten; he was to lie down after bathing; take a midday siesta, and lead a life which would in every way relieve the demands on the heart. Massage was used and the precordial region was kept slightly irritated for months, and hot foot-baths were given at night. His slight indulgence in both coffee and tobacco was stopped, and he was ordered ether, iodid of potassium and trinitrin, which he took on different occasions, extending over a period of nearly a year. On retiring at night he was ordered to give himself a hot saline enema, which had a most salutary and beneficial effect on the various emunctories, besides giving tone, without tension, to a somewhat enfeebled arterial circulation. His bedroom was changed from the first to the third floor. His trips away were to the mountains instead of to the lowlands and seashore. This patient has not had an attack for eighteen months, and his wife and self are congratulating themselves that there will be no recurrence. He may yet die suddenly of this affection, for we are too often deceived, when we relax our vigilance and persuade ourselves that recovery is in sight.

William Pepper, who suffered for some seven years from premonitions of angina pectoris, was joyous and hopeful that compensation was taking place and remarked to his physician but a few days before his death that "the battle had been won." "At seven in the evening," his physician writes, "I left him gazing upon Mount Diablo, shadowed in the gathering darkness. I was called at 8. I found him in the attitude and with the expression of *angor animi*, from which he never aroused. At the time of his death he was recovering the lost cardiac compensation and appeared on the clear road to recovery."

Any affection of the heart or systemic system, which may disturb the regular and proper supply of blood to the coronary arteries, may precipitate an attack of angina pectoris. The danger lies in the myocardium not receiving sufficient blood for its nutrition and so cause the generation and accumulation of toxins,

which poison and depress the heart, and allow of an abnormal distension of its cavities. This distension of the heart and interference with its nutrition accounts for the great pain and distress in the precordial region. Smirnow, in speaking of the dilation and distension with blood, which the heart undergoes in the paroxysm of angina, attributes the torturing precordial pain to the stretching of the net-work of afferent nerve-fibrils, which are immediately beneath the endocardium.

The quality of the blood supplied to the nutrition of the heart is the cause of many true anginas of the heart. The effects of blood changes on the heart have been investigated by several authors. Ringer found that without the presence of a lime salt in the circulating fluid, contractility of the isolated frog's ventricle cannot be supported. He also found that the production of acid, particularly that which is formed in the course of, and as the result of, muscular contraction, is also inimical to cardiac contraction. A toxin within the blood depresses the heart, increases the length of the systole and finally causes a standstill in diastole. The accumulation in the heart's substance of a toxic material seems to act on the cardiac ganglia in such a way as to destroy the heart's excitability. Any slight obstruction of the supply of blood to any part of the myocardium leads quickly to the generation of a toxin. Cohnheim says the closure of one or both coronary arteries is not necessary, any one branch of a certain size, no matter which, is sufficient to produce a limited arterial ischemia, which in such deadly manner paralyzes irrestorably the whole cardiac musculature. The effect of this deficient nutrition is the product of a heart poison, a product of metabolism, which is in a normal state carried off in the course of the blood-current, but which accumulates under defective nutrition in the substance of the heart.

What makes the action of the heart regular? It is the change of nutrition, by which, through the blood circulating in its substance, the waste of its tissue is constantly supplied, the integrity of the fibers preserved and kept consequently in a condition to contract. If the supply of blood is cut off from the substance of the heart, the organ soon loses its irritability and power of contraction. Cohnheim and Schulthess-Rechberg, in experiments on dogs, observed that ligation of any large branch of either right or left coronary "has at first no effect whatever upon the rhythm or vigor of the cardiac contraction, nor consequently upon the blood pressure, but after the lapse of ninety seconds on an average, the heart beats begin to be somewhat irregular and infrequent, yet still without affecting the blood pressure, till suddenly and at the same instant both chambers stop in diastole. From this standstill, which occurs on an average not later than two minutes, there are no means of arousing the ventricles to new life and renewed contraction. It seems as if a deadly poison had forever destroyed the heart's excitability. Accordingly it is in all probability the system of ganglia which is affected in this fashion."

It would seem from all the facts at hand, clinic, pathologic and physiologic, that whereas sclerosis of the coronary arteries or aorta, or dilation of the heart, etc., is found in a large number of cases of angina pectoris, yet attacks of true angina may occur in patients who are entirely free from organic disease of the heart, and, vice versa, we may find most extreme calcareous conditions of the coronary arteries to exist, as in a pathologic specimen I have, from the patient living to some 60 years of age, with no obtainable history of cardiac distress, and dying of a complaint entirely foreign to angina pectoris. Frey maintains that stoppage of the heart in

angina is not a necessary result of closure of a large branch of the coronary arteries, but may be brought about by numerous other interferences with the heart.

In our treatment we must ever keep in view the picture of the heart whose muscular structure is improperly nourished, whose ventricles are suddenly distended, and whose nerve and muscular fibers are stretched, and I might say stunned, by the rapid engorgement of a circulating fluid, which from the apparent embarrassment in respiration, becomes poorly oxygenated. Relieve the tension of the vascular system by inhalation of nitrite of amyl, and if pain persists, give a few whiffs of ether, which, as there is danger of heart failure, is to be preferred to chloroform on account of its stimulating effects. Morphia should not be continuously used because of its resulting depression, and if given should always be combined with atropia, which is an excellent analeptic. Camphor, also, is one of the best of the cardiac stimulants. Irritation, in the way of sinapisms, etc., to the precordia, eases the distress. The ice-bag or very hot applications have the same effect. The kidneys, skin and respiratory functions should be kept active. In the interval nitroglycerin or nitrite of sodium and iodid of potassium should be used, and the doses should be pushed to tolerance. Good results are obtained from the baths as recommended by the Oertel's method or as carried out by Dr. Schott of Nauheim. A saline bath at the temperature of 92 and reduced 5 or 6 degrees in the course of the treatment has many of the good effects of the Nauheim bath. It is well in ordering the bath to see that your patient is gently massaged, and while in the bath, which should last not longer than ten to fifteen minutes, that slight restraint be given to the flexion and extension of the limbs and body, which the patient is directed to follow. These movements draw blood to the parts and lessen the arterial tension in the system, and with the stimulating effects of the bath have a most beneficial and somewhat lasting good effect on the heart and circulation, the blood pressure being lowered, the diastolic rest of the heart lengthened, and the force of the systole increased. The result is, the ventricles are fully emptied, the peripheral resistance lessened, the whole circulation becomes freer and more active and changes in metabolism promoted, and a general well-being of the patient is noticed. With the results of the use of electricity some claim good results, but it may be said that electricity seems to be more of an adjuvant than a curative agent. To obtain any measure of success in our treatment all excesses of every kind must be corrected and the general functions of the body made to act properly.

919 McPherson Square.

TUBERCULAR PELVIC PERITONITIS.

BY CHARLES C. ALLISON, M.D.

Professor of Clinical Surgery, Omaha Medical College; Surgeon-in-Chief to the Presbyterian Hospital, etc.

OMAHA, NEB.

It seems to be generally held that the lymphatic system affords the main avenue for tubercular invasion. Thus in children, whose exposed facial integument, with repeated abrasions, invites a tubercular cervical adenolymphangitis,⁽¹⁾ in the adult the nasopharyngeal space, directly exposed to multiple bacterial irritations, favors this predisposition an entrance into the lymph-channels and, with suitable soil, general dissemination through the larger ducts to the bronchial and mediasti-

¹ De Forrest Willard: *Annals of Surgery*.