vibrio, and as none of the 161 examined were carriers, and as the epidemic seemed to have died out, Toulon thought itself safe; but on Nov. 14th a case developed in the *Suffren*, with cholera vibrios in his stools, and he died next day. The situation was complicated by a coincident outbreak of 40 cases of food-poisoning in the *Justice* on Nov. 15th or 16th; these were recognised not to be cholera. But other cases did occur from Nov. 14 to 21st, 18 in all, of whom 6 died. The clinical discussion of these cases appears this month; the epidemiological consideration will be completed in the next number; so we limit ourselves to the clinical aspects of this epidemic, very fully and usefully detailed as they are. Besides the 18 attacked by cholera, 129 persons were isolated, either because they had slight diarrhoea or because they were contacts; 7 had diarrhoea and 6 had vibrios in their stools. Of the 122 healthy contacts, 2 were found to be carriers, and only 2. One of the 18 was a pensioner employed in the dockyard, the other 17 belonged to ships, and a long account is given of each of these cases. Thirteen cases were severe; one was fulminant (the first). Temperature was always subnormal, and perhaps even 7° F. lower in the axilla than in the rectum. The symptoms were, as usual, diarrhoea, rice-water stools, vomiting, cramps, drying of the tissues, wrinkling of the skin, cyanosis, anuria. One man got a double parotitis on the fourteenth day and died a week later with double pneumonia. The treatment which is approved after experience of the whole epidemic is the intravenous injection of saline solution, about 1500 c.c. an hour, or an ounce a minute, during a couple of hours. The veins are very small, the blood quickly coagulates, the patient is particularly restless. Fix the arm then to a splint or the bed, make an incision in the bend of the elbow, isolate the vein, and draw it out, insert the (cannula or) needle, and pour in the saline. The fixation of the arm is the more necessary as the injection continues for two or more hours until the patient improves. It is the conviction of those who saw this epidemic through that no other treatment is to be compared with this. The cases who died had relatively small injections or the large injections were given too late, when other treatments had failed. This paper should be read. Dr. Lepeuple (surgeon in the French navy) reports a case of septicæmic plague at Saigon, where the liver was chiefly attacked, and on section the bacillus of plague was found growing throughout the vessels of that organ. All contacts were injected with Yersin's anti-plague serum. No further cases occurred and no dead rats were found on board.

Correspondence.

"Audi alteram partem."

HÆMATEMESIS DUE TO ARTERIO-SCLEROSIS OF THE GASTRIC VESSELS.

To the Editor of THE LANCET.

SIR,—In your issue of Oct. 12th you have an interesting annotation on hæmatemesis occurring as a very rare complication during the course of arterio-sclerosis.

For the past 30 years I have paid special attention to the clinical aspects of this protean form of arterial degeneration, and from the material that has passed through my hands I am convinced that a local manifestation of the disease process is far more common than is usually considered to be the case. I believe that if as searching an examination were to be made of the smaller arterial twigs as has been done with nerve tissues, abundant evidence would be forthcoming that arterio-sclerosis, commonly looked upon as being of a more or less general arterial change, will often be found as a local patchy disease affecting a very limited area.

My impression is that in at least some of the cases of fulminating pancreatic hæmorrhage it is the focal condition and the potent cause or underlying pathology. We are so accustomed to regard the affection as being a systemic disease affecting the arterioles that we are apt to overlook the fact, supported by clinical evidence, that there are many cases in which the maximum effect is largely local. We

know that the cerebral, ocular, and coronary vessels are constantly affected, and we know that the cerebral and ocular arteries are prone to bleed in a very early stage; naturally hæmorrhage in either of these situations is accompanied by symptoms intensely objective, far more so than if the hæmorrhage occurred in countless other situations, where a minute bleeding would produce only symptoms trifling in character. Personally I have found great difficulty in attempting to differentiate clinically between chronic renal change of that type known as granular kidney in its incipient stages, with well-known tissue changes and slight hæmorrhages, and the more general disease we usually designate arterio-sclerosis, my difficulty being all the greater as I have had on several occasions seen individuals at different stages—extending over many years—of this illness, where at the early commencement one would have labelled the malady as being renal, but where as years have passed by the conditions have been on all fours with those generally considered as due to arterio-sclerosis. So much have I been impressed by this that I take the view that in a large proportion of those cases of so-called chronic renal disease of contracting type, we are in reality dealing with a general arterial degeneration which happens to be concentrated on a certain definite area—to wit, the kidney. Otherwise I fail to explain a clinical fact which has frequently come under my notice—namely, that in some patients the primary disease, or perhaps more properly the symptoms denoting such local disease, may remain quiescent, or entirely in abeyance, or occasionally subsiding altogether, and marked evidences of arterial changes will assert themselves in another position of the arterial tree. We can scarcely overrate the importance of recognising the early local changes of arterio-sclerosis, as it is in the commencement of the disease we obtain so much from treatment.

I am, Sir, yours faithfully,

Green-street, W., Oct. 22nd, 1912.

DONALD W. C. HOOD.

LEGISLATION CONCERNING PROPRIETARY REMEDIES.

To the Editor of THE LANCET.

SIR,—I have read with interest the excellent editorial in THE LANCET of Sept. 28th under the above title. In no spirit of criticism, may I call attention to a slight misconception that the editorial writer has of the Federal Food and Drugs Act of this country? The editorial states that this Act "prohibited extravagant and misleading claims in advertisements and labels." As a matter of fact, the Act has never prohibited any kind of claims that might be made in advertisements. The most outrageous falsehoods may be published in the newspapers regarding the composition, place of origin, or therapeutic efficiency of any patent or proprietary medicine, and the manufacturer is exempt from prosecution under the Food and Drugs Act. What the Act actually does prohibit is false claims that appear on the label regarding composition or source of origin of proprietary remedies.

When first enacted it was generally supposed, and the lower courts so held, that the Act also prohibited false or misleading statements regarding the curative properties of preparations. The United States Supreme Court, however, has decided (in a divided opinion) that the Act did not apply to therapeutic claims. This decision seriously crippled the efficiency of the law for some time. Within the past few weeks an attempt has been made to strengthen the law by adding an amendment that specifically prohibits any statements regarding the therapeutic effect of medicines "which are false and fraudulent."

The editorial further states, "Substances that are considered dangerous are scheduled in a very complete list of prohibited articles in this Act." There is no list of prohibited articles, but there is a list of articles whose presence and quantities must be declared on the label of the preparations containing them. The list, unfortunately, is a small one, comprising "alcohol, morphine, opium, cocaine, heroin, alpha and beta eucain, chloroform, cannabis indica, chloral hydrate, or acetanilid, or any derivative or preparation of any such substances." It will thus be seen that some of the most dangerous poisons can be used as ingredients of patent or proprietary medicines and a purchaser be given no hint as to their presence. Strychnine, arsenic, hydrocyanic acid, and