GEOLOGY OF THE WICHITA MOUNTAINS

BY H. FOSTER BAIN

(Read before the Society December 30, 1899)

CONTENTS

	Page
Introduction.....	127
Previous work in the region.....	128
Physiography.....	130
Rocks present.....	133
Crystalline rocks.....	135
Classification.....	135
Raggedy Mountain gabbro.....	136
Carrollton Mountain porphyry.....	136
Quana granite.....	137
Later eruptives.....	138
Sedimentaries.....	138
Blue Creek series and Rainy Mountain limestone.....	138
Geronimo series.....	140
Red beds.....	141
Cretaceous and Tertiary.....	141
Alluvium.....	141
Age of mountains.....	142
Report on the fossils from the Wichita mountains by Stuart Weller.....	142

INTRODUCTION

For a region of such geologic interest the Wichita Mountain belt has been but little studied. Rising, as these mountains do, through the latter sedimentaries of the prairie region midway between the Rocky mountains and the crumpled paleozoics of western Arkansas, they afford a key to many of the geological problems of the great plains. A detailed study of the region is of the first importance, if we are ever to know the geology of the plains region. While the present study is far from being sufficiently detailed to answer many of these problems, it is believed that important information worthy of permanent record has been obtained. The survey was made within the present year, the month of May being devoted to the task.

In the course of the work the author had the constant and valuable assistance of Mr J. W. Finch, to whom he is indebted for many facts of observations and suggestions of value. He is also in debt to Professor R. D. Salisbury for private notes collected on a reconnoissance trip earlier in the present year. Permission to publish the results is generously given by General Manager H. A. Parker, of the Chicago, Rock Island and Pacific railroad, under whose direction the work was done.

PREVIOUS WORK IN THE REGION

The "Wichitas" are a detached group of mountains of general east-west trend located within the Kiowa and Comanche reservation, in southwestern Oklahoma. They are not within the area open to settlement, and perhaps in part for this reason have long been attractive to the prospector. They have never been geologically surveyed in detail, though several geologists have made reconnoissance trips through the region. The first of these was George G. Shumard, who was attached to Marcy's expedition to the sources of the Red river in 1852.* The specimens collected by this expedition were studied by Edward Hitchcock,† and chemical analyses of the ores and soils were made by C. U. Shepherd.‡ In 1889 Messrs T. B. Comstock and W. F. Cummings, of the Texas Geological Survey, made a trip through the mountains, and their results are given in the First Texas Report.§ This is the only geological paper especially devoted to the Wichitas which we have had up to the present. There have, however, been several brief papers dealing with the Wichitas in connection with neighboring areas. Among them is a paper published by T. Wayland Vaughan last July.|| The latter paper includes petrographic notes by Doctor A. C. Spencer. Robert T. Hill determined the height of mount Scott,¶ and has made several incidental references to the geology of the region.**

Cummings and Comstock devoted more time to the area than any previous investigators, but their opinions seem to have been very largely colored by what they had previously seen in central Texas. The present work has shown that they are fundamentally wrong in referring the granite to the pre-Cambrian, since it cuts and metamorphoses Ordovician strata. They are also clearly mistaken in indorsing Hitchcock's †† opin-

* Senate Doc., 2d sess., 31st Congress, vol. 8, Washington, 1853, Appendix D.

† Op. cit.

‡ Op. cit., Appendix C.

§ Geol. Survey Texas, vol. i, pp. 319-328.

|| Amer. Geologist, vol. xxiv, 1899, pp. 44-57.

¶ Ibid., vol. vii, 1891, p. 119.

** Ibid., vol. vii, p. 254; vol. vi, pp. 252-253. Amer. Jour. Sci., (3), vol. xlii, 1891, pp. 122-123.

†† Geol. Survey Texas, vol. i, p. 321.

FIGURE 1.—Sketch Map of Wichita Mountains

H. Foster Bain

ion that there has been recent volcanic activity in the region. The greenstone dikes are certainly the most recent igneous rocks. The tuffs and rhyolites found are pre-Cambrian, and the resemblance of certain of the mountains to "an ancient crater" is fanciful only. It has been impossible to recognize all the peaks named by this party and to perpetuate these names. Wherever the recognition seemed secure the names have been used. Mount Webster, located and ascended by Marcy's party, seems now unknown. From the general references to its position and the sketch given of its outline it seems probable that it is the peak now known as Baker, but in the absence of accurate maps there will for some time be confusion regarding the geography of the area between Saddle mountain and the Mesquite valley.

The maps of the region are quite imperfect. The land survey seems to have been more than usually inexact. The sketch map (figure 1) given with this paper is from a blue print of a map made by the railway engineers. The position of the various mountains is set down from checked compass readings, where other data are lacking.

PHYSIOGRAPHY

The Wichitas rise abruptly from the great prairie plain about midway between the Washita and Red rivers. Cache creek flows through the low, circling hills of the east end, and on the west the North fork of Red river separates the Wichitas from the Navajo mountains which lie in Greer county, Oklahoma.

The core of the mountains is a rugged mass of igneous rocks. On the north side, circling around the east as far west as the south side of Signal mountain, is a lower range of limestone hills of less rugged aspect and with rounded slopes. These hills rise 70 to 400 feet above the plain and are in part detached, such as Rainy mountain and the elevation near it, and in part massed together. These limestone hills are separated on the north from the main mountains by a wide valley, down which for much of the way Medicine Bluff creek flows. This stream, which by Marcy's party was evidently mistaken for the main branch of Cache creek, cuts through the porphyry of Carrollton mountains, forming a sharp bluff 250 to 300 feet high, from which the stream takes its name and which is regarded with a certain amount of awe by the Indians.

The mountains themselves consist of a series of detached and semi-detached mountains, and are crossed by wide passes which are nearly as level as the surrounding plain. Between the mountains flat grassy plains are found and spurs protrude through the plain at some distance from the main mountain mass. The exceedingly rugged topography of

the mountains proper and the great detached blocks of granite which cover their surface give as a result the appearance of a deeply buried range with the tops just protruding through the plain. In a way this is a true conception of the case.

The northernmost series of hills consists mainly of limestone, but includes some porphyry, granite, and gabbro. In the vicinity of Rainy mountain they are made up of a series of detached and rounded limestone hills rising out of the prairie 70 to 400 feet and with a general dip north. East of this region, and extending from north of Saddle mountain to a little northeast of mount Scott, the hills are traversed by no pass

FIGURE 2.—Sketch of Lower Narrows.

Showing sharp granite hills with flat plains between.

and consist of a complicated, much folded and faulted mass of limestone, with porphyry, granite, and a little gabbro. This series of hills stretches north, with a triangular outline culminating about 10 miles southeast of Mountain view. The pass north from mount Scott is traversed by the Fort Sill-Cheyenne trail, which follows the narrow canyon of Blue creek, separating the range just described from another limestone range. The latter shows a complicated structure near the pass, becoming simpler to the northeast, where the limestone crops out on Lime and Chandler creeks. To the north the hills swing round, crescent-like, to meet the point of the limestone triangle already noted.

South of the series of limestone hills, and separating them from the first of the granite ranges, is the broad open valley occupied in part by

Medicine Bluff creek, 5 to 7 miles wide, and trending with the mountains. The granites immediately south of this valley form the north range of the mountains proper. They extend on the east almost to Cache creek in a series of detached knobs, rising to the west and culminating in mount Cummings. The latter is separated from mount Scott, the highest peak in the mountains, 2,305 feet above sealevel and about 1,500 feet above its base, by an open pass. Mount Scott extends for about 5 miles to the west with decreasing altitude, and is separated from the next peak, mount Sheridan, by a narrower, higher pass. From mount Sheridan west to Haystack mountain, which lies a little east of south from Rainy moun-

FIGURE 3.—Sketch of Lower Narrows in Raggedy Mountains.

Showing semi-detached character of the peaks.

tain and is in the same locality as Tymatee mountain, as located by Vaughan, there are no passes proper. About midway the granite runs out to the north nearly 4 miles and culminates in the striking peak called Saddle mountain. From Haystack west nearly 10 miles is a broad mesquite plain, extending south across the entire mountains and separating the main Wichitas from a series of detached and irregularly placed mountains extending to the Red river and known locally as the Raggedy mountains.

South of mount Scott and the granite range of which it is the most conspicuous feature is an open valley 3 to 5 miles across, broken by irregular granite and porphyry knobs rising 150 to 400 feet above the general level of the plain and closing in to the west. Signal mountain

and the range of porphyry hills, including Medicine bluff, and called the Charlton mountains by Comstock, pierce this valley from the southeast with a series of elevations of decreasing height and extending as far west as the longitude of mount Sheridan. South of the porphyry hills is the main granite mass of the south range. This begins in an outlying bunch of hills east of Beaver Creek pass and extend unbroken westward to the broad pass north of Quana Parkers, a Comanche chief living on West Cache creek. About 2 miles east of Quanas the range throws out a long low spur to the south. West from Quanas the mountains are unbroken to the Mesquite valley, already described. The intermontane valley is more and more broken by mountains to the west until it becomes too obscure for recognition, and the mountains face the Mesquite valley along the eastern edge in a practically unbroken range.

West of Mesquite valley and extending to the Red river is a group of detached peaks known locally as the Raggedy mountains. Otter creek flows through them and at two points on its course cuts gorges known as the Upper and Lower narrows. The former is located in township 4 north, range 17 west, and latter in the southern portion of the first township south of this. Iron mountain, as located by Vaughan, is one of the peaks lying northeast of the Upper narrows, and Round mountain would lie west of them, in the region where there has been so much surreptitious prospecting.

ROCKS PRESENT

The rocks forming the Wichita mountains include granites, gabbros, and porphyries in their various phases, and a series of Cambrian and Ordovician sediments, principally limestones. The oldest rocks in the region are the Raggedy Mountain gabbro and the Carrollton Mountain porphyry. The relations of the two are not certain beyond dispute, but apparently the gabbro is the older. In a general way the gabbro is more prominent in the western portion of the mountains, being especially well developed in the Raggedy mountains, while the porphyry is more common in the east. Both are pre-Cambrian and perhaps Archean. These rocks formed an old land-mass or island around which the sediments were laid down. The latter begin with a conglomerate carrying fragments of porphyry and gabbro, but none of granite, and run up through the quartzose sandstone, calcareous sandstone, and arenaceous limestones of the Blue Creek series, and so into the Rainy Mountain limestone proper. At its base the series carries definite Cambrian fossils, and higher up Ordovician (Lower Silurian) fossils are found. Allowing for dip and faults, a thickness of a trifle over 4,000 feet of these beds is exposed.

They are best displayed on the north side of the mountains, but sweep around the eastern end, and are found so far west as Signal mountain on the south side. Beyond this point they seem to be buried beneath later material.

The Quana granite, so named from a prominent chief, near whose lodge it is well exposed, is eruptive, cutting the porphyry and the gabbro at many points and exhibiting all the usual contact phenomena. A great dike of granite running from mount Scott north has disturbed and metamorphosed both the porphyry and the limestone, proving that the granite is later in age than either. Additional proof is found in certain

FIGURE 4.—General View of Granite Hills of the South Range near Quanas.
Looking south.

beds, called here the Geronimo series, east of Cache creek, near Fort Sill. In some mesas nearly east of Fort Sill agency, there is a series of sandy shales capped by limestone. These are east of the Fort Sill limestone quarries and from the dip presumably represent a higher horizon. The shales include a conglomerate bed which shows fragments of the pre-existing limestone, porphyry, and granite. They belong accordingly to a series later than the Ordovician, already described. The age of the series is not certain, but its appearance suggests Paleozoic.

There is a series of dikes of fine-grained greenstone of as yet undetermined character, which cuts granite, gabbro, and porphyry alike.

There are also quartz veins in the granite and porphyry and in the gabbro in connection with granite dikes.

Recapitulating, the various rocks found in the mountains are shown in the following table:

Formation.	Character of the rocks.	Age.
Red beds.	Sandstone and shales.	Permian.
Geronimo series.	Sandstone conglomerate, shales, and limestones; carries asphaltum.	? ?
Dike rocks.	Fine-grained greenstones of various types.	May be later.
Quartz veins.	Cut by dikes.	May be in part earlier.
Quana granite.	Eruptive through earlier formations and furnishing debris to the Geronimo series.	
Rainy Mountain limestone.	Blue and gray limestone folded and metamorphosed by the granite; conformable on the earlier sediments.	Ordovician.
Blue Creek series.	Conglomerates, quartzites, sandstones, running up without break into the Rainy Mountain limestone and unconformable on the earlier rocks.	Cambrian.
Carrollton Mountain porphyry.	Including rhyolites, amygdaloids, and some tuffs. The Saddle Mountain porphyry, of slightly different type, probably belongs here.	Archean (?)
Raggedy Mountain gabbro.	Including a considerable variety of basic rocks here regarded as facies of one magma. Possibly eruptive through the porphyry, but not certainly so.	Archean (?)

CRYSTALLINE ROCKS

CLASSIFICATION

As has already been stated, the central core of the mountains consists of crystalline rocks. These include three well marked and usually easily distinguished types: (a) granite, (b) porphyry, (c) gabbro. There are also quartz veins and greenstone dikes. There has as yet been no opportunity to make microscopic studies of these rocks, but there is no reason to believe them greatly different from the rocks examined by Spencer*

* Amer. Geologist, July, 1899, pp. 47, 48.

and collected in the same region. What is said here is based on macroscopic observation only.

RAGGEDY MOUNTAIN GABBRO

With the gabbro are classed here a considerable variety of dark basic rocks, some of which, occurring in dikes, may prove to belong rather with the diabases. The normal type is a dark holocrystalline rock showing pyroxene and labradorite. West of Otter creek the rock shows frequently great masses 6 to 8 inches across of pyroxene. Ilmenite is not uncommon, and has been mistaken for tin ore. The gabbro areas are characterized by low, almost flat surfaces (see plate 15, figure 1). The rock apparently weathers down rapidly, and east of Mesquite valley nowhere rises in sharp hills. It occurs along Medicine Bluff creek on the north flank of mount Scott, where it is coarse grained and apparently properly an anorthosite, and in the embayment between mount Sheridan and Saddle mountain. It is found in limited areas north of Medicine Bluff creek, rising from beneath the limestone. It was noted also in what has been called the central valley of the mountains, and is abundant along the west flank of the mountains facing Mesquite valley. Low knobs of this rock project through the floor of the valley, and near the Upper Narrows it forms hills rising 500 feet along the creek. The relations of the granite to the main gabbro mass is beyond dispute. Northwest of mount Scott the granite was seen to cut the gabbro at two points. Along the west flank of the mountains granite dikes running out into the gabbro are quite common. On Otter creek, about half way between the Upper and Lower narrows a boss of granite rises through a sheet of gabbro and sends out stringers into the latter on all sides. Figure 2 of plate 15 represents a horizontal granite dike cutting the gabbro of one of the high hills west of Otter creek. The presence of a small stringer of granite running off from the dike and a sharp covered block of gabbro imbedded in the granite is conclusive.

•The relations of the gabbro to the porphyry and associated rocks is not so clear. In the Blue Creek canyon a gabbro area was noted wholly within the porphyry and apparently under it. Near the same area a dike of gabbro cuts the porphyry. The gabbro, as well as the porphyry, has furnished material to the Cambrian basal conglomerates. Not all, however, of these basic rocks belong with the main gabbro mass, since in the Carrollton mountains a quartz vein cutting the porphyry is in turn cut by a greenstone dike.

CARROLLTON MOUNTAIN PORPHYRY

There are two types of porphyry found in the region. The first has a fine pink ground-mass, is set with phenocrysts of orthoclase and clear

FIGURE 1.—TYPICAL GABBRO SURFACE
Showing low, rounded boulders

FIGURE 2.—GRANITE DIKE IN GABBRO
Showing stringer and horse

GABBRO SURFACE AND GRANITE DIKE

FIGURE 1.—CHARACTERISTIC VIEW OF GRANITE TALUS ON SIDE OF MOUNT SCOTT

FIGURE 2.—GRANITE BOULDER ON TABLELAND SOUTH OF MOUNT SCOTT

Illustrating tendency to form large surface boulders

GRANITE TALUS AND GRANITE BOULDER

quartz. The latter is especially common in the porphyry of the Carrollton mountains. In general, the porphyry does not take on a granitic phase, though a few such instances were noted. In Blue Creek canyon there are associated with the porphyry certain euryolites, showing lithophysæ and other rocks apparently rhyolitic and tuffaceous. It is this pink porphyry which forms the common type of the mountains and is seen south and east of mount Scott, in the Carrollton mountains, in Signal mountain, and in the limestone hills north of the main range, where it forms the platform upon which the Cambrian conglomerate rests and has been brought up by faulting. The second type of porphyry was noted only along the north flank of the main granite range west of Saddle mountain. It differs from the usual type in having a much darker ground-mass, which may be a wholly inconsequential difference.

Wherever the porphyry outcrops it shows a characteristic topography, forming rounded hills with smooth flowing contours. It is often much shattered, and the surface of such hills usually shows many small sharp-edged fragments of rock, as distinguished from the granite and gabbro outcrops, which are characterized by rounded boulders of disintegration.

QUANA GRANITE

This is by far the most common crystalline rock in the region, and shows but little variation in character. The rock is predominantly feldspathic, with a subordinate amount of quartz and a still smaller amount of a green mineral, presumably hornblende. So far as was observed, the granite is wholly free from mica, and in fact mica was only found in two localities, where it occurs in small yellow flakes in connection with certain dikes of granular white quartz radiating from the granite mass and cutting the gabbro. Granite forms the bulk of mounts Scott, Sheridan, Baker, Haystack, and Saddle mountain, and in fact all the more prominent peaks. It weathers characteristically into large boulders marked off by joint cracks. On the top of a large tabular mountain just east of the Mesquite valley one such boulder of disintegration was seen, measuring 60 by 40 by 30 feet. Figure 1 of plate 16 shows a characteristic view on the side of mount Scott, and figure 2 of the same plate gives one a correct impression of the size of the blocks on the lowland. These boulders are so common that over much of the granite area it is difficult to find exposures of the solid rock.

The granite has suffered much deformation. Shear zones are not uncommon, and faulting with well developed slickensides was repeatedly observed. The rock is eruptive through the gabbro, as already shown, and its relations to the porphyry are quite as clear. At contacts observed on the southwest flank of mount Scott the granite was seen to

send off stringers into the porphyry. It becomes, too, notably finer in grain, while the porphyry is not changed in general character, though it is badly shattered near granite outcrops. In Blue Creek canyon the porphyry platform upon which the Cambrian conglomerate was deposited has been sharply tilted by the granitic intrusion. The relations of the granite to the limestone are clear from the three facts: (a) At an actual contact observed on the west side of the Blue Creek canyon the dark earthy limestone had been recrystallized into a white, coarse marble; (b) in the same vicinity the intrusion of the granite had tilted the porphyry and its covering of sedimentaries; (c) the basal conglomerate contains everywhere fragments of both porphyry and gabbro, but no granite, though the latter could not have been more difficult of access, assuming it to have been pre-Cambrian.

It is only in the later conglomerates of the Geronimo series that granite pebbles appear.

LATER ERUPTIVES

At several points the granite is cut by greenstone dikes. At one point in the Raggedy mountains three generations of the greenstone, including the original gabbro, can be counted. Quartz veins, some of them granular and carrying mica, also cut the granite. The best known of these veins is the one separating the easternmost from the middle porphyry hill at Medicine bluff, in the Carrollton mountains. This vein is 12 feet wide and shows plainly on both sides of Medicine Bluff creek. It is cut by a small greenstone dike and was noted by Marcy.

SEDIMENTARIES

BLUE CREEK SERIES AND RAINY MOUNTAIN LIMESTONE

The character and relations of the sedimentary series can perhaps be best made out from the exposures around Canyon Creek camp, which was located on the Fort Sill-Cheyenne trail about 7 miles north of mount

FIGURE 5.—Crushed Anticline and Overthrust Fault in Limestone, north of Mount Scott.

Scott and in the canyon cut by Blue creek. The trail enters the limestone hills from the south and runs for some distance on granite, believed to be a great dike running off from the Mount Scott mass. To the west is a great complex of highly contorted limestone, changing into marble at the contact with the granite, and extending with many conflicting dips nearly to Rainy mountain. Figures 1

FIGURE 1.—CONTORTED LIMESTONE WEST OF FORT SILL—CHEYENNE TRAIL NORTH OF MOUNT SCOTT

FIGURE 2.—TRUNCATED PLUNGING ANTICLINE IN ORDOVICIAN STRATA NORTH OF MOUNT SCOTT

CONTORTED LIMESTONE AND TRUNCATED ANTICLINE

and 2 of plate 17 will show how sharply the limestone is folded, and figure 5 is a rough sketch of a crushed anticline and overthrust fault chosen as one of many possible illustrations. There are anticlines with almost vertical pitch, and the rocks have evidently suffered very much

FIGURE 6.—Sedimentaries: Blue Creek Series resting on Porphyry, north of Mount Scott.

disturbance. It may be tentatively suggested, though the data collected hardly warrant any generalization, that the rocks seem to have been first thrown into a series of east-west folds parallel to the main mountain range, and then thrown into cross-folds by the intrusion of the local granite mass.

To the east of the trail are high porphyry hills covered by a sedimentary sequence having a fairly uniform strike and dip. Near the mouth of the canyon a fault repeats the sequence as if to enable the geologist to check

FIGURE 7.—Repetition of Cambrian Strata by Faulting, north of Mount Scott.

his work. Figure 6 represents the basal portion of the section as far as the fault. In figure 7 the fault and repetition are shown. The general section of the beds from the base up is as follows :

	Feet
1. Porphyry, red, usual type.	
2. Sandstone, white, rounded grains, quartzitic, with one small dip fault; dip, 28 degrees.....	50
3. Conglomerate, dark colored, porphyry pebbles set in matrix of green sand seemingly derived from disintegration of basic rocks; Cambrian fossils; dip, 30 degrees.	250
4. Coarse grained crystalline limestone with particles of magnetite and horn- blende; dip, 29 degrees.....	30
5. Dark sandstone, conglomeratic, with porphyry pebbles and much horn- blende; dip, 24 degrees.....	180
6. White calcareous sandstone weathering cavernous.....	70
7. Limestone, blue, with fossils (Cambrian) ranging from 40 to 50 feet near the middle; dip, 29 degrees at base, becoming 5 degrees at top.....	110
Fault repeating first seven numbers.	

	Feet
8. Limestone, thin shaly, with some sandy material.....	50
9. Limestone, blue, heavily bedded.....	450
10. Limestone, thin bedded, sandy.. . . .	80
11. Limestone, heavy, blue; dip, 35 degrees.....	510
12. Limestone, brecciated, rounded, and angular; fragments imbedded in calcareous matrix; no evidence of especial crushing.....	10
13. Limestone, blue, earthy, with chert for a thickness of 100 feet 600 feet above base, and fossils near top.....	1,250

It is possible that numbers 10 and 11 are repetitions of 8 and 9, due to an obscure dip fault, but it was impossible to determine this absolutely. Number 13 seems to represent the same horizon as that from which numerous fossils were collected in the hills east of Rainy mountain, and would accordingly be *Calciferosus*. According to this determination the section was not at this point carried up to the Trenton. A total thickness of more than 4,000 feet was measured in this vicinity, and it is doubted whether the combined Cambrian and Ordovician much exceed this thickness, though in view of the changes in dip and liability to faults the question must for the present remain open.

At Rainy mountain the lower beds are not shown, and the Trenton only is exposed. The *Calciferosus* is exposed in the group of detached hills east of the school, and on the plain between these hills and the mountains there is a suggestion of the basal conglomerate at the proper horizon. These basal beds are shown again at the granite and porphyry hills, about 10 miles southeast of Mountain view, and at the east end of the Carrollton mountains, just west of the rifle range at Fort Sill, as well as on the south side of Signal mountain. No trace of any sedimentary rocks was found on the south side of the mountains west of this point.

*GERONIMO SERIES**

Rocks later than the Trenton were only seen southeast of Fort Sill. East of Cache creek and near the southeastern corner of the military reservation are certain limestone mesas known because of the presence of a number of asphaltum springs. This series of rocks include much shale and sandy limestone. It is characterized by low dips and extends some distance east of Fort Sill, on the Marlow trail. Near the base of the series the shales are interbedded with a conglomerate having a calcareous matrix and carrying large, 2-inch, rounded pebbles of granite, porphyry, and limestone. Unfortunately, no recognizable fossils were obtained from either the shales or the overlying limestones, and hence

*In the absence of any more available name, this series is called after the famous Apache chief, who now lives as a prisoner of war at Fort Sill.

all evidence as to the age of the beds must be derived from the stratigraphy. They are manifestly later than the granite intrusion, and in composition and character resemble the known Paleozoic rocks of the region. They are earlier than the Red beds (Permian), as they pass under the latter. It is to be hoped that further study may be devoted to them.

RED BEDS

The sandstones and shales of the Permian cover much of the prairie between the Washita river and the mountains. They were observed at many points in the course of the present work, but nothing with regard to them can be added to previous knowledge. They were nowhere seen to have been involved in the dynamic movements which have given a slight dip even to the Geronimo beds. They were seen quite near the mountains and wholly undisturbed. It would not be strange if at some point escaping observation they showed a certain amount of disturbance which might be traced to relatively recent and slight movements.

CRETACEOUS AND TERTIARY

Mr Robert T. Hill reports the former entirely absent from the mountains, and this accords entirely with our own observations. Comstock has assigned to the latter, under the name of Fort Sill beds, certain of the surface materials covering the intermontane pastures. There seems no good reason for this. The material is, as Vaughan remarks, lithologically similar to that which makes up the Red beds. Community of genesis probably accounts for this. The outwash from the mountains would be of much the same character, regardless of the time when it was deposited, and, while the matter may in its present condition be fully conceded to be one mainly of opinion, the author is inclined to treat these beds as of recent origin.

ALLUVIUM

Whatever may be the ultimate decision in regard to the beds just mentioned, it is perfectly certain that there is along the south side of the mountains a vast quantity of material which can not be considered to be other than recent. On West Cache creek beyond Quanas there are great beds of sand and recent conglomerate rising 50 feet above the present stream. The passes in the south range of mountains show great alluvial fans, which curiously enough in one or two instances seem to have choked the old passes and led to the capture of the headwaters of intermontane creeks by younger streams, which have cut new and sharp walled canyons

through the south granite range. This outwash is far heavier on the south than the north side of the mountains, and boulders 6 to 8 inches in diameter were quite frequently found as much as 15 miles south of the mountains in such quantity as to preclude any hypothesis of their adventitious character. The suggestion is that there has been a marked differential rise to the north, and that any early limestone hills on the south have been buried by Permian and recent outwash.

AGE OF THE MOUNTAINS

The preceding sketch of the geology has perhaps shown clearly enough what the conclusion must be as to the age of the mountains, but possibly a word or two of summary may not be out of place. It is evident that there was a pre-Cambrian land-mass of igneous rocks, and that over this was laid down an undisturbed sequence stretching from the Cambrian up to and including the Trenton. Then came the main upheaval and the intrusion of the granite. Around the edge of the new mountains the Geronimo series was laid down. The shearing and faulting of the granite and the presence of greenstone dikes cutting it, with the true though slight dip of the Geronimo beds, indicate later disturbances of lesser degree. Since the intrusion of the granite, however, the main history of the region has been one of vigorous and long continued erosion, through which the mechanical sediments of the Red beds and later deposits were prepared and distributed. That there was a notable interval of erosion between the intrusion of the granite and the Geronimo beds is shown by the presence of granite pebbles in the latter. Granite being intrusive, these pebbles could only be obtained after the erosion of a considerable amount of rock.

REPORT ON THE FOSSILS FROM THE WICHITA MOUNTAINS BY STUART WELLER

The following determinations of the geologic horizons in the Wichita mountains have been made from a small collection of fossils submitted to me by Mr H. F. Bain, supplemented by a small collection secured by Professor R. D. Salisbury. The fossils are all poorly preserved and in almost every instance are imperfect or fragmentary, so that definite identification of the species is in most cases not practicable. They indicate, however, the presence of three distinct horizons, the lowest of which may be referred to the Cambrian and the two upper ones to the Ordovician.

Most of the material identified as Cambrian was collected at "Canyon Creek camp."* The species represented are as follows:

1. *Dikellocephalus*? sp. This is the most abundant species in the fauna and is placed in this genus with much hesitation. The glabella is usually all that is preserved. This portion of the head is rather strongly convex, subquadrangular in outline, with the lateral margins nearly parallel or slightly diverging anteriorly. The occipital furrow is strongly defined, with one much fainter glabella furrow situated at about one-third the distance from the occipital furrow to the anterior extremity of the head. The fixed cheeks are rather broad and moderately convex. In the largest and best preserved specimen the glabella is 19 millimeters long and 15 millimeters wide. The smallest specimen is 6 millimeters long and $4\frac{1}{2}$ millimeters wide.

2. Free cheek of a trilobite, possibly belonging to the species referred to *Dikellocephalus*.

3. *Ptychoparia* sp. A single imperfect head shield.

4. Undetermined coiled gasteropod shell with an elaborate spire.

5. *Oboella*? sp. Two fragments.

A few specimens bearing the label "Fossils from conglomerate in sandstone east of granite dike"† also prove to be Cambrian, the species recognized being as follows:

1. *Ptychoparia*, sp. undetermined. A single well preserved cephalon.

2. Fragments of several trilobites too imperfectly preserved for identification.

All of these fossils would seem to indicate an Upper Cambrian horizon.

The fossils of the second horizon, the Calciferous of the Lower Ordovician, are for the most part from the limestone hills east of Rainy mountain. They were collected from several localities, but as they all seem to belong to a single general horizon, they will be listed all together.‡

1. Pygidium of trilobite. *Bathyurus*? sp.

2. Pygidium of undetermined trilobite.

3. *Cyrtoceras*, 1 or 2 species.

4. *Hormotoma* sp.

5. *Lophospira cassina*, Whitf.?

6. One or two species of coiled gasteropods like *Holopea*.

7. *Raphistoma*? *trochiscus*, Meek?

8. *Ophileta complanata* var.?

9. *Bucania* sp.

10. Two or three additional species of coiled gasteropods represented by fragments.

11. *Orthis* or *Strophomena* sp. Several undeterminate fragments belonging to one or the other of these genera.

* Located in Blue Creek canyon, about 7 miles due north of mount Scott.

† Number 2 Blue Creek section.

‡ Vaughan (Amer. Geologist, July, 1899, p. 49) collected *Raphistoma* and *Ophileta complanata* var. *nana*, as determined by Girty from this general locality.—H. F. B.

From the limestone of the Fort Sill quarry a few species were secured, as follows:*

1. *Ophileta* sp.
2. Fragments of trilobites, genus and species undetermined.

While all these fossils are in a very unsatisfactory condition for determination, there seems to be no doubt of their Calciferous or Chazy age.

The third horizon is that of the Trenton limestone. The only locality from which Trenton fossils have been examined is Rainy mountain. A few of these were submitted by Mr Bain, but most of them were collected by Professor Salisbury. The species recognized are as follows:

1. *Ceraurus*, sp. undetermined.
2. *Isotelus gigas* De Kay or *megistos* Locke. A single imperfect thoracic segment.
3. *Trinucleus concentricus* Eaton.
4. *Rhynchonella* sp.
5. *Orthis* (*Dalmanella*) *testudinaria* Dal.
6. *Rafinesquina alternata* Emm.
7. *Lingula rectilateralis* Emm.

* From Fort Sill Vaughan reports *Paleophycus*, *Murchisonia*, *Cyrtoceras*, *Linguella* cf. *L. lamborni*, and *Asaphus*, from which it is suggested that the beds may be as late as the Trenton.—H. F. B.