cases a revulsive bleeding brings on flooding. I have lately seen an example of the latter, produced by an abstraction of four ounces of blood. These are undoubtedly rare exceptions, but we should not lose sight of them. In such cases, general, and, *a fortiori*, local, bleeding should be avoided. Some females, essentially nervous, cannot support bleeding, without suffering various unpleasant nervous accidents. Here the quantity of blood taken away may be diminished to a very few ounces, and this precaution sometimes succeeds; but should the same accidents occur, we must give up the idea of bleeding altogether. If we except these cases, venesection, aided by narcotics, is the surest means of calming and removing the pain. This idea is not new. Stahl had long ago remarked, that in cases of cancer, the patient experienced a marked amelioration whenever the veins gave way. You have often seen me, gentlemen, prescribe bleeding to twenty women at the same time, in this hospital; fifteen of them, at least, are benefited; the pains disappear for a longer or shorter period, unless some foreign influence, as a moral affection, or a change of atmosphere, interrupt the action of the remedy. In this very ward we have supported a woman for ten years who was affected with a very dangerous uterine disease, not so much by employing narcotics as by repeated small bleedings, varying from two to six ounces, according to the strength of the patient. When the disorganization of the uterus is very far advanced, and accompanied by the discharge of cancerous matter, the quantity of blood drawn should be very small, in order to avoid the danger of facilitating its absorption. If the patient be very feeble, if we remark any symptoms of indifference, a tendency to sleep, or stupidity, in a word, if adynamia exist, we must avoid employing venesection altogether, as a means which can only hasten the fate of the patient.

The treatment with cupping-glasses, blisters, &c., must for the present be deferred.

AMPUTATION IN AN INFANT.

To the Editor of THE LANCET.

SIR,—The subjoined case has been handed to me by a Member of the Council of the Royal College of Surgeons in London, with a view to its publication in *THE LANCET*. It was originally intended for publication by the College, but as this course has not been deemed expedient, the gentleman in

question is desirous that the utmost publicity may be given to the case through the medium of "*The leading Medical Journal*."

The case (an amputation of the thigh at nine weeks) possesses its peculiar interest from the early age of the patient operated on, and I am informed no member of the college senate is cognizant of a similarly severe operation performed on one so young, not even the veteran Sir Astley, nor that patriarchal English surgeon the venerable Sir William Blizard. I am, Sir, your obedient servant,

GEORGE FREDERICK KNOX.

14, George-street, Adelphi,
Dec. 9, 1833.

CASE OF CONGENITAL FUNGUS HÆMATODES, IN WHICH

AMPUTATION OF THE THIGH

WAS PERFORMED IN THE TENTH WEEK
OF THE CHILD'S LIFE.

With Remarks.

By JOHN PAUL, M.D., Surgeon to Gray's Hospital, Elgin.

A MALE child, seven weeks old, was brought to me by his parents in the month of September, with an enormous swelling of the right leg, which had all the characteristic marks of fungus hæmatodes. The swelling was soft and elastic, bulged out in various parts, and presented a livid hue at the most depending point, with enlarged cutaneous veins. At birth two tumours were observable, the one running into the other; the lower one was said to be about the size of a turkey's egg, and the other somewhat less. The child's health was tolerably good.

The whole leg being involved in the morbid action of the disease, nothing but amputation above the knee even required the consideration of a moment, and the patient being so young this alternative was deemed almost too desperate; at least some delay was thought prudent. The parents were therefore advised to take their child home, and return when the swelling burst.

They had not been at home many days when the swelling did burst, and so profuse was the hemorrhage that in less than one minute the infant was in a state of syncope, and for two days life appeared to be almost extinct, so much were the vital endowments depressed. The hemorrhage was controlled by pressure, and the little patient rallied after a few days.

He was admitted into *Gray's Hospital*,