shaken, no small part of the power of treatment for good he believed to be lost.

For a number of years past the doctor was afflicted with a degree of deafness precluding anything like satisfactory auscultation. It was remarkable how little this circumstance lessened his confidence in diagnosing diseases of the chest. He trusted entirely to acute observation, and shrewd interpretation of the ordinary symptoms.

The doctor looked upon the placenta blocking the os uteri or vagina after labor as a natural and most useful tampon, and he regarded too early an interference with it as a mischievous interruption of nature in her efforts to close and occlude the uterine vessels. Naturally he agreed with Braithwaite, that in certain conditions of hæmorrhage after labor, the uterus being contracted, to tampon was good practice; and he went so far as to suggest that probably in such cases the re-introduced placenta might be the best plug that the accoucheur could employ. The doctor had for nearly forty years of his practice an unusual number of cases of post-partum hæmorrhage. He had meantime conducted the third stage of labor according to the standard precepts of the day. Of late years he had fewer cases of hæmorrhage, and he referred the change in his experience to a change in his practice. As an almost invariable rule he had during the first decades of his work given ergot immediately after the birth of the child. It was following a discontinuance of this practice that the comparative freedom from post-partum bleeding occurred.

In placenta prævia centralis Dr. Groesbeck believed that perforation of placenta gave as good results as the passing of the hand between it and the uterus, and he regarded the former method as easier, and more speedy than the latter. When under any circumstances turning had been resorted to, Dr. Groesbeck delivered promptly, impressing on the child the necessary turns, and never waiting for nature to do so. In breech presentations the doctor thought it better practice to break the wedge as soon as the os was sufficiently dilated, bringing down both feet and delivering at once.

The doctor was an advocate of the rather frequent use of the forceps, using the old model Bedford for higher operations. When the head was lower in the pelvis he used the short forceps, always speaking highly of their convenience and efficiency. It should be remarked, however, that in the selection of the short instrument, Dr. Groesbeck rejected those in the construction of which the maker, in shortening the handles, shortened also the cephalic curve.

Dr. Groesbeck fully endorsed the views of the anatomy of the cervix uteri gravidi held by Braune, Bandl, Barnes and others. He believed that the placenta prævia was located below the os internum.

The doctor was possessed of an unusual degree of candor and was entirely void of professional pretense. He often admitted his shortcomings; thus, he did not hesitate to state that the mechanism of labor as regards the mutations of the foetal head in passing the parturient canal were to him inexplicable. He declared that he was generally unable to determine the position of the head, and that the reputed

"planes and spines" effecting changes in it were to him a "terra incognita."

His favorite obstetrical work was "The Principles of Midwifery, by John Burns." In the preference here indicated our colleague complimented his own judgment as well as the classical production of the Glasgow professor.

The doctor had a keen appreciation of the ways and means which tact and experience teach as aids to success in practice. In a knowledge of the subtle methods by which a physician may ingratiate himself with the people, magnify his importance, and obtain and retain practice, Dr. Groesbeck was not second to the author of "The Physician Himself." But while he was prodigal of advice in such matters to others, he took, in his own conduct, little heed of his own wisdom.

After he had been in the profession half a century he felt less confident that he would in any case do all that was possible for a patient than he had felt fifty years before. He on one occasion expressed surprise at the recollection of the nonchalant unconcern with which as a novice he had undertaken the management of an adherent placenta or placenta prævia.

As to the profession as a business, Dr. Groesbeck held an opinion contrary to that of the great mass of practitioners. He always insisted that the practice of medicine was upon the whole as easy and as pleasant a mode of earning a livelihood as any of which he had knowledge.

Dr. Groesbeck felt a fatherly interest in the younger members of the profession, and never tired of stimulating their thought and enlarging their knowledge by the narration of instructive cases and the enunciation of cardinal precepts.

Mr. President: From the shores of the mystical river we call to our vanishing friend the final word of mortals: Vale! Vale! An adieu laden with uncertainty, yet buoyant with hope. As we turn from the dark waters to the duties yet before us, let us pause to inscribe our index memorabilium with the honorable name of Abram Groesbeck.

FOREIGN CORRESPONDENCE.

COCAINE.

WIESBADEN, GERMANY, Dec. 1, 1884.

Mr. Editor:

E. Merck, the celebrated pharmacist of Darmstadt, has kindly furnished me with the following items in regard to the now famous remedy—cocaine.

The chemical formula according to Lossen is: $C_{17}H_{24}NO_4$. Cocaine is the alkaloid contained in the coca leaves (*Erythoxylon Coca Lam*) which was first isolated by Niemann. In 1862, Lossen discovered a second principle contained in the same leaves, viz.: hygein, which is of a volatile nature, and has received but little investigation so far; it seems to have a weak and hardly characteristic action. The leaves further contain ecgonin, coca tannin and a peculiar wax.

The coca crystals belong to the monocline system, melt at $98^{\circ}C.$, easily soluble in alcohol, more easily

in ether, but only dissolve in 704 parts water. The cocaine *salts*, however, are easily soluble in water.

The first reports of the internal use of coca leaves are handed down from the sixteenth century (Dr. Monardes, Seville, 1569). Its first appearance in Europe dates from 1749—it was described by Jussieu, and was named *erythoxylon coca* by Lamarck. Tschudi, Markham, Poppig and others who travelled in South America, found that the natives were in the habit of chewing coca leaves as a remedy for or preventive against the effects of extraordinary exertions. It seems, however, that the Indians chewed the leaves, together with the ashes of *chenopodium*. Quinea, the alkali in these ashes, probably eliminated the tannin from the coca leaves, leaving free the alkaloid.

Cocaine is believed to be the really active principle of the coca leaves. At first, on apparently good grounds, it was believed that this alkaloid possessed a property analogous to caffeine, theine and theobromine, viz.: the checking of waste in the body, but Merck declares it impossible to absolutely confirm this theory. Only the other day I saw it in the same list with guarana, caffeine, theine, etc., but was not considered to be as powerful as caffeine!

Cocaine has in small doses an exhilarating effect on the nerve centres, and it takes but a very small dose to produce toxic effects, especially in cold-blooded animals, but fortunately the effects of the drug do not seem to be cumulative. Schroff, who first experimented with cocaine in 1862, observed that 0.05 grm. given internally to rabbits caused great variation in the pulse and respiration, also temporary mydriasis; the same dose, subcutaneously, caused death in convulsions of an epileptic form, also mydriasis in a strong degree, which disappeared at once after death. In frogs 0.001 grm. produced complete loss of motion, preceded by excitement, while a dose of 0.002 grm. proved fatal.

According to Fronmüller, in 1863, a dose of 0.03–0.33 grm. given internally did not have any important effect upon human beings, simply causing sleep in some, accelerated and then retarded pulse and respiration in others. In a case of attempted suicide 1.5 grams produced no serious injury. Explanatory of this Mr. Merck suggests that the so-called cocaine of that time may not have been really pure cocaine. The preparation with which experiments have been and are being made on this side is Merck's cocaine *urias sol.* and the dose is 0.05 grm.; however, as much as 0.5 grm. of this solution has been administered subcutaneously. The effects of a subcutaneous injection are first a feeling of warmth, then an insensibility to feeling in the neighborhood of the part, and finally a reddening of the skin. In thirty minutes' time these all disappear.

Dr. Aschenbrandt has found in cocaine an admirable restorative from the exhaustion of diarrhoea, as stated in No. 50 of the *Deutsche Medicin. Wochenschr.*, 1883. Prof. Dr. E. V. Fleischel, of Vienna, has proved, to his own satisfaction at least, that cocaine subcutaneously administered is a valuable aid in preventing and curing the morphia habit. (I myself have had excellent results from it with tobacco

chewers who were anxious to break the habit.) In America, W. H. Bently observed the good effects of cocaine in overcoming the taste for morphia. Dr. Freud, of the Vienna General Hospital, reports a case of the morphia habit cured within ten days by the use of 0.1 grm. injected subcutaneously three times daily. His theory is that morphia and cocaine are antagonistic. Dr. Freud likewise credits this alkaloid with aphrodisiac properties.

According to the experiments collected by Merck, the therapeutical value of cocaine may be thus described: Cocaine is a stimulant which quickly increases and serves to sustain, in a harmless manner, the physical powers of the body; it is much more strengthening than alcohol—hence its value on all occasions of extraordinary bodily exertion—in doses of 0.05 grm. at intervals. Whether the powers of the mind can likewise be increased and strengthened is still unknown, as also whether the activity of the nerve centres can be permanently increased. Some good results have been obtained in the treatment of melancholia by the subcutaneous method. It is valuable as a stomachic, particularly after a debauch in eating or drinking. It is credited with the permanent cure of atonic indigestion. The various cachexias, phthisis, exhausting fevers, mercurial poisoning, etc., are said to be averted by its use. Its value in dipsomania is not so well established as in the morphia habit, though there are many satisfactory experiments in that direction. As is well known, cocaine paralyzes the sensation of the mucous membrane when brought in contact with the membrane; hence its possible future in diseases of the throat, and in operations about the mouth and throat. (I have not seen any account of experiments of others, but I am making some observations and trials in regard to the use of the drug subcutaneously in perineorrhaphy and trachelorrhaphy. It will be a happy day for surgeons if these minor operations can be done without ether.)

Of its ophthalmological value enough has already been said.

Cocaine *citras* is used by dentists to anæsthetize the dental nerves. Mr. Merck tells me that this salt can be readily kneaded into pills, which may be wrapped in wadding, moistened and placed in the hollow of the tooth, when lo! the tooth can then be cleaned and filled without pain. If this be true, the man who discovered it is worthy of a high place among the benefactors of the race.

Merck's preparations include the pure cocaine alkaloid, also the salts, viz.: Hydrochlorate, salicylate, hydrobromate, tartrate and citrate. The experiments mentioned above have been made with the hydrochlorate in about 0.05 gramme doses. Von Hoffman, of Baden-Baden, recommends the salicylate for ophthalmological use, but Merck observes that it still remains to be seen which salt is the best for the eye. Finally, cocaine is not a simple substance. When treated with concentrated hydrochloric acid, it splits up into ecgonin, benzoic acid, and methyl alcohol. Mr. Merck informs me that he hopes to be able ere long to report on the physiological action of ecgonin.

That your readers may possibly be as much interested in this subject as I am, is my only apology for thus intruding upon your valuable space.

Very sincerely yours,

SARAH HACKETT STEVENSON, M.D.

BOOK REVIEWS.

COURIER REVIEW CALL BOOK. J. H. Chambers & Co., Chicago, St. Louis and Atlanta.

PHYSICIANS' POCKET DAY BOOK. By C. H. LEONARD, Detroit.

THE 1885 MEDICAL RECORD VISITING LIST. Wm. Wood & Co., N. Y.

THE PHYSICIAN'S VISITING LIST. P. Blakiston, Son & Co., Philadelphia.

The character of these books is too well known to need comment. They are all excellent for the purpose for which they were intended. They are well bound, and in every way convenient.

MISCELLANEOUS.

COMFORTS FOR TRAVELLING INVALIDS.

An article on Winter Health Resorts, in the *Practitioner*, contains some valuable hints to invalids, to relieve the discomforts found by them abroad in travelling and at hotels. One of the discomforts of the long railway journey which is required in order to reach most health resorts, is the constant vibration and the fatigue it occasions. This may be, to a considerable extent, avoided by the proper use of air cushions. It is well for an invalid to be provided with two or more of these. The one to be used for sitting on may be either round or square, and may be either with or without a depression in the centre. Air cushions of a horseshoe shape and furnished at the ends with tapes so that they can be opened out, and again tied together, are most useful. One of them tied round the loins supports the back, and another just like a collar round the neck supports the head, so that in whatever position it is put, sideways, backwards, or forwards, it always rests against the cushion.

Another comfort to invalids travelling, is an India rubber bag for hot water with a flannel case. Indeed it is well to have two of these if the invalid be very delicate. If any sudden chill is felt, or if any pain in the chest or elsewhere should come on, relief is often obtained by the application of a hot-water bottle which takes the place of a poultice. The invalid's friend can easily get these bottles filled with hot water by giving a small gratuity to the guard of the train, and asking him to fill them when stopping at a station. Stoppages occur on the French lines at least every two hours, and the water retains its heat quite long enough to make it useful between the stoppages. Another thing which is a great comfort is an eider-down coverlet, which should be both long and broad.

It is useful, not only during travelling by land or by sea, but during a stay at the health resorts. One would imagine that an eider-down quilt would be bulky and inconvenient to travel with, but this is not the case. In most "hold-alls" there is a large pocket, and the eider-down quilt can be readily rolled up and put into this pocket. When the straps are once around it, it can, from its softness, be easily pressed into a small bulk.

Two other things that the invalid should not be without are a small etna, with can or kettle, to boil water, and a small package of tea, also beef-tea in a portable form, with a small box of biscuits. Even when such things are to be had at a hotel, the prices put upon them are sometimes so exorbitant that the invalid would hardly care to order them, however much he might desire them.

OFFICIAL LIST OF CHANGES IN THE MEDICAL CORPS OF THE U. S. NAVY DURING THE WEEK ENDING DECEMBER 13, 1884.

Craig, T. C., Passed Assistant Surgeon. Detached from Alliance and placed on waiting orders. Dec. 6, 1884.

Curtis, L. W., Assistant Surgeon. To the Naval Hospital, Chelsea. Dec. 9, 1884.

Gaines, J. H., Passed Assistant Surgeon. Present duty continued.

Green, E. H., Passed Assistant Surgeon. Special duty at Museum of Hygiene, Washington. Dec. 11, 1884.

Murray, J. M., Passed Assistant Surgeon. Detached from Naval Hospital, Chelsea, to the Flagship Hartford. Dec. 9, 1884.

Wolverton, T., Surgeon. To the Shenandoah. Dec. 6, 1884.

OFFICIAL LIST OF CHANGES IN THE STATIONS AND DUTIES OF OFFICERS SERVING IN THE MEDICAL DEPARTMENT, UNITED STATES ARMY, FROM DECEMBER 6, 1884, TO DECEMBER 12, 1884.

Hammond, John F., Colonel and Surgeon, retired from active service, by operation of law, on Dec. 7, 1884, under provision of Act of Congress approved June 30, 1882. (S. O. 287, A. G. O., Dec. 8, 1884.)

Irwin, B. J. D., Major and Surgeon, granted one month's leave of absence. (S. O. 112, Dept. Arizona, Nov. 28, 1884.)

McKee, J. C., Major and Surgeon. Leave of absence still further extended one month. (S. O. 288, A. G. O., December 9, 1884.)

O'Reilly, Robt. M., Captain and Assistant Surgeon, assigned to duty as attending Surgeon, Washington City, D. C., to date from October 20, 1884. (S. O. 284, A. G. O., December 4, 1884.)

Porter, Joseph Y., Captain and Assistant Surgeon, sick leave of absence extended four months on surgeon's certificate of disability. (S. O. 286, A. G. O., Dec. 6, 1884.)

Kane, John J., Captain and Assistant Surgeon, from Dept. East to Willets Point, New York. (S. O. 286, A. G. O., December 6, 1884.)

Banister, J. M., Captain and Assistant Surgeon. (Ft. Adams, R. I.) Granted one month's leave of absence on surgeon's certificate of disability. (S. O. 251, Dept. East, December 9, 1884.)

Barrows, C. C., First Lieutenant and Assistant Surgeon, in addition to other duties, to take charge of Medical Division Office, Dept. Arizona, during absence of Surgeon B. J. D. Irwin. (S. O. 112, Dept. Arizona, Nov. 28, 1884.)

Gray, Chas. C., Major (retired.) Died at Geneva, N. Y., Nov. 22, 1884, instead of Nov. 26, as heretofore announced. (Circular Orders A. G. O. December 8, 1884.)