was also normal in appearance. The liver was fairly large, smooth, light in color, and on section showed no macroscopic evidence of necrosis; it appeared slightly fatty, but otherwise perfectly normal. The gall bladder contained a small amount of bile.

The left kidney was large, thick, fairly firm and rounded; there were a number of small subcapsular hemorrhages. The capsule stripped readily, leaving no trace of the hemorrhages on the kidney. Section of the kidney showed a cortex of normal width; there was no congestion. The parenchyma was clouded and granular in appearance. The right kidney showed exactly the same condition and in addition an area of hemorrhage in the parenchyma about a centimeter in length. The pancreas was normal. The right suprarenal was normal. The stomach contained a small amount of dark brown liquid. The mucous membrane was not ulcerated but was merely dotted here and there with a number of small reddish spots. There did not seem to be any break in the continuity of the mucous membrane at these points. The mesenteric lymph glands were normal in size and appearance. The intestines were not opened. The bone marrow from the middle of the shaft of the right tibia was normal in appearance.

Pathologic Diagnosis.—Severe anemia. Jaundice. Numerous small intrapulmonary hemorrhages. Dilatation of the heart. Acute parenchymatous nephritis with subcapsular ecchymoses.

Microscopic Examination of Tissues.—Lungs: Hemorrhagic extravasation into alveoli, congestion and edema. Number of large epithelial cells containing pigment (Hertzfehlerzellen). Kidneys: For most part the epithelium of the tubules presented normal characteristics. In some areas, chiefly in the convoluted tubules the epithelium was granular, the demarcation between cells obscure, and the nuclei invisible. The lumina of a number of these tubules were filled with granular debris; many others were filled with red blood corpuscles, more or less disintegrated. Glomeruli presented no unusual features. Proportion between interstitial and parenchymatous tissue was normal. In small scattered areas were found bodies presenting the characteristics of emboli composed of bacteria, but unattended by any inflammatory reaction. Liver: Some deposit of pigment, probably blood pigment, in the liver cells. (No areas of necrosis.) Suprarenal: Moderate congestion. Pancreas, spleen and bone marrow normal.

HEMIPLEGIA AS A COMPLICATION IN TYPHOID FEVER,

WITH REPORT OF A CASE.*

FRANK SMITHIES, M.D.

Instructor in Medicine in the University of Michigan.

ANN ARBOR MICH.

Hemiplegia, during the course of typhoid fever, is a rare affection. Certain of the acute infectious fevers, e. g., smallpox, scarlet fever, measles, diphtheria, whooping cough, influenza and pneumonia, are not infrequently complicated by more or less complete unilateral paralysis. This affection is sometimes associated with aphasia. It is usually coincident with some disturbance in cardiac mechanism. It may occur at any stage of the special disease under consideration. The greater number of cases have been found to occur in children—perhaps because the diseases mentioned are most common in the first decade of life. But considering the severity of many of the typhoid infections, and the widespread prevalence of the ailment, is it surprising to note the infrequency of paralysis of any form as a complication, and especially of hemiplegia.

Numerous cases are on record in which paraplegia has complicated the course of the disease. Paralysis of single muscles, as, for example, the external rectus of the eye, or of certain muscle groups, such as the

muscles of deglutition or the muscles of the posterior part of the thigh, is rather more common. However, one seldom meets with more than a passing mention of unilateral paralysis. Isolated cases are scattered throughout the voluminous literature of typhoid fever, but systematic tabulation is lacking. In view of the fact that the complication may occur in the course of any given case, and that, when present, it brings up an important point in the prognosis of such a case, it has been thought that a brief review of the subject might be of interest and value.

HISTORICAL.

Gubler, in 1860, first called attention to hemiplegia occurring at the "height of fever" in a young girl, a syphilitic, seven months pregnant. Aphasia was present. A few years later Scoresby Jackson published a careful analysis of a case in his own practice, in which hemiplegia and aphasia had suddenly appeared in the course of typhoid fever. Jackson paid more attention to the speech disturbance than to the paralysis, but nevertheless gives a most fascinating description of the progress of both lesions. Soon after the appearance of Jackson's paper Benedict reported two cases, in both of which there were disturbances in speech. The same year Karmin described hemiplegia without aphasia in the case of a young adult. The paralysis came on as late as the eighth month of a particularly severe attack of typhoid with prolonged convalescence. Studies by Berger and Nothnagel soon followed, the latter considering the subject far more completely than it had been heretofore. Between 1872 and 1889 only four cases appear in the literature that was accessible. Caley mentions hemiplegia during convalescence in a lad of 11, without aphasia; Gee cites its occurrence in a boy of about 7 years, with aphasia, and five years later Sorel reports hemiplegia and aphasia at the nineteenth day in a young adult male. A writer in the Middlesex Hospital Reports of the same year recorded hemiplegia coming on the ninth day, with convulsions and unconsciousness, in a man of 24. In this case also there was aphasia. It was not, however, until Sir Francis Hawkins' address before the London Clinical Society in 1889 that hemiplegia, in the course of typhoid fever, became recognized as a complication which was worthy of serious consideration. Thus far it had been looked on as a curiosity in medicine. Hawkins reported seventeen cases which he had collected from the literature or personally observed. He gave a masterly discussion of the conditions noted and added much of value to the pathology. No further summary of the lesion occurs until that of Osler's in 1899, although several cases had appeared in the literature, and a surprising number of them had been observed at the Johns Hopkins Hospital. Osler added four cases to the growing list, and was fortunate enough to be able to tabulate the postmortem findings in two of the cases reported. Since 1900 about a dozen cases have been mentioned, some with very complete detail, others with most meager observations.

FREQUENCY.

Holscher, summarizing 2,000 fatal cases of typhoid fever at the Pathological Institute in Munich, mentions twenty cases as showing "cerebral apoplexy," but gives no details concerning the locations of the lesions nor the antemortem manifestations. Perusal of the Charité Annalen of Berlin from 1874 to 1903 reveals the surprising fact that of 4,793 cases of typhoid fever, of which 875 were fatal, no cases of hemiplegia occurred.

*From the Clinic of Internal Medicine of Professor Dock.

In an analysis of 1,000 cases of typhoid fever in America Herriek cites no instance. McCrea's tabulation of 707 cases at the Montreal General Hospital includes one case of paralysis of one side of the face and one case of paralysis of the muscles of deglutition, but no mention of hemiplegia. A somewhat careful review of the Guy's Hospital Reports, London, fails to find mention of hemiplegia complicating typhoid fever.

With the case herewith reported, the number of cases of which some details can be given has been increased to more than forty. It is probable that more extensive search may reveal a greater number, but further literature than that recorded was not available. For purposes of reference and convenience of comparison, a tabulation of all the cases reviewed is given, but before considering the reports in detail it is deemed advisable to present a summary of the patient which the writer recently had opportunity of watching.

CASE.

Patient.—A male, aged 36, a farmer, was admitted to Prof. Dock's clinic Sep. 30, 1906, on account of fever of about six days' duration, cough, headache, general weakness and discomfort. He came from a village in which there existed a small epidemic of typhoid fever.

History.—Family history was negative. Patient was puny as a child; had the usual diseases of childhood; made a rather slow recovery from measles, suffered with chronic bronchitis for some time afterwards; contracted syphilis at age of 24; had ulcer on penis, sore mouth, lost some hair, and had a rash. He took no treatment at that time. Patient said that the disease left him with a "feeling of weakness below the knees," that the knees "played out suddenly" and that for some time he had difficulty in walking any great distance. He gradually recovered, but six years later again suffered from "weak feelings" in his legs, but was not disabled. In April 1906, he had a third, similar attack, and had to give up his farm work. He then began specific treatment, and managed to get about again, but did not feel strong enough to work. He gave no history of alcoholism.

Present Illness.—On Sept. 24, 1906, he began to suffer from headache, "flashes of heat and cold," cough, with a little sputum, severe pains in back and top of head, flatulence; thinks that he had fever; called a physician, who sent him to the hospital.

Examination.—On admission at 10 a. m. his temperature was 101.5 F.; pulse 110; respiration 36. He was able to walk with slightly spastic gait; he was rational; face flushed; had anxious expression, lips tremulous and slightly cyanotic; tongue very red and moist; extremities cold. His abdomen was moderately distended, the abdominal wall lax, with slight tenderness over both lower quadrants; the spleen extended to the rib margin in left nipple line at the end of deep inspiration. Heart showed slight enlargement to right; all sounds weak; pulse was full, rapid, regular, compressible with no diastolic. Lungs disclosed harsh breathing over both upper anteriors, with few coarse, mucous râles.

Blood: Red blood cells, 5,560,000; white blood cells, 5,348; hemoglobin, 95 per cent.

Urine: Positive diazo and indican tests.

Progress of Case.—On the second day in hospital, temperature ranged from 100.4 F. to 102. F.; pulse 88 to 120; respiration 28 to 32. Patient became delirious and rather violent towards evening. Splenic dullness disappeared. On the following morning, was rational, but stupid. Temperature ranged this day from 99.8 F. to 103.4 F.; pulse 80 to 112; respiration 28 to 36.

Oct. 3.—Widal positive, moderate clumping in 45 minutes with dilution 1:40 (Dr. McMichael). Towards evening the patient became delirious and very restless, and stood his tub-baths, at 65 F., rather poorly.

Oct. 4.—In the afternoon, patient complained of "cramping" of the last two fingers of his left hand, and of pain in his back, head and ankles. He became extremely restless, eyes be-

came staring, arms and legs rigid, had difficulty in swallowing, and lay with his head drawn sharply to the right. Temperature ranged, 99.2 F. to 101.8 F.; pulse, 96 to 122; respiration 28 to 42.

Oct. 5.—Slight tremor over entire body, extremities cold, skin mottled and dusky. Heart sounds very weak; reacted poorly to bath at 65 F. Blood pressure: Systolic, 112 mm. Hg; diastolic, 65 mm. Hg, with the Stanton 12 cm. cuff. During the night the patient's head continued to be strongly drawn to the right, there was marked tremor of the jaw and the upper part of trunk; the arms were rigidly flexed at the elbows. Cheyne-Stokes respiration appeared; the patient urinated involuntarily. On account of the tremor the blood pressure variations could not be followed.

Oct. 6.—Condition much the same; tremor slightly less marked. Patient became conscious during the early part of the day. He was unable to protrude his tongue, and his head was noticeably retracted. The heart sounds were weak at the apex, and a soft, blowing systolic bruit could be made out. Later in the day his face became very deeply flushed, violent coarse tremors and twitchings appeared over both upper extremities, especially over the right, the patient became delirious, and Cheyne-Stokes respiration again appeared. Temperature 99.2 F. to 103 F.; pulse 98 to 120; respiration 28 to 44.

Oct. 7.—Tremors more noticeable, especially over the right arm and the right face; ptosis of the left upper eyelid, slight photophobia; tongue could not be protruded; involuntary urination; abdomen slightly distended; heart sounds weak, systolic bruit at apex still present. Temperature 98.2 F. to 100.4 F.; pulse, 88 to 116; respiration, 28 to 40. Blood pressure: Systolic, 112 mm.; diastolic, 75 mm., with the Erlanger 12 cm. cuff.

Oct. 8.—(Nine days after the patient entered the hospital, and about fifteen days after the beginning of his illness.) Patient lay with head drawn to the left; left angle of mouth drawn down; left arm flexed rigidly at right angles across thorax; very coarse tremor over both upper extremities, especially the right. Legs could not be freely extended on the thigh, when thigh was flexed on trunk; tongue deviated to the left; involuntary urination. Patient in stupor greater part of day. Blood pressure: Systolic, 102 mm.

Oct. 12.—Patient rational, but dull; pulse very soft and irregular; moderate dilatation of the left heart. Blood pressure: Systolic, 99 mm.; diastolic, 60 mm., with the Erlanger 12 cm. cuff.

Oct. 11.—Distinct ptosis of the left upper eyelid; moderate deviation of left eyeball to left; pronounced lowering of the left angle of the mouth; marked leftward deviation of the tongue; difficulty in swallowing; flattening of the left nasolabial fold. There was slight rigidity of the left forearm, but a fair degree of motion on the right. The patient was unable to open his left hand, and only partly could extend his left forearm, slowly dragging it along the bedclothes. Patient could not elevate the left arm from the bed; there was no grip in the left hand; there was moderate movement on the right with fair grip. The legs were slightly flexed, the right moved slowly, the left could not be moved. There was marked sensitiveness to stimulation of the skin, with marked spasm of the leg and thigh muscles on such stimulation. Sensation otherwise was negative. There was slight Babinski's sign on the left; ankle clonus on both sides, especially on the left; patellar reflex was reduced on both sides. The patient complained of severe pains in the muscles of both legs. Temperature 96.6 F. to 102.2 F.; pulse 80 to 96; respiration 20.

Oct. 12.—*Laryngoscopic examination.* (Prof. Canfield) Right side negative; partial paralysis of the left vocal cord; throat negative. Deep reflexes (Dr. Klingmann) showed no change in character, but a general diminution in response on the left, with increase on the right; no reaction of degeneration.

Oct. 13.—Patient felt much better; could move right arm and leg freely; there was little motion on the left side, but the rigidity had almost disappeared. Eyes, mouth and tongue as before. There was still difficulty in swallowing; patient urinated involuntarily; slight Babinski's sign still present on the left; pronounced left wrist-drop, with flexion of the fingers of the left hand. From this time on the patient passed through

a rather eventful convalescence, the main events of which are summarized in the following notes:

Convalescence.—Oct. 16.—Intense muscular tenderness in both legs and arm, especially the left; moderate but painful swelling of the glands in the left axilla. Oct. 20: Marked cardiac irregularity, with pronounced extra systole and loud systolic murmur at apex and in axilla. (Patient had been taking digitalis for dilated heart.) Babinski's sign disappeared on the left. Oct. 25: Moderately severe phlebitis in the left femoral. Nov. 1: Patient complained of severe general headache. Put on potassium iodid. Nov. 6: Right arm and leg practically normal; patient sat up in bed for a few minutes. Nov. 16: Moderately severe dacryocystitis on the left. Nov. 23: External strabismus of the left eye had disappeared; no deviation of tongue; patient could control his urine. Dec. 12: Patient walked with assistance: dragged left foot at toe, and swung left leg from trunk when walking; could not raise left leg over right knee. Dec. 16: Moderate edema of left leg and left hand. Patient had been moving about rather freely. Jan. 5, 1907: Patient was feeling fairly well, and was gaining in both weight and strength. Jan. 7: (Prof. Dock) Patient looked thin; lips and ears were red; vesiculo-papular eruption on forehead (potassium iodid?) walked fairly well; left arm very weak; tongue pushed slightly to the left. Jan. 25: The patient was about to be discharged; looked fairly well; was within about ten pounds of his normal weight, and could walk about twenty rods without feeling fatigued. Had difficulty in climbing stairs. Still slight ptosis in the left upper eyelid; pupils negative; could pucker lips so as to whistle, but could not whistle a tune; had slight tremor of both lips; the tongue was fuller on the right and deviated slightly to the left. The right hand and arm showed no disability. The left arm hung loosely at the patient's side; the left hand was puffy and cooler than the right; the arm was slightly flexed at the elbow and the fingers were flexed into the palm. The patient could approximate his fingers to his lips, and could raise his arm laterally about one-third of the distance to the shoulder-level and about half way, anteriorly. He could extend to about an angle of 160 degrees at the elbow. Flexion, without assistance, was very limited. He could extend fingers to about an angle of 170 on the hand, but distal phalanges were only capable of slight extension; flexion of fingers was fairly complete, with moderate grip. He could hold a full cup in his fingers. In walking, the patient swung his left leg stiffly from the trunk and shuffled his left foot slightly. He could move his toes, but there was a moderate degree of stiffness at the left ankle joint. Patient could cross his left leg over his right knee with little difficulty. There were no disturbances of sensation. He had no difficulty in swallowing. The jaw reflexes were increased, as were also the elbow and patellar deep reflexes; no ankle clonus; no reaction of degeneration.

Subsequent History.—The patient returned to the hospital for re-examination about the middle of February, 1907. He had gained considerably in weight and had steadily improved in strength. Movement of the left side was in every way freer, more definite and firmer. The patient had been on potassium iodid since leaving the hospital, with massage and faradism. Headaches were rare; mentally he seemed much brighter. He was advised to continue his medication, to get massage and passive motion when possible.

LESION.

The somewhat gradual onset of the hemiplegia, with the succeeding progressive recovery from the condition, points strongly to a thrombotic process in one of the cerebral arteries or its branches or both. The artery involved was doubtless the right middle cerebral, with its off-shoots to the various motor centers. The early disability of the right side may be explained by temporary disturbance in the circulation in the left middle cerebral or its branches. However, it is not impossible that extension of the thrombosis on the right side could have resulted in material interference with the flow of blood in the circle of Willis, and, as a consequence, the

free passage of blood to the left motor area was retarded. The absence of any definite cardiac lesion, and the relatively slow onset of the paralysis, together with the extent of that established, are reasonably strong evidence against the process being primarily embolic. The large extent of the motor cortex simultaneously involved, and gradually rendered functionless, argue against hemorrhage as being the chief cause of the trouble. However, it is impossible to say that there was not slight hemorrhage, with early thrombosis following. The history of syphilis in this patient gives a significant hint as to the underlying pathologic alterations in the cerebral arteries, which rendered interference with the circulation possible—in fact, it is a question as to just how much the patient would have suffered had he not been luetic.

THE TABLE IN SUMMARY.

In the table herewith presented I am able to furnish information concerning forty cases of hemiplegia in typhoid fever. There are several cases mentioned to which I have not had access, and so am unable to properly complete the list. In addition to the cases herewith tabulated, I have to mention cases reported by Courtade,¹ Cohin² and Moore.³ I regret that I have not been able to consult the literature for these cases. The interesting features of the summary will be considered in detail.

Sex.—Of the thirty-three cases in which the sex is mentioned, twenty-three are males, nine are females; in two instances the sex is not recorded.

Age.—Four cases were 5 years or under, five cases were between 5 and 10 years, eight cases between 10 and 20, thirteen cases between 20 and 30, and two cases were more than 30 years of age. In the remaining cases the ages were not stated. It is readily seen that the lesion is more common in young adults, and this is explained, doubtless, by the greater prevalence of typhoid fever at that period of life.

Associated Conditions.—Syphilis is mentioned in two cases, including the one reported in this paper. Pregnancy complicated in one; in another there was purpura. Several cases were accompanied by hemorrhage from the bowels. Cardiac lesions were not frequent. It seems that in few instances were any lesions other than those of the febrile state present. In the case herewith presented at times there were present a softly blowing systolic bruit at the apex, which became more marked under digitalis medication. There was considerable irregularity in rhythm, with marked diastolic incompetence.

Aphasia.—In thirty-two cases in which disturbances in speech were recorded aphasia occurred in twenty-six; it was absent in four cases, and in the remainder of the cases there was unconsciousness, with later faltering speech or death before the presence or absence of aphasia could be determined. In the case given here the patient did not answer well for several days. He appeared to understand questions, but responses were very tardy.

Time of Onset.—This varied widely. It is not reported in all of the cases. In thirty cases in which some details are given, in but one case did the hemiplegia occur during the first week of the disease. Eight instances are recorded as occurring during the second week, eight during the third week, two during the

1. *Encephale*, Paris, 1886, 431.

2. *Bull. general. de Therap.*, Paris, 1887, 397.

3. *Trans. Royal Acad. Med.*, Dublin, 1900, xvii, 492.

TABULATION OF CASES OF HEMIPLEGIA COMPLICATING TYPHOID FEVER.

No.	Reported by	Sex and Age.	Occurrence.	Symptoms of Onset.	Aphasia.	Side.	Result.
1	Gubler: Arch. gen. de Med. 1890.	F., 19.	"Height of fever."	Syphilis; pregnant; seventh month.	Present.....	R.	Recovery.
2	Jackson: Edinburgh Med. Jour., 1857.	M., 21.	Eighteenth day....		Present.....	R.	Recovery almost complete.
3	Benedict: Electrotherapie, 1863.	?	?	?	Present.....	R.	?
4	Karmin: Wiener Med. Press, 1868.	M., 24.	Between eighth and ninth month of prolonged and severe case.		No cerebral disturbance.	L.	Persisted.
5	Benedict: Electrotherapie, 1863.	?	?	Atrophy of optic nerve.....	Speech faltering.....	L.	?
6	Berger: Berlin klin. Wochschr., 1870.	M., 21.	Third week.....	Unconscious.....	?	L.	Persisted.
7	Nothnagel: Dent. Arch. f. klin. Med., 1872.	M.	During febrile state.	Convulsions.....		R.	?
8	Eulenberg: Quoted by Nothnagel.	?	?	?	Present.....	R.	Recovery.
9	Same.....	?	?	?	Present.....	R.	Recovery.
10	Calley: Med Times and Gaz., 1878.	M., 11.	During convalescence.		Not affect speech.....	R.	Recovery.
11	Gee: Med. Times and Gaz., 1878.	M., 7-8.	During convalescence.		Present.....	R.	Recovery.
12	Sorel: L'Union Med., 1883.	M., 24.	Nineteenth day....		Present.....	R.	Persistent one year after.
13	Hawkins: Clin. Soc. Trans., 1889.	F., 18.	Twenty-ninth day.	Hemorrhage from bowels; purpura.	Present.....	R.	Died. Clot in right middle cerebral artery and in left auricle.
14	Middlesex Hosp. Rep., 1893.	M., 24.	Ninth day.....	Convulsions; unconscious.	Present.....	L.	Recovery.
15	Newbolt: Lancet, Aug. 27, 1892.	M., 21.	Forty-first day....	Hemorrhage from bowels.	Present.....	L.	Recovery.
16	Strümpell: Lehrbuch.....	F., 10.	"Fourth fever-free day."	Convulsions; severe attack.	Present.....	R.	Recovery in fifteen weeks.
17	Same.....	M., 15.	Third week.....		Present.....	R.	Recovery in ten weeks.
18	Kummell: Ztschr. f. klin. Med., II, 2.	M., 30.		Severe attack; much meteorism.		?	?
19	Curshman: Nothnagel's Syst. Med.	F., 19.	Third week.....	Moderately severe attack.	Present.....	R.	?
20	Haynes: Johns Hopkins Hosp. Bul., 1893.	M.	Twelfth day.....	Delirium.....		R.	Recovery partial.
21	Blumer: (Same).....	M., child.	Seventeenth week after typhoid.	Violent, sudden, convulsions.	Present.....	R.	Recovery partial.
22	Thayer: (Same).....	M., 21.	Tenth day.....	Stupor.....	Present.....	R.	Recovery partial.
23	Thayer: (Same).....	F., 10.	Nineteenth day....	Typhoid state; involuntary passage of feces and urine.	Seemed to understand but did not answer.	R.	Much improved.
24	Rollleston: Brit. Med. Jour., 1893.	M., 30.	Twenty-fourth day.	No convulsions.....	Present.....	L.	Gradual recovery.
25	Herringham: (Same).....	F., 9.	Third week.....			R.	?
26	Emerson: Johns Hopkins Hosp. Rec., 1899-1900.	M., 45.	Eighth day.....	Headache; diplopia.....		L.	Gradual improvement.
27	Osler: Johns Hopkins Hosp. Rep., 1899-1900.	M., 22.	Fifth day.....	Sudden, uneasy feeling in head; general, clonic convulsions, beginning in both arms simultaneously.	Unconscious.....	L.	Died. Hemorrhagic enteritis of ileum; ulcers in Peyer's patches; thrombosis in ascending parietotemporal artery, including points of origin in mid-cerebral, also recent thrombosis in branches running into Rolandic fissure.
28	Osler: (Same).....	F., 7.	Tenth week.....	Fever; sudden convulsions.	Present.....	R.	Partial recovery; both lesions.
29	Osler: (Same).....	M., 25.	End second week.	Delirium; no convulsions.	Not present.....	L.	Partial recovery.
30	Osler: (Same).....	M., 43.	Third week.....	Cheyne-Stokes respiration; cardiac arrhythmia; nausea.	Not present.....	L.	Died. Ordinary intestinal lesions of typhoid fever, softening of upper part of right internal capsule and thrombosis of artery.
31	Ellis: N. Y. Med. Rec., Oct. 1, 1892.	Young lad.		Symptoms pointing to poliomyelitis.			Complete recovery.
32	Diller: N. Y. Med. Rec., 1903.	M., 5.	Three and one-half weeks.	Moderate fever; much tympany.	Present.....	R.	Recovery partial.
33	Dankin: Lancet, Apr. 23, 1892.	F., 30.	Thirtieth day....	Pains in head.....	Present.....	R.	Died. Clot in middle cerebral artery; heart dilated.
34	Hrach: Wien. Med. Wochschr., No. 42, 1902.	M., 23.	Fourteenth day....	Moderately severe attack: convulsions; facial paralysis first three days then hemiplegia.	Present.....	R.	Improvement of both lesions; contractures of right hand.
35	Barrett: Mich. State Asylums for Insane.	F., 5.	During second week.	Moderately severe attack.	Not present.....	R.	Epileptic attacks followed; contractures, with marked shortening of right arm and leg; athetoid movements; patient died at age of 17 of pulmonary tuberculosis. Postmortem examination of brain showed early thrombosis of second branch of Sylvian artery, atrophy of entire motor tract and replacement of parenchyma with dense scar tissue.
36	McCaskey: Studies.....	Child, 5.		Severe attack.....	Present.....	?	Fatal.
37	Stevens: Glasgow Med. Jour., 1897.			Severe attack; onset gradual.			
38	Dantsig: Ejened. Jour. Prakt. Med., St. Petersburg, 1898.				Present.....	R.	?
39	Stein: Pest. Med. Chir. Pres., Budapest, 1904.				?	?	?
40	Baylac: Toulouse Médicale 1904.			Convulsions.....	Jacksonian type.....	R.	
41	Teissier: Progrès Médical, 1835, -440.				Present.....		
42	Sakorrhaphos: 'Ιατρικὸς μηνύτωρ Ἀθῆναι, 1903, iii, 75.				?	R.	
43	Smithies.....	M., 36.	Seventh day.....	Convulsions, delirium; stupor; Cheyne-Stokes respiration; moderate temperature.	Did not answer well for two days; seemed to understand.	L.	Gradual improvement contractures hand and leg; muscular atrophy.

fourth, and the remainder during convalescence; in one case as late as the eighth month of a particularly severe attack with prolonged recovery period.

Mode of Onset.—In ten cases the hemiplegia was preceded by convulsions. In four instances these came on suddenly and violently. With the convulsions there was usually associated unconsciousness, either temporary or long continued. In four instances there was delirium—in the case here given it was quite violent at first. In three cases there was stupor. Several instances of very severe headache preceding the onset of paralysis are recorded. Hemorrhage from the bowel was noted in one instance, and in another there was choked disc. In the majority of the cases the height to which the fever rose is not mentioned. In a few instances there was very high temperature, with low muttering delirium and subsultus tendinum. In the case given there was but moderate rise in temperature.

The Side Affected.—This is definitely mentioned in thirty-five of the reported cases here tabulated. In twenty-one instances the paralysis was on the right side; in ten cases it was on the left side. In the remaining cases the side is not definitely mentioned. In only two cases of right-sided paralysis was aphasia absent. In three of the left-sided paralyzes aphasia was present.

Postmortem Findings.—In Dankin's case there was a clot in the left middle cerebral artery. The heart was dilated and there was a hemorrhagic infarct in the lower lobe of the left lung. In Hawkins' case there was a clot in the left middle cerebral artery and thrombosis of the right auricle. Two of Osler's cases were fatal. One exhibited hemorrhagic enteritis and ulcerated Peyer's patches as the primary typhoid lesion. Section of the brain revealed thrombosis of the ascending parietal branch and the parieto-temporal branch of the middle cerebral artery, including their points of origin. There was also recent thrombosis of the branches running into the Rolandic fissure, the sulcus between the frontal gyri and the ascending frontal convolutions and the branches supplying the temporo-parietal region generally. The other case reported by Osler showed ordinary typhoid lesions in the bowel. In addition there was softening in the upper part of the right internal capsule. The area was about the size of a hickory-nut and lay lateral from the caudate nucleus and a little median and slightly dorsal from the upper border of the cortex of the island of Reil. Barrett's case died twelve years after the onset of the hemiplegia, from pulmonary tuberculosis. The autopsy showed evidence of early thrombosis of the second branch of the left Sylvian artery, with atrophy of the entire motor area. The cerebral parenchyma was replaced by scar tissue. In the remaining fatal cases the postmortem findings were not recorded.

Results.—In six of the cases the patients died. Twelve recovered completely. The shortest time to complete recovery was about twelve weeks. In the great majority of the remaining cases there was usually gradual improvement of the paralysis. This was especially so as far as the gross movements of the parts involved were concerned. The finer movements requiring delicate co-ordination were, in most instances, late in returning, or were lost completely. In three instances the hemiplegia persisted with no improvement. Recovery from the paralysis was generally noted as being more rapid and more complete in the lower limbs, with the muscles of the feet and legs regaining their functions earlier than

those in the arms and the hands. Muscular weakness, more marked than that usually noted in post typhoid states, was frequently commented on. In many cases there was moderate atrophy of the parts involved. This usually was more apparent in the muscles of the hands, arms, shoulder girdle and legs. Facial hemiatrophy was occasionally observed. Contractures, both early and late, were commonly present, even with the most careful therapeutic measures adopted to prevent such. In Barrett's case there was shortening amounting to several inches of the right arm and the right leg, associated with athetoid movements. The contractures led to complete disablement. This case was also complicated with epileptiform attacks. In the patient, whose progress furnishes the basis of this paper, after six months' time the man was still greatly incapacitated. There was slight shortening of the left leg, with spastic paralysis of the posterior leg muscles, and partly of the posterior thigh muscles, contractures of the flexors of the left hand and paralysis of the anterior forearm muscles and the left trapezius. Co-ordination in the movements of both arm and leg was imperfect. Slight ptosis persisted in the left upper eyelid, the left face muscles were sluggish and incomplete in their action.

In the patients in whom the hemiplegia was associated with aphasia the recovery from the latter disturbance was much slower and less complete than from the former lesion. In some cases there was but slight improvement in the aphasia. In others only after persistent teaching efforts and various exercises did the patients succeed in speaking. In but few of the patients was there complete recovery. It might be said, indeed, that in no instance was the speech so perfect as before the illness.

In the non-fatal cases loss of bladder control has been but temporary. Involvement of the muscles of deglutition has not been permanent. Sensation has usually been but little interfered with. In the case here reported there was not the slightest sensory disturbance at any time, except the rather painful neuritis of the early period. Athetoid movements on attempted voluntary movement have been reported by both Osler and Barrett.

The mentality seems to suffer no serious alteration. The delirium of the early stages of the paralysis has not been observed to have very lasting effects. In a few patients, even without aphasia complicating, slow cerebration has persisted for some time after the febrile state. Confusion and hesitancy of speech is fairly common, persisting for some months, as in the case of the patient recently under our care. There seems to be, in some instances, more than the ordinary tendency toward post typhoid neuroses, particularly of the psychic nature. But this is not to be wondered at in view of the serious nature of the disability. For a man who has been previously well and active to suddenly find himself a helpless cripple, with but a prospect of partial alleviation at the best, is certainly sufficient ground for pessimism, if not actual melancholia.

Causes of the Hemiplegia.—In two instances syphilis doubtless predisposed. In the fatal cases coming to autopsy thrombosis has been the rule. This has occurred, irrespective of whether or not the lesion has appeared, as Hawkins suggests, early or late in the course of the typhoid fever. Whether or not the damage was done as a result of an early cardiac involvement can not be stated, inasmuch as observations on the heart condition are lacking in the majority of the reported cases.

In hemiplegia complicating other acute febrile diseases, cases are on record in which the cardiac valve-flaps have been found to be normal, and yet clots have been found in the heart and thrombi in the arteries. Hawkins cites Henoeh as mentioning an instance in which embolus was found in the right Sylvian artery and a thrombus on the left side of the heart. In many cases dirotic pulse and cardiac irregularity were marked, especially in those reported by Hawkins, Osler and that here given. When the lesion comes on early in the course of the fever, with sudden, severe convulsions and loss of consciousness, the evidence appears to be that the causative factor is either an embolus or a small hemorrhage involving the motor areas of the cerebrum. When the condition is a complication of the later stages of the disease the damage is doubtless due to thrombosis, although one can not eliminate the possibility of a hemorrhage occurring as a result of sudden strain on the arterial walls. The recent studies of Thayer on the blood-vascular changes in typhoid fever might lend strength to the opinion that there are some cases in which thromboses may develop without much coincident cardiac weakness. The possibility of the injurious effects on the circulation from a diffuse encephalitis must not be overlooked.

Prognosis.—Death occurred in about 15 per cent. of the cases in which hemiplegia complicated. In these cases the lesions in the bowel did not appear to have been the most serious factor in causing the fatal termination. In the non-fatal cases the prognosis is for gradual partial recovery. When the lesion is associated with aphasia the recovery is apt to be less rapid and less complete. The aphasia is apt to persist. However, this rule does not apply in all cases. In about 8 per cent. of the cases there was no improvement.

Treatment.—This naturally falls into (a) treatment during the onset of the paralysis, (b) treatment of the condition after hemiplegia has been established, and (c) treatment of the primary infection.

DURING ONSET OF PARALYSIS.

Measures whereby cerebral congestion is relieved are indicated. Ice-bags may be applied continuously to the head. They are of great service in controlling the delirium, and certainly, in my experience, exert a favorable influence on the convulsions when present. Doubtless they are of value in preventing the diffuse cerebritis, as suggested by Sir Thomas Barlow. The Brand bath should be discontinued during the convulsive stage, but in the absence of convulsions with simple delirium the cold tubs, with constant pouring of cool water on the head, do much to relieve the mental symptoms. Frequently by this treatment the onset of severe convulsions may be prevented. In those cases in which cerebral congestion is extreme, as shown by the swollen vessels, the purple face, the wandering mind, the high temperature and the dilated heart, venesection—if quickly done—is of undoubted service. In the cases with less evidence of cerebral congestion, and with slow onset of the paralysis, blood-letting is contraindicated, as predisposing, by retarding the circulation, to increased thrombosis. Local blood-letting, as by the application of leeches over the mastoid area, as suggested by Berger, may be tried and may be of some value in selected cases.

The action of brisk purges, as the use of salines or a few drops of croton oil, may be of advantage in the cases in which the paralysis comes on early, but would, of course, be out of place in the later stages of the disease when ulceration of the bowel has taken place.

Warm bags to the feet or mustard cloths to the legs may be serviceable in some instances. For the cardiac weakness digitalis and strychnin may be necessary, although on account of their action on the blood pressure they should be used with caution. Aconite is of value in cases with high pressure and rapid heart. In many instances the constant application of an ice-bag over the precordia will do much to prevent sudden dilatation and to control such when once present. The convulsions seldom require chloroform. It should be given cautiously. Bromids in large dosage, in the early stages, may be useful for controlling spasm and convulsions, and are quite harmless if cautiously given. For the intense headache nothing is so grateful to the patient, or so harmless in its action, as the ice-bag or frequent sponging with cold water. An occasional dose of acetphenetidin is sometimes of value, but on account of its depressing effect its use should not be persisted in.

Surgical interference may be of value to relieve pressure or remove clots or thrombi in cases where an operator of the highest skill is available.

AFTER HEMIPLEGIA IS ESTABLISHED.

When the hemiplegia has been established supportive measures are indicated. In the early stage the patient may have to be frequently catheterized on account of the loss of sphincteric action. He may have to be fed through a tube or by the rectum, the latter procedure being inadvisable except in the very early cases or those coming on during late convalescence. It is to be remembered that an ordinary patient will go several days without much nourishment, and with little harm to himself, provided sufficient liquid is furnished in the form of water. Burns from hot water bags and the resulting bed sores should be carefully avoided or, if present, should be kept scrupulously clean, protected and dusted. In those cases which come on late, or after the acute infection has subsided, the indications are to prevent atrophy and contracture. Massage with warm olive oil several times daily is of much value, but must not be too vigorous, inasmuch as it occasionally is largely instrumental in the production of muscle tenderness and phlebitis. It should always be from the periphery. Passive movements of all the involved joints should be performed, gently at first and not over too long a period. In this way contractures and ankylosis are largely prevented. Faradism is often of value in promoting nerve and muscle nutrition.

Careful feeding, outdoor life and frequent bathing will do much to restore function. In those cases in which syphilis is present or suspected, full doses of the iodids are indicated. For the obstinate edemas, which frequently occur in the dependent parts, e. g., as in the hand, rest, with elevation, is indicated. If such a condition persists elastic web bandages, not too strong, may be worn when the patient is in the upright position. The patient should be encouraged to make voluntary movements as often as possible, and should be told the importance of attempting the finer movements of co-ordination. Exercises calling for such movements are readily invented, and frequently the patient performs them with enthusiasm. When aphasia is present exercises for phonation, for recognition and description by writing or by speech should be arranged.

TREATMENT OF THE PRIMARY INFECTION.

The carrying out of the measures for the relief of the symptoms associated with onset of hemiplegia, and for its treatment after once established, do not inter-

fere with the routine treatment of the acute infection. The cold tubs may have to be suspended for a time; the patient will require more care during convalescence, which may be long drawn out, and the prolonged stay in bed may render him more liable to complications, but by taking care of such as they arise one can usually control the situation.

I wish to acknowledge my sincere thanks to Prof. Dock, Prof. Canfield, Dr. Klingmann and Dr. Freund for aid in the examination of the patient herewith presented and for suggestions in the collecting of data.*

THE TREATMENT OF PSEUDARTHROSIS AND RETARDED CALLUS FORMATION BY INJECTION OF BLOOD.*

VICTOR SCHMIEDEN, M.D.

Privatdocent of Surgery; Chief of the Surgical Clinic in the
University of Bonn.
BONN, PRUSSIA.

It is now more than two years since Professor Bier reported his first experience in treating pseudarthrosis and retarded callus formation in a new and interesting way. It comprises the injection of venous human blood between the fragments, which blood is taken from the arm of the patient or from that of another healthy man, in order that callus formation may be stimulated.

As the manner in which this injection excites the formation of callus is not perfectly clear, I will make a few explanatory remarks regarding it. In every case in which there is a fresh extravasation of blood one sees the signs and symptoms of a slight inflammation. The affected part is painful, edematous, and many times the patient has fever and a feeling of heat in the part, at times also the patient has albuminuria.

These phenomena, which are but the manifestations of the reaction of the body to the extravasation of blood, seldom fail, and are especially marked when, instead of the patient's own blood, the blood of another person is injected. The most intense reaction is called forth by injection of the blood of animals.

These immediate symptoms of absorption are followed by others, which are active for a long time. Hard, new connective tissue is formed in which blood pigment may be deposited, and often more highly organized tissue is destroyed by these scars which follow such extravasations, for instance, in joints, muscles and in the central nervous system.

The inflammatory irritation produced by such extravasations can be easily demonstrated in simple fractures. There is intense pain, marked swelling and increased amount of heat in the part. Bier believes that in these cases the inflammatory reaction is due to the extravasation of blood, and that the formation of callus is caused by this inflammatory reaction.

The fact that cases which often heal with the formation of a large amount of callus are those cases in which a large extravasation has occurred seems to point to the above-mentioned logical conclusion. On the other hand, we often see delayed callus formation in compound fractures on account of the escape of the blood. Further, in hemorrhages near bones we see at times a thickening of the periosteum; also in cases of repeated traumatic hem-

orrhages we see the formation of osseous nodules in the muscle substance. In the same manner the intentional irritation of the fractured ends of bones and the pounding on the site of the fracture, as practiced by Thomas, excited callus formation, not by the direct irritation, but by irritation excited by the hemorrhage which this treatment caused. These various considerations prompted Bier in his efforts to induce callus formation by injection of blood, and these injections have proven the truthfulness of his suppositions.

One can divide the pseudarthrosis into two classes, the one due to inefficient treatment (marked dislocation of the fragments), the other due to a general condition of the patient which is not favorable to the formation of callus. In the latter class there are many conditions which I will not enumerate. All surgeons have seen cases in which the fractured ends have been perfectly reduced and the best treatment followed, and yet no callus formation takes place. In such cases all treatment is in vain, even suture of the bones, which in these days of aseptic technic is the best, but is not always to be recommended.

On account of better technic in the treatment of fractures we see to-day comparatively few cases of pseudarthrosis as compared with former times. With the help of the skiagraph we are able so perfectly to reduce a fracture that pseudarthrosis from malposition of the fragments is very rare, indeed. Therefore, we may place almost all cases of pseudarthrosis in the second group, in which the system, from some known or unknown reason, has lost its ability to form callus.

The injection of blood in these cases is similar in its action to the injection of other irritating agents (tincture of iodine and oil of turpentine), the purpose of which was the same, namely, stimulation of the fractured ends to callus formation. The injection of blood is much better, because it is free from danger, it causes less pain, and is similar to the natural irritation, which results in union by callus.

In Bier's clinic, where this treatment is carried out, the technic is as follows: The site of the "false joint," as well as the arm from which the blood is to be taken, are rendered aseptic. By light application of a constricting band the veins of the skin are made prominent, then one of the largest vessels is punctured and 30 or 40 c.c. of blood are slowly drawn into the syringe. If any difficulties are encountered in puncturing the vein, the operation may be facilitated by a small skin incision. The syringe must be large and its needle sufficiently strong in order that the blood may be readily forced through it. Above all, the syringe must be very strong, since much force is necessary to inject the blood into the hard scar tissue.

It is desirable to inject the blood into the space between the fractured ends of the bone, and one may be assisted very much by a skiagram. Not only must one inject the blood between the fractured ends, but also around them into the places, where, from periosteal and paraosteal connective tissue, callus formation shall take place.

It is important that one should immediately inject the blood after its removal, using the same syringe into which it is drawn, in order that it does not coagulate in the syringe. The venous blood does not coagulate as rapidly as arterial blood.

Before using the syringe it should be proven to be in perfect order, normal salt solution being used for that purpose, as it tends to prevent coagulation in the syringe.

4. There is a bibliography, omitted from THE JOURNAL, which appears in the author's reprints.

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.