This article was downloaded by: [130.132.123.28]

On: 03 February 2015, At: 07:18

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 5

Publication details, including instructions
for authors and subscription information:

<http://www.tandfonline.com/loi/tnah11>

The Mediterranean species of Fierasfer

Prof. C. Emery

Published online: 07 Oct 2009.

To cite this article: Prof. C. Emery (1882) The Mediterranean species of Fierasfer, Annals and Magazine of Natural History: Series 5, 9:50, 137-138, DOI: [10.1080/00222938209459011](https://doi.org/10.1080/00222938209459011)

To link to this article: <http://dx.doi.org/10.1080/00222938209459011>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever

caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

The observations of F. E. Schultze on *Quadrula symmetrica*, and those of Leidy on *Trinema acinus*, show that in these two genera the same things evidently take place as in *Euglypha*; that is to say, the platelets protecting the body are produced in the interior of the parent individual, and afterwards conveyed round the body of the daughter individual. M. Gruber has almost completely traced these same phenomena of division in *Cyphoderia ampulla*, the carapace of which is not formed of a comparatively restricted and tolerable constant number of plates, but rather of an infinity of little particles which gives this envelope the aspect of shagreen.

The phenomena of division seem to be nearly the same in the *Arcellæ*.

In the Monothalamia with carapaces formed of foreign materials, such as the *Diffugiæ*, which are covered with grains of sand &c., individuals have been observed united by their buccal poles; and this state has been regarded as the result of conjugation. According to Dr. Gruber this interpretation is incorrect, and the individuals thus joined must be the product of a division on the point of completion. With Bütschli he assumes that these Rhizopods first of all introduce into their bodies the foreign substances which are to serve for the formation of the envelope. The sand-grains, Diatoms, &c. are then transported to the outer surface of the newly-formed individual, just in the same way as the platelets of the *Euglyphæ*, *Quadrulæ*, &c.

The forms which are protected by an inflexible chitinous carapace also present the same mode of multiplication. This would seem to be proved by Dr. Gruber's observations on *Microgromia socialis*, and Schneider's on *Diffugia cinchelys*.

On the other hand, in the genera which have an envelope formed by a flexible membrane adherent to the sarcode of the body, division takes place, as in the *Amœbæ*, in the mode that may be denominated normal, because it is that which is by far the most frequent in animal cells. A constriction is produced in the middle of the body, and causes the formation of two individuals.—*Zeitschrift f. wiss. Zool.* xxxv. p. 431, & xxxvi. p. 104 (1881); *Bibl. Univ., Archives des Sciences*, December 15, 1881, p. 624.

The Mediterranean Species of Fierasfer. By Prof. C. EMERY.

Fierasfer acus, the commonest species in the Mediterranean, attains a length of 19 centim. (about $7\frac{1}{2}$ inches), and takes up its abode preferently in the large Holothuriæ, such as *Holothuria tubulosa* and *Stichopus regalis*. The author has frequently observed the process adopted by the little fish for introducing itself into the body of the Echinoderm. It commences by examining the whole length of the latter until it has discovered at which end the anus is situated. It places its muzzle against this orifice, and then, at the

moment when the sphincter dilates to allow the escape of the water which has served for respiration, it bends round quickly, and gliding its slender tail along its body, passes it in an instant into the cloaca of the Holothuria. This first step taken, the rest of the operation may occupy more or less time. A small *Pierasfer* attacking a large Holothuria sometimes succeeds in making an entrance at once. But should there be any disproportion of size the parasite waits for the respiratory stream to dilate the anus, and then pushes further in; and it is only by long-continued efforts that it finally enters. Prof. Emery has seen as many as seven of these fish successively enter the body of the same individual.

The *Pierasfer* lodges at first in the respiratory tree of the Holothuria, which opens into the intestine not far from the anus; but it is also found in the perivisceral cavity, because the respiratory tree is most frequently torn by the efforts of the little fish, especially when it receives several of them at the same time. The *Pierasfer*, however, is not a true parasite feeding at the expense of its host, but gets its nourishment from the sea by pushing its head out of the Holothuria. The position of its anus, which is placed very near the head, also enables it to evacuate the faecal matters and the sexual products without quitting its domicile.

This singular fish consequently makes use of the Holothuria as a habitation, or as a refuge from its enemies. It is therefore what we may call a *commensal* in the words of Van Beneden, or, as Prof. Emery expresses it, a lodger-parasite (*inquilinus*).—*R. Accad. dei Lincei, Atti*, ser. 3, vol. vii. 1880; *Bibl. Univ., Archives des Sciences*, December 15, 1881, p. 627.

Mode of Capture of Lizards in Southern Europe. By Dr. T. EIMER.

In my memoir on *Lacerta muralis cærulea* I described the peculiar method, usual in Italy, by which the boys there catch lizards: they make a noose at the end of a long stiff haulm of grass, and fill this with saliva so as to appear like a shining mirror. They hold the grass-haulm towards a lizard, which, being very inquisitive, comes nearer and nearer in order to examine the apparatus, and in the midst of its curiosity easily allows the noose to be drawn over its head.

The celebrated statue of the Sauroctonus*, as is well known, represents a youth, still of tender age, who, leaning with his left arm upon the trunk of a tree, and holding in his right hand a piece of a rod, in a watchful attitude follows with his eyes a lizard running up the trunk of the tree, with the object, as the archæologists think, either of tickling or transfixing it with the above-mentioned rod, as with a dart, a fragment of which the rod would represent. The latter opinion, so far as I know, relates to the statement of Pliny†,

* *Σαυροκτόνος*, lizard-killer.

† Hist. Nat. xxxiv. 70.