Dr. Watson mentions, in his "Lectures on the Practice of Medicine," that he has known two or three cases in which the diseased artery all at once cracked across (the inner and middle coats), and death very rapidly occurred. In one of these the crack extended round a considerable part of the circumference of the aorta, and looked like a clean cut made by a sharp knife. Dr. Wardrop relates, in his monograph, "On Diseases of the Heart," that when, from any powerful muscular effort, there is a strong regurgitating stream of aortic blood, it has happened that either the semilunar valves have been torn, or when they have been able to resist the retrograde stream, (as occurred in the present instance,) the coats of the aorta have given way. He gives a couple of instances in illustration.

The following case, therefore, is one of considerable interest. The notes were taken by Mr. Jackson, one of the hospital pupils:—

William S—, aged forty-five, was admitted into the hospital on the 3rd of July, 1860. This patient, a stout and short-necked man, was driving a spring-cart, and while in the act of rising to whip the horse he fell backwards. He was brought into the hospital a few minutes after, and when first seen he was insensible, and was kept on the sofa with great difficulty, as his body was constantly writhing about. He frequently cried out in a sharp, abrupt manner, gasping and struggling violently at the same time. His respiration was irregular, his face livid and perspiring, pupils dilated, and jugular veins turgid. From his short, thick neck, livid face, insensibility, and dilated pupils, some concluded he was suffering from apoplexy; but as there was no paralysis, and from the anxiety and constant convulsive struggling, the diagnosis of cardiac syncope was made; and on feeling for the pulse over the radial artery, none was detected. The hand was then applied to the chest, to ascertain if there was any impulse of the heart, but this was attended with a similar negative result. The skin was covered with a profuse cold sweat, his struggles became fewer and weaker, his respiration more and more interrupted, and he died within a few minutes after admission, never having recovered consciousness. His mouth was fixed in a sardonic grin.

At the autopsy, the scalp was found to be full of blood, the arachnoid membrane congested, and the puncta vasculosa numerous. The lateral ventricles contained about three ounces of limpid serum. No clot was found in the brain. Right lung healthy, left adherent, but substance healthy. The pericardium contained about seven ounces of bloody serum, and a clot ten ounces in weight, forming a thick layer, moulded to the surface of the heart and great vessels. This clot could be traced backwards between the appendix of the left auricle and pulmonary artery to the posterior surface of the aorta, where the blood forming it seems to have issued from an oval valvular rent in the walls of that vessel. The aorta, from its origin to the commencement of the arch, was greatly dilated and spindle-shaped, its greatest circumference at the middle being about six inches and a half. The coats of the artery at this point were extremely thin and brittle; the lining membrane was either rugous, opaque, tubercular, or studded with calcareous deposits. On the posterior wall of the aorta, in its ascending portion, was a narrow, rather elongated rent in the two inner coats of the vessel. From this opening the blood had escaped between the middle and external coats, but chiefly into the pericardium, through the small, irregular valvular opening already mentioned.

ST. GEORGE'S HOSPITAL.

ANEURISM OF THE AORTA, AND GENERAL DISEASE OF THE HEART; FATTY DEGENERATION OF THE LATTER; FATAL RESULT.

(Under the care of Dr. PAGE.)

IN dissections of aneurisms of the aorta and its branches, the lining membrane of the contiguous bloodvessels is commonly found affected in various degrees by atheromatous ulcerations in different states of development. This peculiar condition in most cases accounts for the presence of the aneurism, which originates within the thoracic cavity in a similar manner to that in the vessels of the abdomen or extremities. It is probable that if any intra-thoracic vessel should suddenly give way, as happened in the preceding instance, death would speedily occur. Fortunately the process of ulceration is slow, and the aneurismal dilatation of the outer coat, at first insignificant, when the two inner coats have been absorbed gra-

dually becomes augmented. The following example (for the notes of which we are indebted to Dr. W. H. Dickinson, medical registrar to the hospital) is an illustration of a double aortic aneurism, originating from the causes above referred to—causes in which the heart participated. Two aneurismal pouches were developed upon the aorta, one above a valve-flap, as large as a filbert, and the other an inch higher up, of the size of a small apple.

As the natural elasticity of any artery is diminished or destroyed by the morbid changes which it undergoes when invaded by atheroma, the consequence of this process on the aorta is shown by its dilatation; the "perpetually recurring strain" upon it of the blood coming from the left ventricle, causes its calibre to be enlarged. Such was the result in the case of rupture of the aorta. In the present instance it existed to a slighter extent; but this was compensated by the yielding of the weaker parts of the vessel, and the formation of the aneurisms mentioned.

When aneurismal pouches form very close to the commencement of the aorta, their detection is a matter of extreme difficulty. It is otherwise when they occur a little higher up, or upon the transverse portion of the arch; for their increase in size then gives rise to symptoms which at a later period point to their existence.

John G—, aged thirty-two, admitted September 26th; is a tripe-dresser, and attributes his illness to sleeping in the open air for several nights six years ago. He then appeared to have caught cold. He suffered from rheumatic pains and shortness of breath, but he was not confined to bed, and soon got better. He never had rheumatic fever. During the last three years he has suffered much from palpitation of the heart, and has latterly been unable to rest, except on the right side. He never had any dropsy. When admitted, he had a pallid countenance, and suffered much from dyspnoea. He had some cough, which had come on three days before. On examining the chest, bronchial râles were heard all over the lungs, which obscured the sounds of the heart to a certain extent. The præcordial dullness was much increased, but the sounds of the heart were not unusually distant. At the apex a rough murmur could be heard in the place of the second sound, the systole being natural. The murmur increased in intensity towards the upper part of the heart, and attained its maximum at the left edge of the sternum, about the level of the second rib. In this position only there was a little roughness with the systole. The impulse of the heart was much extended; the blood could be seen regurgitating into the veins of the neck. The pulse was 92, regular, very weak, but with a sharp jerk. The hepatic dullness extended beyond its proper limits, reaching into the epigastrium. There was no ascites. He was allowed a roast slice for dinner, had a night draught, and took a mixture every six hours of chloric ether, eight minims; aromatic spirit of ammonia, tincture of henbane, of each half a drachm; and camphor mixture, an ounce and a half.

Sept. 27th.—The palpitation is the source of much discomfort, and some tincture of arnica was ordered to be applied to the upper part of the præcordial region, where the impulse and dullness were not perceived. The cough got much better—in fact, it had quite ceased after a few days under the influence of tincture of squills and linctus occasionally. In other respects he became much worse.

29th.—He became rapidly cedematous, and his strength began to fail.

Oct. 1st.—Some wine was now ordered. His bowels were generally confined, and he suffered from piles. Confection of senna was given every other morning. He grew weaker as the dropsy increased; he became delirious, especially at night, yelling and screaming in such a violent and objectionable manner that he had to be removed to a ward by himself. This condition did not last long. He lapsed into a state of semi-coma, and quietly expired on the 9th.

Post-mortem examination, nineteen hours after death.—The veins of the surface were congested, and the superior half of the body was very livid, especially on the upper surface. There were partial old adhesions of the right side of the thorax. The lungs were healthy, except slight emphysema of their edges. The heart was of rather large size. The pericardium was tolerably firmly adherent at the upper part to the base of the heart, and to the aneurism of the aorta, presently to be described. The muscular fibres of the heart showed considerable fatty degeneration, and there was also a large quantity of oil-globules amongst the fibres. The blood was fluid, and the lining membrane of the heart and arteries deeply blood-stained. The right auricle was of very large size; the tricuspid orifice admitted all the fingers of the one hand, and the valve-flaps

seemed unable to close it. The mitral valve was healthy. Above one flap of the aortic valves there was an abnormal pouch, about large enough to contain a filbert, and empty. The corresponding valve-flap was much stretched, and irregularly thickened. Just above this (at a distance of about an inch) was a round opening in the wall of the artery, about the size of a florin, leading to a pouch the size of a small apple, lying between the right auricle and the artery. The liver and kidneys were healthy.

GUY'S HOSPITAL.

OBLITERATION OF THE SUPERIOR VENA CAVA; INFLAMMATION OF THE RIGHT AURICLE, WITH ATHEROMATOUS DEPOSITS; FATAL RESULT.

(Under the care of Dr. OWEN REES.)

ONE of the consequences of the pressure of a thoracic aneurism is obstruction of the superior vena cava. This may proceed to such an extent as to end in actual obliteration, when of course a fatal result soon ensues. From the symptoms present (which we regret are not given more in detail), a tumour was suspected to be the cause of the obstruction in the subjoined case. The autopsy, however, showed that complete obliteration of the superior cava had taken place, not from a tumour, but probably from extension of inflammatory action from the right auricle. This is a peculiar and rare cause of obstruction in this particular vessel.

The diagnostic signs of serious interrupted circulation through the superior cava are: anasarca of the entire upper half of the body, which may be enormous; the surface of the thorax and abdomen is covered with distended and tortuous veins; and the breathing and circulation are greatly embarrassed. All these were present, to a tolerable extent, in the patient whose case we give from the notes of Mr. D. M. Sergeant, clinical clerk. In other instances these symptoms are so much exaggerated as to cause the upper half of the body to appear as if stuffed, with the arms projecting from the sides from oedema, the eyes starting from their sockets, and the integuments ecchymosed. A remarkable case, in which the vena cava became impervious from the gradual pressure of an aortic aneurism, is related by Dr. Watson, in his "Lectures on the Practice of Physic."

Elizabeth L—, aged forty-eight, married, mother of one child, admitted on October 6th last. She states that she was taken ill about four months ago, with what she considered to be ordinary cold, accompanied, however, with swelling of the face, and considerable difficulty of breathing. Up to the time of seizure her general health had been remarkably good. She complained of great pain of the left side, had great difficulty of breathing, and much congestion of the face and upper extremities. The urine was not albuminous. Dulness was present posteriorly on the right side of the chest, and the respiratory murmur was nearly absent. There was ægophony; some dulness was present also on the left side anteriorly. The diagnosis was some tumour pressing upon the superior vena cava. She was ordered three grains of bromide of potassium, and half a drachm of liquor potassæ, in infusion of cascarrilla, three times a day.

Oct. 14th.—Congestion of the face is greater; arms very painful; some swelling of the left leg.

Nov. 2nd.—Died very calmly.

Post-mortem examination.—Much effusion in right pleura; heart healthy, except the right auricle, which presented traces of inflammatory action, covered by atheromatous deposit. The superior vena cava was completely impervious, organized deposit having obliterated the cavity for at least an inch and a half from the auricular opening. Fibrous ante-mortem clots extended up into both jugulars. The right lobe of the liver congested, having a deep sulcus across it. The other organs, with the exception of the uterus, which contained a small fibrous tumour, were healthy.

THE COMPARATIVE APPRECIATION OF SOUND BY HUMAN EARS.—The sensibility of the human ear for the pitch of musical notes has been recently illustrated. M. Fessel, of Cologne, on testing the new Parisian tuning-fork, observed that he heard differently with his two ears, the note heard with his right ear being somewhat higher than that heard with his left. On examining his musical friends, he has not yet found one, even amongst part musicians, whose ears are precisely alike in the estimation of the pitch of musical notes.

Medical Societies.

OBSTETRICAL SOCIETY OF LONDON.

WEDNESDAY, DEC. 5TH, 1860.

DR. TYLER SMITH, V.P., IN THE CHAIR.

THIS being the last meeting of the second session, the following gentlemen were appointed to audit the accounts for the year 1860:—Dr. Braxton Hicks and Mr. J. T. Mitchell, F.R.C.S.

Dr. TANNER exhibited a

LARGE COLLOID TUMOUR OF THE OVARY.

This tumour, which weighed upwards of 8 lbs. avoirdupois, was removed from a patient of Dr. Robert Fowler. The woman was thirty years of age, and was doing very well, eleven days having elapsed since the time Dr. Tanner operated. The operation was employed as a last resource to save life, since the patient was not only suffering from the tumour, but also from ascites produced by it. Five days before the performance of ovariectomy tapping was resorted to, and fourteen pints of ascitic fluid withdrawn; yet at the operation it was found that about five pints of fluid had been again secreted. With regard to the operation itself, there was one point deserving of mention,—namely, that after applying ligatures to the pedicle, the tumour, instead of the pedicle, was cut through, and a portion about the size of a small hand left attached, which was retained outside the abdomen. By this proceeding the use of the clamp was rendered unnecessary, while all risk of hæmorrhage was avoided. The latter point was of no little importance, since bleeding from the pedicle had been the cause of death in no less than 16 per cent. of the fatal cases. The edges of the wound were carefully brought together with twine sutures; these latter not being passed through the peritoneum.

In reply to a question from Dr. Tanner,

Mr. SPENCER WELLS said he had attached the greatest possible importance to the practice of passing the metallic sutures or hare-lip pins through the peritoneal edges of the wound. He had been led to originate this practice by observing the inner aspect of the wound in the first fatal case of ovariectomy which had occurred in his practice. The sutures had been passed through integuments and muscle, but not through the peritoneum; and the consequence was that a raw surface of considerable breadth was only partially protected from the general peritoneal cavity by a portion of intestine, which was already adhering to it, although the patient died the day after the operation. Had she lived longer, there could be no doubt that pus or other secretions from the wounded surface would have been poured into the cavity of the peritoneum. Struck by this observation he (Mr. Wells) made a number of experiments, in concert with Dr. Richardson (who narcotized the animals by puff-ball smoke or chloroform), upon guinea-pigs, rabbits, and dogs, opening the peritoneum and closing the wound in some cases by sutures which included the membrane, and others which only passed near it, and killing the animals at various periods after the operations, which were made as accurately comparative as possible. The result was that nearly all the animals recovered; but in those where the more superficial sutures were used there was a portion of the parietes left destitute of peritoneum, and this defect was supplied by adhesion of intestine or of omentum. In two cases where omentum became adherent, the motions of the dogs were materially interfered with. In those cases where the peritoneal edges of the wound had been included, the union was so perfect that it was difficult to detect the line of union two or three weeks after the operation. This was explained by the observation of a case in which a woman died within two days after ovariectomy. The hare-lip pins which had passed through the peritoneal edges were quite concealed from view and protected from contact with intestine, partly by folding of the peritoneal edges together, and partly by effused lymph. Mr. Wells said he had preserved many of the most striking specimens, and would be happy to show them to any gentleman who was interested in the subject.

Dr. TYLER SMITH had recently attended a case which strongly exemplified the good effects of bringing the edges of the peritoneum together by the sutures. After union of the external wound and the removal of the sutures, suppuration took place along the whole line of incision, extending through the entire thickness of the abdominal wall. The peritoneum had, however, healed so perfectly that no pus found its way into the peritoneal cavity. In this case, silk sutures had been