Clinical Remarks

ON THE

TREATMENT OF TIGHT URETHRAL STRICTURES.

By REGINALD HARRISON, F.R.C.S.,

SURGEON TO THE LIVERPOOL ROYAL INFIRMARY, AND LECTURER ON CLINICAL SURGERY IN THE VICTORIA UNIVERSITY.

THE circumstances under which retention of urine occurs in the male are so varying that it is impossible to lay down general rules for its management, other than, where practicable, the employment of the catheter with as little delay as possible. In the majority of instances where retention happens in connexion with a urethral stricture the relief thus afforded is not only sufficient for the emergency, but serves as the starting-point for the local treatment which in due course is to follow. Hence in those cases where, from a variety of causes, retention of urine supervenes upon the slighter forms of urethral stricture, the introduction of the catheter followed by the appropriate local treatment—such, for instance, as gradual dilatation—generally suffices. The cases now under notice differ importantly from these, both in the conditions under which retention presents itself and in the treatment to be employed. I refer to the severest forms of organic stricture, where, by the conversion of more or less of the urethral wall into a mass of tissue in every respect resembling the worst kind of cicatrix, not only has the urethra been reduced to the smallest dimensions compatible with the escape of urine from the bladder in drops, but the bladder and associated parts are also in a condition of active inflammation. The processes to which I have thus referred are of a gradual kind, and, as a rule, the result of months, or even years, of continued neglect. It is remarkable how often we meet with instances of this kind, where persons have gone on, hoping, I suppose, against hope, and quite regardless of the increasing difficulty with which their urine was obviously voided. The conditions here are very different from those where retention supervenes upon spasm of a more or less transient kind. Taking the cases I am now passing under review, in the course of time one of two things happens: either abscess in immediate relation with the stricture, which may open either with or without extravasation of urine, or complete retention of urine. It is with the latter contingency I am now dealing. The best that can be hoped for under the condition thus sketched out is that the surgeon will be able to pass a small catheter through the contraction, and thus relieve the urgency of the retention. Let us now take a case where the operator has succeeded, probably under an anæsthetic, in passing a No. 1 metal English catheter through a densely contracted stricture of half an inch or more in length, and that he has verified his position (1) by the escape of urine in drops by the catheter, and (2) by his finger in the rectum. The instrument is firmly grasped by the stricture, and there can be no doubt as to the correctness of its position. Will this expedient serve to meet all the exigencies of the case, even supposing that the urine can be slowly drained off in this way? To tie such an instrument in the bladder, and to leave the urine to drain away indefinitely, is, I am sure, a most hazardous proceeding, though it may be accepted as an alternative. In all these cases more or less acute cystitis has been in existence for some days, and the urine is high-coloured, offensive, and loaded with deposit, and what could thus be done would be to drain off the most fluid and least harmful of the contents of the bladder, leaving the more poisonous remainder to add to the mischief already going on in the bladder, ureters, and kidneys. Such a plan usually ends with the death of the patient in a few days of what may with truth be called a catheter fever.

A considerable experience in the class of cases coming under this category has taught me very forcibly that the introduction of the small catheter must here be regarded only in the light of the initial proceeding. In addition, the bladder must be put in a position for thoroughly draining itself, otherwise the patient's chances of recovery are

exceedingly small. Where there is a perineal tumefaction indicating that suppuration is occurring, no question can then be entertained as to the immediate necessity for perineal section. The small catheter will serve as a guide, and upon it the stricture can be completely divided, though the proceeding is easier when a grooved staff, however small, can be substituted. Where there is a fear as to the possibility of being able to make this change, the catheter should be utilised as a guide for the incision. I have often used a small metal catheter under such circumstances, and have never had much difficulty in completing the operation of perineal section, with division of the stricture, to my satisfaction. When the perineum is opened in the median line, far more can be done with the index finger as a guide than is generally supposed, so long as we have the means of ascertaining the precise line the urethra takes. When, however, there is no perineal tumefaction, and the catheter is merely persistently grasped within a hard fibrous stricture, other means must be taken at once to bring up the calibre of the urethra to something like its natural dimensions, so that the bladder may be kept thoroughly drained of all its contents. The mode of doing this will be best illustrated by the narration of the following selected cases, where the same object was aimed at, but in different ways. I have very little preference for any one way of bringing the contracted urethra up to its normal dimensions so long as thorough urine drainage and cleanliness can be obtained. These are the most important factors in the antiseptic surgery of the urinary organs.

The first case I will relate is that of a gentleman who had been suffering for some years from a traumatic stricture of the urethra following an injury to the perineum. He had had many attacks of retention when abroad. Catheterism was always difficult. He had been aspirated several times above the pubes, and on one occasion before I saw him had perineal section performed. The last operation had not been a success. When I first visited him he was not suffering from retention, but only a No. 1 English metallic bougie could be passed through an extremely hard stricture. I then advised that perineal section should again be practised, but upon somewhat different lines from the proceeding he described to me. However, nothing was then done, and I heard no further about him for some months, when I was asked to see him for urgent retention of urine, where catheterism had been attempted, but without avail. Under chloroform, and with difficulty, I passed a No. 1 metallic catheter, and this, with much patience, was gradually increased up to a No. 4 English. This, with some misgivings on my part, was tied in. The stricture, however, was so tight that I feared the catheter might give way in my attempt to increase the size. In the early morning the instrument became plugged, and was eventually withdrawn. Further attempts were made at catheterism, but without avail, and I was again summoned. Returning, I took with me a fine stricture stretcher (elsewhere described¹), by means of which I succeeded in dilating the stricture up to a No. 13 English. A full-sized catheter was then passed, and the bladder emptied of its most offensive contents. As the patient lived in a rather inaccessible position, the catheter was tied in and retained for three days. This process of divulsion was the course of proceeding which in the first instance ought to have been practised; as I had no instrument of this kind with me sufficiently fine for this purpose, I reluctantly contented myself with retaining the small catheter. I saw this patient a month afterwards. His urine was voided in a full stream, and was healthy. I passed a No. 12 bougie for him, and taught him how to do this for himself.

The next case is that of a man aged about twenty, who presented himself at the infirmary a few months ago with a very tight stricture of some years' duration, for which he had received no previous treatment whatever, in spite of the gradual diminution in the size of the stream. He appeared to have been drinking heavily. His stricture was very fibrous and contracted, and I could not pass even the smallest metal catheter. However, I succeeded in getting through with the fine filiform bougie of a Maisonneuve's urethrotome. In this way I was able to perform an internal urethrotomy, and then to empty his bladder of a quantity of most offensive urine with a large catheter. The patient would not remain in hospital; he was, however, brought back in forty-eight

¹ Lectures on Urinary Diseases, third edition.

hours in a state of delirium tremens, but passing his urine freely. He remained in the hospital for ten days until this complication was over, when he was discharged, passing urine normally and able to introduce a full-sized bougie for himself.

The third and last case I shall introduce was also recently admitted into the Royal Infirmary. It was that of a man aged thirty-two years, who had long suffered from stricture, and had undergone some operation for it in hospital seven years previously. The meatus of the urethra was involved in a cicatricial mass connected with the foreskin, and in addition there was a long fibrous subpubic contraction. He had extreme retention of urine, which had evidently been only very partially relieved by the introduction of a small catheter prior to his admission. With much difficulty, owing to the double nature of the obstruction, I could only succeed in passing a No. 1 metal Lister's bougie. This was most firmly grasped. Under an anæsthetic I managed to introduce the entire series. The bladder was then emptied of much offensive urine by a large catheter. The stricture at the orifice was freely divided with a probe-pointed bistoury. This patient convalesced rapidly, and was able to leave the infirmary in the course of ten days, passing a full-sized instrument.

In connexion with these cases there are some details to which I could refer, as I am satisfied that much of the success attending them, as well as of others of a like kind which have come under notice, is due to their recognition. Every care was taken to prevent the occurrence of any form of urethral septicæmia, as indicated by the occurrence of rigors and fever following the performance of the operation that was selected. This consisted in the employment of local and general measures. As soon as the bladder was emptied with a large catheter, it was thoroughly washed out with a perchloride solution of 1 in 5000; three or four ounces of this fluid were left behind, so that the first urine which was spontaneously passed was largely impregnated with this antiseptic. As a rule, no catheter is tied in, nor can I remember an instance where its subsequent introduction became necessary. Before the patient is sent to bed the urethra is distended with carbolic oil (1 in 20), which is well rubbed into the part where section or divulsion may have taken place. Five-grain doses of quinine are given every four hours; in some instances boracic acid appears to have answered equally well. They seem, by sterilising the urine, to prevent rigors and fever.

For rapidly dilating very fine tight strictures, I have found Banks's filiform bougie of great service. It is an instrument that is well known in America, but not in this country. It is made of highly polished whalebone, on the principle of the wedge, and can be used with great safety. When completely passed through the stricture, a much larger-sized ordinary catheter or bougie can usually be made to follow easily. These instruments, of which there are several sizes, have been supplied to me by Messrs. Tiemann of New York.

It is not to be assumed that the treatment this class of cases immediately received, as illustrated in the three cases selected, represents in each instance that which was considered best so far as the future of the stricture was concerned. On the contrary, I feel sure that in two of the illustrations perineal section with complete external division of the stricture would have given far more reliable permanent results, and may yet have to be brought under consideration. Retention of urine, however, sometimes happens where the surrounding conditions render the performance of such an operation, on an emergency, quite out of the question. It is in view of this that I have ventured to illustrate how in my own practice it has been found possible to tide over the difficulties which attend this complication when it occurs in connexion with the worst forms of organic urethral stricture.

Liverpool.

THE CONVALESCENT HOME AT GILMERTON.—This new home for children was formally opened by the Rev. Dean Montgomery on the 18th inst. It appears that it was not exactly the opening of a new institution, but simply the transference to another and more favourable site of the home, which had been in existence for some time. The new building provides enlarged and improved accommodation, and has cost £1330, of which £200 remained to be paid.

A CASE FOR DIAGNOSIS.

By FRED. J. SMITH, M.B., M.R.C.P.,
MEDICAL REGISTRAR TO THE LONDON HOSPITAL.

By the kind permission of Dr. Warner, under whose care the patient was in hospital, and with the ready assistance of Mr. H. A. Debenham, the house physician, I am enabled to publish the following case, which I watched with the greatest interest, not unmixed with stronger feelings of baffled curiosity.

L. W.—, aged twenty-three, a female fancy worker, was admitted to the London Hospital complaining of a dull pain in the back and abdomen. The family history was without bearing on the case, except that the father was said to have died from a tumour of the lungs. The personal history showed an entire absence of any previous illness of sufficient severity to cause her to lie up. The history of her present illness was that the pain began somewhat suddenly in the lumbar region about three weeks only before admission, and was of such severity as to cause her to faint; for this she was treated by a general practitioner. On admission the patient was seen to be a well-nourished girl; slightly anæmic; very restless in bed; complaining of severe pain, which appeared to shift from place to place, at one time being in the middle of the back, and again appearing at the sides of the abdomen; it was increased by lying on either side, and by sitting or standing; the pain was of a throbbing character, likened by the patient to a "gathering." General superficial examination and palpation revealed absolutely nothing to account for the pain. An examination of the alimentary tract showed some slight digestive disturbance, with a furred tongue, discomfort after eating, with slight nausea and obstinate constipation. The heart and lungs yielded entirely negative results on examination, except for occasional dyspnoea and slight pain on taking a deep breath. The patient was treated with a saline mixture, and morphia was administered hypodermically. For the next fortnight no change took place practically in her condition; the pains were occasionally a little easier, but generally got much worse at night. On May 25th a thorough vaginal examination was made by the assistant obstetric physician, who reported that the uterus was freely movable, and, in fact, that there was no abnormal or pathological condition to be detected, either per vaginam or on bimanual abdomino-vaginal examination; but, at the same time, he advised that a surgical opinion should be obtained as to the condition of the spine, inasmuch as the patient seemed to refer the pain pretty consistently to that region. For some days after this she seemed and expressed herself as being much better, though the notes state that she was very hysterical. On June 5th the patient was again most thoroughly overhauled in an attempt to arrive at some definite grounds on which to base a diagnosis. The reflexes appeared entirely normal; the bladder and rectum were performing their functions in a perfectly healthy manner; passive movements of all kinds were borne without the slightest alteration in the severity of the pain; and, in fact, nothing whatever could be detected in the nature of a clue to the origin of the pain; and about this time it was almost definitely decided that the case was one of hysteria. During June and July this diagnosis seemed to be confirmed, in that the patient improved under disciplinary measures, though her physical condition was such as to cause considerable uneasiness, as the dyspeptic troubles, especially the constipation and the anorexia, still continued, and the patient lost strength and flesh very rapidly, notwithstanding the improvement in her habits and manners. Towards the end of August it became evident that the patient was dying, though the cause was still as mysterious as ever. On the 25th a loud mitral systolic bruit was heard, evidently due to a failing heart; and on the 28th she died, apparently from exhaustion.

The post-mortem examination revealed the following condition. On the inner side of the lower ribs, one inch to the right of the sternum, was a small mass of a soft, melanotic growth; all the lumbar and several of the lower dorsal vertebræ were affected with a similar growth and were very carious. The glands along the bodies of the vertebræ and those of the anterior mediastinum were enlarged, and some of them were blackened, as were also the pelvic and inguinal glands. The heart was dilated and fatty. The liver was fatty and anæmic; the lungs were œdematous. The intes-