little more than 1800 cubic feet of air to each. Connected with each ward, in the centre of the building, is a nurses'-room and dining-room. To each dining-room is attached a closet and refrigerator, and it is connected with the kitchen by a steam dumb-waiter. At the extreme end of each ward is a corridor, connecting it with the end wings, which contain each a bath-room, a hot-air bath, a foot and hip bath, a lavatory, and waterclosets. All these are so separated from the wards, as to be independently ventilated. Each ward, again, has two closets for linen, and the main hall on each story is provided with closets for patients' clothing. On the street face of each ward is a broad balcony, with seats built permanently into the windows. All corners and angles are rounded to aid ventilation. The first story ward differs from the others in having bidet baths for women.

The west side of the main hall in the fifth story contains a surgical ward for nineteen beds. It is 26 ft. wide, 78 ft. long, and 16 ft. 5 in. high, and allows 1800 cubic feet to each bed. Its general arrangements are the same as for the other wards. In the centre wing of this story is an isolation ward with two beds. The capacity of the hospital, exclusive of the children's wards, is 163 beds, including the private or paying wards. Each bed is connected with the nurse's-room by an electric communicator, which may at pleasure be connected with or disconnected from a bell. The button is so arranged that it can, at the discretion of the nurse, be within reach of, or withheld from the patient. Sockets have been built into the wall behind each bed, into which metal brackets can be hooked, which swing over the beds. From these handles are suspended beneath the patient, to enable him to raise himself or change his position without assistance. The nurses'-rooms are connected with the janitor's-room on the first floor by speaking-tubes. The janitor's-room is connected with the entrance doors, the office, the private wards, the first story and basement offices, the autopsy-room, and stables, by speaking-tubes or electric communicators. Each nurse's-room is connected with the office by speaking-tubes. The ward dining-rooms have speaking-tubes connected with the kitchen.

The east side of the hall in the fifth story is occupied by the operating theatre, and rooms connected with it. This theatre is very carefully arranged, and has a top and north light supplied through 261 square feet of hammered glass.

The sixth story contains the kitchen, pantry, store-room, laundry, drying-room, and linen closets. All are provided with the most modern and complete apparatus and machinery.

The seventh story is occupied by wash-rooms, engine and blower-rooms, water-tanks, exhaust-fan blower, and disinfecting-room.

To the administrative building, directly in the rear of the hospital main building, at a distance of 10 ft. 6 in. from it, two stories have been added, which are connected with it by an enclosed bridge at the level of the staircase landing. This is also approached on the first floor by a balcony running across the administrative building, and by a bridge. The lower part of these stories contains bedroom accommodation for thirty nurses in the training school of the hospital, also the necessary offices. The rooms are provided with separate entrances and corridors, leading respectively to the rooms for male and female nurses. The upper story forms a single room, with windows on all sides and two skylights, of which that in the centre is 22 ft. by 51 ft. in size. This is used as a convalescent-room, and is divided by partitions to separate the males from the females. This room is 64 ft. wide, 90 ft. long, and 18 ft. high. It is filled with rare plants, aquaria, and means for occupying and amusing the patients; is maintained at a uniform temperature; is light, bright, and cheerful, and is a beautiful summer and winter garden, of which it would be difficult to over-estimate the value, in accelerating the recovery of the sick and injured.

The arrangements for heating and ventilation are combined, and contain several ingenious features of novelty for the admission of fresh and removal of foul air in all parts of the building with the avoidance of draughts and careful regulation of temperature. The whole is under control, and can be regulated with minute accuracy, even to the supply of each particular bed, which could, if necessary, be practically isolated. The windows in the wards are opposite to each other, and are so arranged as to supply air with the minimum draught when open. The hospital occupied two years in building, and cost 595,069 dollars. The architect is Mr. George B. Post, of Broadway, New York.

This truly excellent hospital, in the economy and per-

fection of its structural arrangements for an institution constructed on the multiple ward system under the same roof, is by far the most complete and perfect that I have seen in any country. Whether it will ultimately effect the object of eliminating septicæmic disease entirely from its catalogue of ills, due to local insanitary causes, remains to be seen, for it has been too short a time in effective operation to determine such a question. So far as ample space, prodigal ventilation, and the prevention of noxious agencies finding their way to the sick are concerned, everything appears to have been done which science could suggest and far-seeing philanthropy adopt. The provision of a permanent convalescent home within its own walls, attests the wisdom and forethought of its founders.¹

Another of the recent hospitals in New York, which is on a different principle, of which the basis is the pavilion plan, is the Roosevelt Hospital—a special endowment somewhat similar in character to that of Guy's in London: it was opened in 1871. It is devoted to poor and paying patients in equal proportions, and as it is at present, so far as I know, unknown generally in Europe, it equally deserves description. In the Soldiers' Home at Washington, built under the supervision of Dr. Billings, of the United States Army, the kitchen and laundry are on the upper floor, as in the New York Hospital, and the result has been equally successful in maintaining freedom from the usual pollution of air inseparable from the operations carried on in such places. The system of heating and ventilation is also somewhat similar.

In all recent additions to the older hospitals of the United States, the pavilion plan, pure and simple, has been adopted, and nearly all the surgical pavilions are single-storied, as shown in the Sturges Pavilion of the Bellevue Hospital, of which I published an account in the *Builder* in January last. The first of these, I was informed, was constructed in connexion with the Presbyterian Hospital at Philadelphia, where there are at present two wards on this plan.

By far the most important, however, of all the hospital constructions in the great Republic is the Johns Hopkins Hospital at Baltimore, which is now not quite half built. It is the outcome of a munificent bequest, is to be attached to a great medical school, and is being arranged under the advice and supervision of Dr. John Billings. It stands on fourteen and a half acres of ground, bounded by streets on all sides, and is strictly on the pavilion plan, complete in all particulars. In it are to be illustrated all the most advanced views in hospital construction and arrangement, and it will probably still take some years to complete. Hence it will be better to defer an account of it to the period of its finishing and occupation, as changes will doubtless be made in many of the details, during the progress of the building. So far as it had gone when I saw it in September last, it appeared to me to promise to become one of the finest institutions of its kind and class in existence.

(To be continued.)

HÆMORRHOIDS OR "PILES."

By JOHN GAY, F.R.C.S.

THE study of hæmorrhoidal swellings, or piles, cannot lead to any very satisfactory results unless based upon, and associated with, an exact knowledge of the venous system, not only as to its local or special, but its general or common, anatomy. Such a remark may be said to be trite; yet I cannot but think that there are treasures of fact concealed beneath our relatively slender and actual acquaintance with that system, both anatomical and physiological, that would, if realised, largely augment the sum of our practical resources in remedying the diseases that appertain to its various parts.

For any very satisfactory accounts of the anatomy of the vein system we have to go back to the works of Vicq d'Azyr, Walter, and Loder. Since these were published its study has gradually relaxed, so that now in our best works it is entirely disposed of in a very few pages, and the most

¹ To understand the arrangements of this institution properly needs detailed plans of every portion, for which it is manifestly impossible to find room in THE LANCET.

complex and interesting of the phenomena appertaining to the circulatory system have had the smallest amount of attention bestowed upon them, whilst the simplest have exhausted whole volumes. Without such an amplification of the means of studying these curious products—piles—neither their *raison d'être* nor the anomalies of their clinique can be fully and satisfactorily explained. Thus portal congestion and allied hepatic affections have been held to be a constant source of disturbance to the currents in the hæmorrhoidal veins, and therefore accountable for the diseases that befall them; and anatomy has been called upon to explain the etiological relationship. But clinical observation shows that the cases of piles traceable to hepatic disease are comparatively few; and in searching for their etiological antecedents the ground comes to be enlarged as we go, and with it the difficulty of tracing them to any common cause; for, as Mr. Allingham says, "they happen equally to the robust and the healthy, to the rich and the poor, the active and the sedentary," and I may add, without fear of contradiction, to the young and the old, to the abstemious as to the indulgent, and I think to the healthy as to the diseased, for I have seen them repeatedly in persons below the age of twenty who have led unimpeachable lives, and are otherwise without trace of disease—perhaps, most frequently in young married women; still, I have seen them in young people of both sexes. Gather together, if you will, the data of an extensive clinical inquiry, and, as regards *remote causes*, heritage will, I think, be found to take a very considerable part in the rôle; and as to *direct*, perhaps those modern usages of society which induce inaction in the lower bowels—sedentary habits, prolonged railway travelling, free indulgence in meat diet and highly alcoholic drinks, and that inattention to those calls of nature which "busy lives" are apt to disregard as of little consequence, compared with those of the mart, the field, the tennis lawn, and the ball-room, will go far to complete it. Thus modern customs have a tendency to exclude from use the means of counteracting those disordered actions and diseased tendencies which they are too apt to engender. There is no doubt, as Mr. Quain remarks, that sluggishness of the bowels aggravates, whilst their free action tends to relieve, hæmorrhoidal malady. At the same time the converse also sometimes holds good, and diarrhoea has been known to be causatively associated with its production. If we take the anatomy of the hæmorrhoidal veins, so far as it is known, the relation of dilatation of their coats to portal obstruction is, nevertheless, to some extent obvious, and certain forms of "piles" may, in part, be thus accounted for. It might, indeed, explain some, but of others, such as those limited dilatations which are usually known as "external" hæmorrhoids, it does not give so satisfactory an account. The hæmorrhoidal, according to abstract or bare anatomy, form a portion of a vast pelvic plexus, in the formation of which other veins, such as the uterine, the vaginal, the vesical, the prostatic, and the spermatic, take their respective share. Together they constitute a network or congeries, and, as it occupies the lowest and most dependent part of the trunk and is very scant of valves, it would *a priori* seem impossible that there should be any disturbance in the larger branches or common trunks, causing reflux and peripheral blood-tension, that did not involve in its consequences more or less the whole of the series; at all events, that it should affect one exclusively. And this would seem to be still more improbable as the cause of disturbance, supposing, for instance, that it should be in the portal system, is so remote from the seat as more or less completely to cover the whole series. Such is not the case; and, as a rule, hæmorrhoidal dilatations or piles are not found to co-exist with like disturbances in any of the neighbouring or even associated organic plexuses. I have not met with a case in which varicocele could be said to have been an etiological co-ordinate of piles, to whatever cause the latter may be traced; nor have I found any satisfactory evidence of the existence, with the worst forms of piles, of allied over-repletion of other vein systems. I have seen the hæmorrhoidal and the vaginal veins very largely distended at the same time through the pressure of large pelvic tumours, or gravid uterus; but this result does not contravene the principle for which I am contending, that hæmorrhoidal veins may be, and usually are, exclusively dilated or varicose. After injury to these veins, as by operations, retention of uterine, as everybody knows, is common, but it is clearly due to sympathetic and not to any compensatory act on the part of the vesical plexus. And, certainly, obstruction of the cava, below the confluence of the hepatic veins, does not

produce piles, and if met in close contiguity to the heart, seems to find compensatory outlet for the pent-up streams rather in hæmorrhage from mucous surfaces, varicosity of superficial trunk veins, or general œdema; so far, anatomy simplifies hæmorrhoidal disorder.

But what are hæmorrhoids or piles? Perhaps this question should have been answered before, as well as in anticipation of, what is to follow. I will, then, just state that they may be taken to include every form of dilatation, of which certain veins are susceptible, that is liable to become permanent by pathological changes in the vein coats. And, as I shall endeavour to show, these changes are in conformity with like tendencies in certain homologous sections of the general system; and are, generally speaking, limited to two well-known forms—varix and varicosity.

Mr. Hunter seems to have dwelt on these disorders with considerable interest, as is clear, not only from the specimens that he collected and placed in his museum, but from his comments upon them. In his MS. cases in Surgery he remarks that piles would appear to be principally increased or varicose veins. "They are, however," he continues, "disease, and in this respect differ from the plexus retiform of the cavernous bodies, to which they assimilate, but only in this respect." Here there is seemingly an intuitive perception of the real nature of pile tumours, for Mr. Hunter does not appear to have gone beyond a superficial examination of them. In his collected specimens, the mucous membrane which covered the veins on which the piles are situated has not been removed in any case so as to expose them to close observation; and where they have been exposed *by section of the bowel or ulceration of the pile* they certainly do not generally exhibit that resemblance to the cavernous vascular tissue that Mr. Hunter speaks of. Lately, however, I have had an opportunity of dissecting some pile tumours, and in one or two cases have found this, what I shall term, trabecular variety. "The most singular thing," adds Mr. Hunter, "is the blood coagulating in them." In the specimen just alluded to, and in many others I have seen, the blood in the veins is coagulated. Thrombosis is, indeed, a very frequent condition of hæmorrhoidally affected veins.

The specimens which illustrate this disease in the Museum of the College of Surgeons begin with the number 1279, and are in continuity. They are well worth study.

In the first (Fig. 1, *d, d, d*) there is a circle of swellings near

FIG. 1.

a a, The cut edges of the bowel. *b b b*, Hæmorrhoidal tumours. *c c*, A circlet of such tumours. *d d*, A fold or fringe of mucous membrane. *e e*, The nates.

the extremity of the rectum, with a few more sparingly dotted about on the inner surface of the adjoining segment of the bowel. They do not extend above an inch or two in the upward direction—that is, they are limited to its lower portion. Amongst these little swellings a large vein takes its somewhat tortuous course, apparently in a direction towards the higher regions, away from that of the anal series, whilst a fringe or fold of mucous membrane, that does not invest a vein, hangs loosely from the submucous textures just above and within the anal orifice. As Mr. Hunter remarks of a similar specimen, "they (the pile swellings) stick out on the surface of the bowel like so many 'knobs,'"

whilst in the interstices the soft parts shrink in as intervening septa." These pile tumours assume certain figured forms. Thus in other specimens (1778B and 1277) they range as a circlet around and protrude from the anal orifice, whilst other such circlets are formed, with free intervening spaces, on the surface of the bowel above. This tier upon tier arrangement is common, but it is usually limited to two or three such tiers, whilst the usually upper ones appear as if they had been more or less broken up and the piles thinly scattered about. They are not often met with above the area of distribution of the tributaries of the lower hæmorrhoidal veins.

The several piles forming a ring or circlet often differ in size, and of two or more such tiers those of the lowest are usually the largest and most clubbed. Mr. Quain has given some excellent illustrations of piles in his work.¹ In one, however, a bunch of pile swellings is seen protruding below the verge of the anus, whilst above the sphincter and on the outer side of the muscular coats of the bowel, dilated and somewhat tortuous veins are seen running upwards, as though their tributaries had passed through the sphincter and had been strangulated into pile swellings by its fibres. But Mr. Quain has given no indication of continuity between the extra-muscular veins and the intra-muscular piles, a point which I think it would be difficult by dissection to establish. At all events, I have not been able to find any evidence of such relationship. Both anal as well as rectal piles, indeed, all pile swellings, may become the seats of ulceration, which often perforates their coats, exposing loose or firm coagula within. The intra-rectal ridges or folds of mucous membrane to which I have alluded are equivalents of those hypertrophied and warty-looking dermal appendages so often met with at the anal verge. These run more or less directly or spirally up the bowel from the rectal extremity (see Fig. 1, *d, d*). Some are loose and floating, but others enclose large or bloated veins or tributaries. In the former case the veins that formed the folds have retreated and are seen lying along at their base. During life in certain conditions of vascular excitement such veins are seen and give trouble, and sometimes even alarm, as they run along lazily beneath the mucous membrane, at times bloated and tense, well-nigh to bursting, losing themselves, however, in the main trunks above. They are usually transient, but might become permanent through hypertrophy of their coats. Such are the forms which piles assume in life on the undivided bowel.

(To be concluded.)

CASES OF FATTY EMBOLISM.

By C. W. M. MOULLIN, M.A., M.D. OXON., F.R.C.S.,
FELLOW OF PEMBROKE COLLEGE, OXFORD; SURGICAL REGISTRAR,
LONDON HOSPITAL.

IT is not a little singular that the subject of fatty embolism, which in other countries has attracted considerable attention during the last twenty years, should in England, with the most ample opportunity, have received only slight notice. In Germany there have been recorded upwards of 140 cases, in no less than 13 per cent. of which it was stated to have been the cause of death. In France Déjérine has published² 12 that occurred under his own immediate notice; but in England, with the exception of a brief mention a week or two ago of its presence in a case of primary amputation, under the authority of Dr. Hilton Fagge, the only instance that I can find is that given by Mr. Southam of Manchester in THE LANCET of July 10th, 1880. Within the last few months I have succeeded in ascertaining its presence in twelve out of fourteen cases examined, of which one or two, from the peculiar circumstances under which they occurred, were almost sufficiently exact to rank as experiments. They were taken consecutively from the post-mortem room without regard to the cause of death, and consequently present a considerable variety; indeed, there is only this one point common to them all, that, with one possible exception, there was no suspicion of its existence during life. The lung was examined in all but one, the outersurface of the right lower lobe in general being selected; the kidney in all but three. The liver and spleen, for

obvious reasons, were not touched, and the brain and pia mater, after two or three attempts, were given up as unsatisfactory, for it was found in the case of the former that the osmic acid solution would not penetrate a sufficient depth into its compact tissue, and in the case of the latter, especially when it was taken from the region of the medulla, the part of most interest, it was felt that there was some danger of confusion from the escape of myelin from the medulla of torn nerves. Besides, if the presence of fatty embolism could be shown in two such viscera as the lung and the kidney, there is nothing strained in the assumption that its distribution throughout the blood is fairly general.

A clean knife was, of course, used; the portions of tissue, which in the case of the lung may be of considerable size, were washed free from any blood that might be on their surface, and then placed in one-quarter per cent. solution of osmic acid for twenty-four hours, and afterwards hardened in alcohol, and cut and mounted in the usual manner.

The cases in which the presence of fatty embolism was not ascertained were two in number; the first, in which the cause of death was hæmorrhage following a wound, and in which the examination, from circumstances that I could not help, was not sufficient to prove a negative; and the fifth, acute croupous pneumonia without surgical injury. The remaining twelve were cases of simple or compound fractures of greater or less extent with the exception of the following: Case 3, pyæmia following acute periostitis of the femur, fatal on the eighth day, with pericarditis and numerous secondary deposits; Case 4, septicæmia following breaking down of fibroid tumours in the uterus after abortion; Case 6, ovariotomy, death in sixteen hours from "shock" combined with the oozing from very numerous torn adhesions; Case 8, removal of large sequestrum, death from erysipelas twenty-two days after operation; and Case 13, dislocation of the hip with much laceration of the soft textures round the joint and rupture of the spleen, fatal on the third day. Of the fractures, those of most interest were: Case 7, primary amputation of the thigh, high up, death in thirty-six hours without the patient having rallied, without rise of temperature or dyspnoea, the pulse remaining throughout small, frequent, and feeble; much post-mortem staining of tissues; Case 10, fracture of the fifth and sixth cervical vertebræ, death in sixty hours; and Case 9, fracture of the second and third cervical vertebræ with disorganisation of the cord. That the substance found under these circumstances in the form of drops intimately mixed with the blood in vessels of all sizes can be nothing but oil would be sufficiently certain alone from its reaction with osmic acid, even if it had not been independently proved by repeated experiments and observations carried out in other countries; and the same may be said with regard to the source whence it comes, for although it was at first regarded as associated in some peculiar manner with the secondary deposits of pyæmia, it has been shown clearly since that this was a mere coincidence, and that in all cases the oil is set free from the crushing of the fat cells somewhere in the periphery, especially when, as in the case of fractures, veins that cannot close are torn across at the same time. It is under these circumstances that it is found in greatest abundance, strictly in proportion to the extent of injury sustained, though occurring also when the softer textures alone are injured. From my own observations, though of course it would be absurd to formulate any general rule with regard to the frequency of its occurrence from the examination of fourteen cases only, yet, seeing that it was found in twelve, and that in the other two the examination of one was insufficient, and no surgical injury was present in the other, it may not unreasonably be admitted that the presence of fatty embolism after injury of any kind is, if not general, at least very much more common than is usually supposed; and that in the majority of cases it is not of very serious import.

There remains the consideration of the ultimate destiny of the fat globules when once they have entered the circulation, and of the symptoms to which they give rise during their passage through various organs, two points at least as of great importance, and not yet clearly made out.

There can be no question that although the oil extruded from the ruptured fat cells may be taken up, and probably is to a slight extent, by the lymphatic rootlets in the connective tissue spaces, by far the greater portion enters the circulation through the veins, and, as Dr. Hilton Fagge among others have pointed out, in the form of drops suffi-

¹ On Disease of the Rectum.

² Progrès Médical, 1879.