New Inventions.

PILLISCHER'S NEW "KOSMOS" MICROSCOPE.

THE annexed woodcut illustrates a new student's histological microscope, recently made and brought out by Mr. Pillischer, of New Bond-street. A well-arranged optical body and stage, with a revolving diaphragm, is supported upon a solid bronze base or foot, which is at once

original and most convenient, enabling the worker when using the stand to manipulate with the greatest ease. The fine adjustment is perfectly steady and sensitive, and will admit of the highest immersion objectives being used. The optical parts consist of one eye-piece and two objectives, a 1 in. and $\frac{1}{2}$ in., of excellent defining and penetrating power, giving amplifications from 30 to 380 diameters. In its simplest and cheapest form—viz., the stand, one eye-piece, 1 in. and $\frac{1}{2}$ in. objectives, and a well-made mahogany cabinet with lock, key, and handle—this microscope costs £5; but the stand is so constructed

that any additional objectives or apparatus, such as Abbé's or achromatic condenser, polarising and other apparatus may be fitted to it. The maker claims for this instrument both mechanical and optical excellence to a degree which has hitherto not been attempted in cheap microscopes of this class. We can recommend the microscope as a good and cheap one, well suited to students' work.

KANGAROO TENDONS.

To the Editors of THE LANCET.

SIRS,—With your kind permission I would like to draw attention to a few rather important circumstances respecting the selection, treatment, and preparation of kangaroo tendons which are intended to be used surgically. Having read in THE LANCET at different times the opinions of several operators who have used them, I wish now to refer more particularly to the defects pointed out in your reports of the meetings of the Royal Medical and Chirurgical Society and the Clinical Society of London, in THE LANCET, vol. i. 1885, p. 205, and in vol. ii. 1886, p. 819, in which the following remarks occur—viz., that tendon ligatures "had broken on more than one occasion"; that they "were sometimes slippery"; and that the knots tied in them "are apt to slip." My own experience is so directly opposite that I cannot help believing that the behaviour referred to was due to an imperfect condition of the tendons employed, and that they were either unsound or had been too much softened before being used. To ensure satisfactory results all tendons must be examined to ascertain that they are quite sound, because they are frequently injured by a hasty and careless removal from the animal's tail, a task requiring some little time and trouble, but not unfrequently performed by the rough hands of a bushman, who will, if not properly instructed, pull

them out by main force, and by this heroic method they are seldom obtained of the proper length and strength, for they either break off short or they are frayed at their edges, or split longitudinally, and I have seen many such which had been sold for surgical purposes. A moment's careful scrutiny of the tendon ligature by the surgeon himself is therefore necessary before trusting to it in the deligation of an important vessel. It should be about eighteen inches in length; if less than fourteen it had better be rejected at once; it should also be of the same calibre, and the same width from end to end. Its surface and edges should be even and without flaw, quite free from frayed, broken, or hanging fibres, and without any splitting or longitudinal separation of its substance. The soundness, or otherwise, is at once apparent by holding the tendon out at length between the eye and the light. Who, may I ask, would employ a silk or a gut ligature in which one or more of the strands were notched or broken? In consequence of its natural wavy and irregularly interlaced structure, a tendon is greatly weakened by a very short longitudinal split in its substance. I have always been careful to point out that a split or damaged tendon would easily break. *Apropos* to splitting, there is an article in Heath's "Surgical Dictionary" on Kangaroo Tendons, which seems to me to be rather misleading, for the writer states that "when fresh the tendons may be split up to any thickness required." *Cui bono?* But with seeming inconsistency, the same writer advises, a few lines further on, the rejection of those tendons which show any "longitudinal split." Another surgeon mentions a case, reported in one of the numbers of THE LANCET already referred to, in which he used a "slightly flattened" tendon. If this flattening was made by pressure, as one must presume it was, it is apparent that the process materially weakened the fibre of the ligature. As regards the slippery surface that has sometimes been spoken of in London, it also is probably caused by the preceding treatment to which the tendons have been subjected, and I fear that I may possibly be responsible, though unwittingly so, for some of this treatment. At all events, I desire to point out an error which occurs in a short paper of mine in the Medico-Chirurgical Transactions, in which it is stated (vol. lxx., p. 73) that tendons can be preserved "in 5 per cent. carbolic oil." This should have been 1 to 5 carbolic oil. I regret that this mistake escaped unnoticed by me till a few weeks ago, when some tendons that had been preserved in 5 per cent. carbolic oil were brought to me for examination by a friend, who had followed the directions in the paper just referred to. They were decidedly soft, rather slippery, and altogether different from other tendons which had been preserved in 1 to 5 carbolic oil. This may account for the "slippery nature of the material used," which is complained of in one of the reports. My recommendations, however, when first introducing tendon ligatures—as recorded in the *Australian Medical Journal*, Dec., 1877, and on all other occasions, with the one unfortunate exception—have been that they should be preserved in carbolic oil, precisely in the manner directed by Professor Lister in the preparation of catgut ligatures. And in this way all those I have ever used, or seen used, have been preserved. And there can be no doubt that they improve and become denser by keeping for many months or years in this solution; in fact, they become rather too hard for immediate use, without some previous softening. From recent experiments it appears that the best way of removing the hardness is by immersion for a few minutes (say at the commencement of the operation) in a warm solution of carbolic acid and water, 1 to 40, at a temperature of about 120°, which makes them pliable, yet not slippery. This is better than soaking for a long time in a cold watery solution; such a soaking for twenty-four hours is apt to make them sodden.

A final word about the length, if not trespassing too far on your valuable space. It is hardly possible to tie a tight and efficient knot on a vessel with a short ligature, especially when the operator's hands are wet; therefore it is all the more important to insist that the tendons should be over fourteen inches long, in order to afford a good firm hold, besides the fact that to obtain a tendon at its full length necessitates care in its removal from the tail.

I am, Sirs, yours obediently,

T. M. GIRDLESTONE, F.R.C.S. Eng.,
Surg. and Lect. on Clin. Surg. Melb. Hosp., and
Melbourne, June 27th, 1887. Lect. Surg. Melb. Univ.