The Boston Medical and Surgical Journal

TABLE OF CONTENTS

October 19, 1916

MASSACHUSETTS MEDICAL SOCIETY	
PAPERS READ BEFORE THE COMBINED MEETING OF THE SECTIONS OF MEDICINE AND SURGERY, JUNE 7, 1916.	
SYMPOSIUM ON GOITRE.	
SURGERY OF THE THYROID GLAND. <i>By C. A. Porter, M.D., Boston</i>	551
RECENT ADVANCES IN OUR KNOWLEDGE OF THE ACTIVE CONSTITUENT IN THE THYROID; ITS CHEMICAL NATURE AND FUNCTION. <i>By Edward C. Kendall, Ph.D., Rochester, Minn.</i>	557
CONDITIONS AFFECTING SECRETION OF THE THYROID GLAND. <i>By W. B. Cannon, M.D., Boston</i>	562
THE CLINICAL VALUE OF METABOLIC STUDIES OF THYROID CASES. <i>By Walter M. Boothby, M.D., Boston</i>	564
PARTIAL THYROIDECTOMY WITH LOCAL ANESTHESIA, SCOPOLAMINE AND MORPHIA. <i>By Frank H. Lahey, M.D., Boston</i>	566
THE TREATMENT OF GRAVES' DISEASE BY THE ROENTGEN RAY. <i>By Malcolm Seymour, M.D., Boston</i>	568
ORIGINAL ARTICLES	
THE EXCRETION OF HEXAMETHYLENAMIN BY DAMAGED KIDNEYS. <i>By George Gilbert Smith, M.D., Boston</i>	569
THE ILEO-CECAL VALVE AND THE CHRONIC INTESTINAL INVALID. <i>By John Bryant, M.D., Boston</i>	572

BOOK REVIEWS	
Pulmonary Tuberculosis. <i>By Maurice Fishberg, M.D.</i>	574
Medical Hints for the Use of Medical Officers Temporarily Employed with Troops. <i>By J. Edward Squire, M.D.</i>	574
The Basis of Symptoms, the Principles of Clinical Pathology. <i>By Dr. Rudolph Krehl</i>	574
EDITORIALS	
NEW ENGLAND SURGICAL SOCIETY	575
ELIMINATION OF THE RAT	576
COLORS THINKING	576
THE REGULATION OF VENEREAL DISEASE IN AUSTRALIA	577
A NEW FORM OF POST-GRADUATE TEACHING	578
MEDICAL NOTES	578
MASSACHUSETTS MEDICAL SOCIETY	
STATED MEETING OF THE COUNCIL, OCTOBER 4, 1916	581
MISCELLANY	
NOTICES, APPOINTMENTS, RECENT DEATHS, ETC.	586

Massachusetts Medical Society.

PAPERS READ BEFORE THE COMBINED MEETING OF THE SECTIONS OF MEDICINE AND SURGERY. JUNE 7, 1916.

SYMPOSIUM ON GOITRE.

I.

SURGERY OF THE THYROID GLAND.

By C. A. PORTER, M.D., BOSTON.

THE surgical lesions of the thyroid which we have to consider are infections; simple bilateral diffuse hyperplasia, or hypertrophy of the gland; fetal adenomata; cysts; adenomata and irregular colloid tumors, often in combination; malignant tumors, sarcomata and carcinomata; and finally, toxic goitres and true exophthalmic goitre or Graves' disease. In infections, we have to do with drainage operations; at times, removal of portions of the gland; in the condition of adenomata, cysts, and colloid goitre, removal of the cyst, enucleation of the tumor or, as in the case of colloid growths, a sufficient removal to relieve obstruction to the trachea, esophagus, etc., or for cosmetic purposes; next, removal of tumors which, owing to their appearance late in life, or rapid growth, suggest the possibility of incipient malignant disease; finally attempt at cure or palliation, *i.e.*, tracheotomy, of actual malignant disease.

Of the goitres with toxic symptoms (so-called hyperthyroidism) there are two types: those in which a goitre has been present for varying lengths of time, followed ultimately by toxic symptoms, with or without exophthalmos; and, perhaps the *true type* of primary Graves' dis-

case, in which general toxic symptoms not rarely precede or quickly follow the development of an enlarged thyroid.

1. In many infections there is undoubtedly a process in the thyroid gland which is shown by swelling, tenderness, increased size, fever, leucocytosis, pain (often referred to the back of the ear or neck), difficulty in breathing or swallowing. As a rule, under ice-bags such infections spontaneously subside without the formation of pus. Occasionally a definitely suppurative thyroiditis takes place with pus formation, requiring evacuation on ordinary surgical principles. I have seen a few cases which were diagnosed as ordinary deep cervical abscesses. In spite of the large venous supply of the gland, pyemia is rare, and the results of proper drainage excellent. Rarely, one sees, however, as a late result, acquired myxedema, which develops after destruction of the thyroid cells, which are replaced by scar tissue. Tuberculosis, actinomycosis and syphilis very rarely, but occasionally, affect this gland.

2. Simple bilateral diffuse hyperplasia, or hypertrophy, of the gland occurs very frequently in girls at or near the age of puberty, in many regions where goitre is endemic; not rarely after marriage, or during pregnancy. With very rare exceptions these hypertrophies should not be treated surgically, for they may disappear spontaneously, or through general medical treatment,—iodine, thyroid extract and x-rays. Inasmuch as they are practically without symptoms, and patients desire operations only for removal of the swelling in their neck, the surgeon must carefully consider whether the scar may not be more of a deformity than the original growth; and operations, if performed, must be done in such a manner that the lower

poles are left of symmetrical size in order that one side of the neck may not be smaller than the other.

3. Fetal adenomata occur as single or multiple, nodular, circumscribed tumors in one or both lobes, developing early in life, or at any period. They lead often to marked deformity, to pressure on the trachea or esophagus,—depending upon their location. If the thyroid enlargement takes place forward, between the sternomastoid muscles, pressure signs are often absent. If, however, a relatively small tumor grows behind the insertion of the sternomastoid muscle, pressure symptoms are apt to arise as the muscle holds back the tumor from any outward growth. With these fetal adenomata there occur varying degrees of toxic symptoms on the part of the heart, nervous system, etc., which are rarely, however, accompanied by exophthalmos.

Goetsch believes that these adenomata have the power of secreting an excess of thyroid juice, which explains the marked relief of general toxic symptoms which almost always follow their removal. Operation, though at times difficult, is most satisfactory, and consists either of enucleation of the nodules from the thyroid tissue, or occasionally it is simpler to do a lobectomy, provided enough thyroid may be left behind in the other lobe to carry on the function of the gland.

Cysts are not uncommon, varying in size from a marble to enormous tumors containing a couple of quarts. Into them, now and then, occur sudden hemorrhages, which may cause even alarming pressure symptoms, but apart from this, their harmful effect is purely mechanical. If a cyst is suspected no harm will come from aspiration with a hypodermic needle, for diagnostic purposes, followed, if deemed advisable, by the injection of carbolic acid, corrosive sublimate, or some irritant designed to prevent recurrence. It is important to make this diagnosis, for treatments by iodine, x-ray, thyroid extract, etc., are absolutely contraindicated and useless in cyst formation. If cysts are increasing in size, or causing obstruction, they should be removed with the technic which will be described in thyroid operations.

5. Colloid is the most common form of thyroid enlargement and varies in all degrees. As a rule, operation is desired for cosmetic purposes, or owing to the formation of large bilateral tumors in the neck, with a varying amount of pressure symptoms. It is surprising to see to what a huge size some of these tumors may develop without causing any symptoms whatever. In other cases thyroid enlargements take place posteriorly, encircling the trachea, or trachea and esophagus, and bring about severe obstructive symptoms. Iodine and thyroid extract, with or without x-ray, sometimes cause diminution of these tumors, but in general their effect is slight, and surgery fulfills here one of its safest indications. In such thyroid enlargements, occurring before puberty, operation

should be delayed as long as possible, lest too much of the more or less active gland be removed and interfere with the proper development of the individual.

In regard to the technic of operations for benign tumors of the thyroid, the first question for consideration is the choice of the anesthetic. As my experience increases, I am more and more in favor of local anesthesia, where the mental attitude of the patient is satisfactory. While a change from local to general anesthesia is always possible, it is time-consuming, increases danger of infection if the patient struggles; and much judgment should be used in endeavoring to select those patients who will satisfactorily put up with the idea of being conscious while their neck is operated upon, or occasionally bear a little pain or sense of dragging. I give sulphonal—grains 20 to 30—the afternoon previous to operation, and one or two doses of morphia, commencing an hour and a half before the appointed time. With the second dose, 1/120 of a grain of scopolamine, which may be repeated just before operation, if the patient is not sufficiently drowsy. This combination of sulphonal, morphia and scopolamine is most satisfactory when it works well, but occasionally there is excitement instead of quiet, a condition like coma vigil,—the patient sleeping until the surgeon starts to do something, then waking up and resisting. If a general anesthetic is to be administered in such instances, it must be given with great care, as the patient is often overdressed for a satisfactory ether operation. A pillow should be placed between the shoulders, so that the neck may be hyperextended, the table in moderate Fowler's position, facing a good light. No pains should be spared for a proper posing of the patient before operation, as great difficulty will be found in carrying out a proper technic if the neck slumps backward. There is no doubt that the collar incision of Kocher is the best for these operations. This may be placed across the bottom of the neck, where a necklace will later conceal the scar, or it may be made by a low incision over the sternum, when no scar will be visible unless a low-neck is worn. I use a 1 or 2% novocaine solution with 1-40,000 adrenalin chloride. It is best to mark out the line of incision by an intradermal injection of the solution, which should bring out a white wheal on the skin. The novocaine acts thus quite quickly from tension, more slowly from its anesthetic action. If you will recall your anatomy you will remember that all the nerves supplying the front and side of the lower neck emerge from the posterior border of the sternomastoid muscle, from its mid point downwards for an inch and a half; the superficial cervical branch, the inner, middle and outer branches of the supraclavicular nerves. While we are waiting for the novocaine in our line of incision to act, these nerves at this point should be anesthetized by deep massive infiltra-

tion. The skin and platysma muscle should next be incised, usually from external jugular to external jugular, the posterior incision extending somewhat upward, rather than transversely, in the lines of cleavage of the skin. The upper flap should be dissected to the level of the bottom of the thyroid cartilage, the lower down to the sternum. The enlarged anterior jugular veins next come into view, and must be carefully clamped and tied between ligatures. The sternomastoid muscles are then to be dissected free and retracted. If the tumor is a large one, there is no objection to dividing these muscles a half to two-thirds through; mattress sutures will ensure no deformity. The surgeon has next to consider the advisability of separation of the sterno-hyoid and thyroid muscles in the median line, with lateral retraction, or a transverse division of them at a higher level than the skin incision. There is little doubt that separation is preferable to division on account of less post-operative pain in swallowing, and freer movements when the wound is healed. Large or vascular tumors require free exposure for neat and bloodless work, and the muscles must then be divided. No time will be lost, and convenience assured, by immediate ligation of all skin vessels. Fine plain catgut should be used rather than silk, as in many instances chromic gut or silk ligatures are discharged from just under the skin some time after the wound has apparently healed. The intrinsic capsule of the thyroid varies much in thickness and in adherence to the gland. Any previous infection or treatment with boiling water, or x-ray, makes its removal more difficult. Anteriorly the capsule is of no importance, but if we are to operate upon both sides of the gland, it marks behind the curtain beyond which lie the recurrent nerves and the parathyroid bodies. When the muscles have been well retracted, careful inspection of both lobes is necessary to determine what operation shall be done,—enucleation of a cyst or an adenoma, removal of the anterior two-thirds or three-fourths of both lobes, or hemithyroidectomy on one side, and upper partial lobectomy on the other.

Lobectomy is comparatively simple and bloodless. The gland is grasped with gauze or a double hook, pulled toward the median line, and the upper and lower vessels clamped between forceps and divided. In spite of care there is often some troublesome bleeding near the middle of the gland, where the inferior thyroid breaks up into a number of branches. An injury to the nerve usually occurs from hurried grasping of a bleeding vessel through the capsule. Local anesthesia has the great advantage that it forces the surgeon to work gently and slowly, and any change of voice can be immediately noted. As a rule, these manipulations are painless, but occasionally the patient complains when a ligature is tied, or the tumor is dragged upon. Several times I have enucleated a cyst or tumor from

behind the sternum, and have been surprised at the freedom from pain. If supplementary injections of novocaine are necessary, it must be remembered that several minutes are required for complete anesthesia. The common error made is to begin without this delay. In large, irregular tumors, Balfour's suggestion is valuable, *i.e.*, the division of the isthmus, and separation from the trachea of the two halves of the glands from within outward. After the amount to be removed has been determined, every effort should be made to control the blood supply by deep mattress sutures or pressure by the finger on the vessels at the base of the lobe. When all bleeding has been stopped, the remains of the lobe are then sutured, leaving behind the lower rather than the upper poles. In all cases of tracheal obstruction, local anesthesia is invaluable. When once the trachea has been bared and is clearly visible to the eye, a general anesthetic may be safely administered, if necessary. In spite of all precautions, hemorrhage (sometimes serious) takes place in a definite proportion of cases. Most frequently a vein is torn in delivering the gland,—a danger much increased when patients are cyanotic from obstruction or vomiting under ether. In general, drainage can be omitted. If used, a small stab wound through the lower flap is preferable to its insertion through the line of incision. Whenever there is mild infection of a drain through the incision, there is apt to occur a simple adherent to the deeper tissues, which causes an unsightly deformity, and dragging on deglutition. The platysma must always be sutured with care; the skin incision closed either by a sub-cuticular catgut suture, or interrupted fine silk or horsehair stitch loosely tied and removed within three days. With this technic the scar of the collar incision is practically invisible. A moist dressing for twenty-four hours allows oozing from the drain or between the stitches. At the end of two days the drain should be removed, and in the mean time great care must be taken to prevent infection from looseness of the dressings. I use a single large gauze sponge, snugly held to the skin by adhesive plaster at the top. This is raised like a lid, and the dressing applied, which is held in place by a towel passed behind the neck, crossed and pinned to a swathe in front. For several days flexion of the head is painful, and patients are bothered a good deal by mucus and difficulty in swallowing. If no drainage is used, in spite of careful hemostasis, there occasionally occur collections of bloody serum which have to be evacuated, and not infrequently one notices little collections just under the skin of a sterile bloody serum, which has to be pricked like a blister. The above-outlined technic is applicable to all benign tumors and cysts of the gland which require removal. I have had several rather alarming hemorrhages during the operation, but as yet no case of sec-

ondary hemorrhage. Three times a recurrent laryngeal nerve has been injured, either in the dissection, or carelessly grasped in the attempt rapidly to control bleeding. Though some of the tumors and cysts have been very large—one filling up a third of the superior mediastinum, and another adherent to the sternum beyond its middle—I have had no death in more than 100 cases until last month, when an old lady of 65, who for 40 years had had a very large cystic tumor, died from hemorrhagic nephritis and bronchopneumonia.

There have been a few instances of continued growth of the tumor or formation of a new cyst, making necessary a second, and twice a third operation. The surgery of non-toxic goitre, then, though often difficult, is extremely satisfactory—the cosmetic results good. The exact opposite is true of malignant disease. In the last fifteen years there have been twenty operations at the Massachusetts General Hospital. Of these, 17 were cancer and 3 sarcoma. One case was 25 years old; one 29, one 32, one 33 all the others over 40. The oldest patient (63 years of age) had had an enlarged thyroid for twenty years, with rapid growth in the last two. There were five cases of very long duration of a goitre subsequently undergoing malignant degeneration,—one 35, one 25, one 20, one 12, one 7, another 4 years after onset. This shows quite clearly what my experience confirms,—that the development of an enlarged thyroid after fifty, or rapid increase in a long-standing goitre, is an indication for immediate operation. When once the diagnosis of cancer is clear, from the stony hardness of the tumor, paralysis of the recurrent laryngeal nerve, and enlargement of the adjacent lymphatics, a radical operation is practically hopeless. The only chance of cure rests in a thorough operation before the carcinoma has spread beyond the capsule of the gland. In one private case such an operation was done, with the removal of a portion of the esophagus. For a year there was complete comfort, but recurrence then took place. In two other patients over sixty, with rapid growth in a few months, recurrence followed operation too soon to make it worth while. Where obstruction to respiration makes tracheotomy necessary, local anesthesia is clearly indicated. This will be found to be a most bloody and trying operation, not rarely followed by secondary hemorrhages; pneumonia and bleeding commonly end the life of these wretched patients.

We have next to consider the toxic cases—so-called hyperthyroidism. As my paper deals only with the surgical aspects of Graves' disease, I shall not enter into the various theories as to its causation and complex relations or interrelations with the other glands. One thing remains clear—the etiology is not yet thoroughly understood. It is also, I think, impracticable to divide the toxic cases too minutely into groups. Broadly speaking, however, there are two

types: one, where the patient for months or years has had an enlarged thyroid, and subsequently developed the toxic symptoms; another group where, preceding any manifest thyroid enlargement, there is nervousness, tachycardia, palpitation, etc., or the general symptoms develop with an increasing enlargement of the gland. If any rough distinction can be made, the first class may be called tumors with hyperthyroidism; the second, true cases of exophthalmic goitre. It is in this latter class (2)—though, of course, there are exceptions—that we find the most severe toxemia, exophthalmos, and the greatest loss of weight. What, at present, are the indications for surgery in these two types of cases? What are the dangers of operation, and what has surgery to offer better than medical treatment? Quite recently the x-ray, being safe and favorably regarded, is being used more and more. The invention of the Coolidge tube, which allows measured, massive dosage, may prove a valuable aid in this disease. Personally, I have been in the habit of having one or two thorough treatments previous to operation. I have had little experience with long-continued treatment. While, undoubtedly, marked improvement has occurred, particularly in the milder cases, I have not been impressed with the results, and in one case in particular, I can see very little improvement after a year of treatment. Dr. Seymour, who has recently been following a series of cases, will later give us his impressions. It has one great advantage,—if not persisted in too long,—the element of safety and the absence of scar. There are some surgeons, notably Kocher of Berne, who think that surgery is the ideal treatment for thyrotoxicosis. He operates as early as possible after a very brief period of rest, and refuses operation on all advanced cases. His mortality is small (about 4%), because, in fact, he operates upon a selected group of cases. The general feeling at present is, I think, that all cases of Graves' disease should first be treated by rest and medical treatment. If satisfactory improvement occurs, this may be continued, but is not applicable to wage-earners and those who cannot give up the time. While under neutral bromide of quinine and other drugs, almost miraculous improvement takes place now and then, it is my experience that such cases are apt to recur, and that if lobectomy, single or double partial, has been performed, cures occur more rapidly and are more permanent. I would advise operation of some sort, then, in cases where improvement is unsatisfactory under medical means; in cases which have already lasted more than a year, with commencing myocardial weakness, exophthalmos, etc., or where it is unlikely that medical treatment will act sufficiently quickly to bring about a satisfactory result. Plummer of Rochester believes that after a year, operation is less hazardous than in the early stages. My experience has shown that

Graves' disease in girls from 18-20, particularly if the mental attitude is excitable, has seemed to carry a grave operative prognosis. Another type in which the risk is poor are women between 40 and 50, who are apparently passing through the menopause. The Hebrew race, owing to their neurotic constitution and lack of control, make a calm preparation for operation difficult, and I have had, among them, a comparatively large mortality. There is another group of rather chronic cases of years' duration, with ups and downs,—exophthalmos, rather firm vascular glands, tachycardia, often irregularity of the heart,—in which there is doubt as to whether the patient is actually suffering from Graves' disease or the aftermath of a passed toxemia. It is in this type that the various newer tests are of very great value. I mean estimation of the carbohydrate tolerance, basal metabolism, differential blood counts, injections of adrenalin, etc., in order that we may determine, if possible, whether the symptoms are due to an overactive gland. In these cases it is probable that other glands, the sympathetic system, and the myocardium have been affected; and it is often a nice question of judgment to decide for or against operation.

Before considering operation, there is one thing I wish to emphasize, and that is an increasing conviction that the psychic influences before operation are of extreme importance—in fact, I believe make for recovery or death in certain instances. The surgeon should have absolute control of the situation, and the complete confidence of the patient who, if possible, should look forward to the day of operation, unless we follow Crile in all detail in his "theft of the thyroid." In no other surgical disease do the surroundings, methodical nursing, cheerful mental attitude, etc., count so much; to no other patients do the occasional worries and effects of a large clinic work so great a detriment. If, as I believe, the mental attitude of the patient is so important, it is queer that surgeons have not come to more unanimity upon the choice of an anesthetic. Kocher has always insisted upon local anesthesia, and I have seen him adhere to this rule to, if not beyond, the limit. The Mayos, whose mortality is equally small, have the patient brought to the amphitheater and quietly etherized by the drop method. Crile, as we know, endeavors to carry out his anoci conception, planning to keep the actual date of the operation from the patient, and gives gas and oxygen in the patient's bed. As my experience increases, I am more and more in favor of the morphia-scopolamine sequence with local anesthesia, when the patient's mental attitude is favorable and the drugs act well. In a few operations I have had to abandon local anesthesia, owing to the complete lack of control of the patient, and have felt that this abortive attempt was much worse than if I had started with a general anesthetic. Indeed, I will go further

and say that in bad cases of Graves' disease, if anything goes wrong with the preparation for operation, and a patient arrives in a highly excited condition, it is always best to delay for a more favorable opportunity.

From the 185 cases in hospital and private practice which I have had, I have selected the 85 which, at time of operation, showed definite Graves' disease or hyperthyroidism present, or very definite in the immediate past. I have excluded the milder cases of transient hyperthyroidism, and considered them in the previous part of this paper. In these 85 cases I have performed 31 ligations, 19 hemithyroidectomies; in 17 the right lobe was removed with ligation of the vessels on the left; in 18 the major portions of both lobes, leaving a bit of the upper or lower poles, and a posterior strip of the gland. Double partial lobectomy is, in my opinion, a more severe and bloody operation than hemithyroidectomy; but as I look over my cases I feel that more immediate improvement takes place, and the danger of recurrence of symptoms is less. The most difficult question to determine when once operation has been decided upon, is between preliminary ligation of vessels, or removal of portions of the gland. Upon this question different surgeons are not in agreement. A majority, I believe, feel that in the severer cases a preliminary operation of ligation, usually under local anesthesia, makes a subsequent thyroidectomy less dangerous. Whoever has done a secondary operation on these cases is convinced of two things,—that ligation makes the operation somewhat harder, owing to adhesions, and secondly that the ligature of the superior vessels has not much diminished the vascularity of the gland. I think the benefit which frequently follows is due to division of the sympathetic nerves accompanying the vessels. In two cases only has ligation of the superior vessels been sufficient to bring about a cure; after it there has usually been either no perceptible improvement or a period of improvement followed by recurrence. It should be regarded distinctly as a preliminary operation and should, at varying periods, be followed by further operation. In a number of my cases, before the patients would consent, intervals of from one to three years have elapsed, at which time the element of safety after ligature is probably abolished. In such cases, therefore, we should consider, if they are severe, further ligature of vessels before thyroidectomy. In other words, after ligature of the superior vessels, there seems to be, according to my experience, after ten days or two weeks, a diminished danger in operation. After a long lapse between ligation and operation, the risk seems to be that of a primary thyroidectomy. I have analyzed, very carefully, my fatal cases, some of which date back to 1909, in order to learn, if possible, any avoidable mistakes. I have had five deaths from toxemia,—one in a suicidal, extremely toxic case, with pho-

tophobia and confinement in a dark room for three months. Ligation was simple, but in spite of all treatment, death with all the characteristic symptoms, occurred at the end of three days. This was a desperate risk taken deliberately. Of the four other cases the operation was a primary lobectomy; two were in young girls eighteen and twenty-five years; two middle-aged women undergoing the menopause,

The one aged eighteen deceived me in her degree of toxicity. Right hemithyroidectomy with ligation of the left vessels was performed under ether, in 1910, and was followed by a typical toxic death in three days. The one aged twenty-five was a very toxic case; had been sick for eight years, and did not improve under three weeks in bed with rest treatment; in fact, on the whole, seemed worse. Right hemithyroidectomy, under gas and oxygen with ligation of the vessels on the left was performed; operation comparatively easy; died on the third day of typical toxemia.

A girl of thirty had extreme symptoms for two years; was unable to work for a year; fell exhausted on the street several times; treated in medical ward without improvement; heart irregular; marked systolic murmur with some enlargement. Right hemithyroidectomy under gas and oxygen, and partial left thyroidectomy were performed with extreme reaction, which persisted unusually long, leaving marked condition of prostration; patient died nine days after operation. All of these three were early cases, and at present I believe that a preliminary ligation should have been done, or the patients should have had x-ray treatment previous to operation.

A Jewish woman of forty, undergoing the menopause, suicidal for a year, refused operation four successive mornings. Finally she consented to it if I would perform it in the afternoon. Rather against my judgment, at the earnest solicitation of her husband, I agreed to do it, under gas and oxygen, removing the right lobe and half of the left. There was considerable bleeding, and with excessively rapid pulse the patient died in three hours. This was in 1909, and did much to influence me against operation. I have broken this rule since then once or twice, always to my regret, and with a severe operative, though not fatal, toxemia; until in March of this year I had another death, the first since 1912. A Jewess of 47 had been ill for five years with nervousness, headache, irregular menstruation, edema and dyspnea, pain in the upper abdomen, rapid pulse, with some enlargement of the heart, some glycosuria, unable to work. Under careful medical treatment for two and a half weeks, she lost seven pounds, and tachycardia and nervousness persisted. I advised ligation under local anesthesia as a preliminary. She had her morphia in two doses and was apparently drowsy, but before operation could be begun she jumped up from the table and insisted upon going home. Three days

later she consented, provided her daughter was present, to have operation under gas and oxygen. I performed a double partial lobectomy without difficulty, but from the time of her recovery from the anesthesia, extremely rapid pulse with excessive nervousness began, and increased until death, at the end of forty-eight hours. This illustrates strikingly, I think, a bad risk in a woman who was steadily running downhill. Operation undertaken was the only hope, and owing to lack of control, toxemia started before operation, and was only increased by it—I think it an error of judgment in attempting to benefit a patient unwilling to accept one's best advice. In four cases severe bronchitis, pneumonia or bronchopneumonia, for reasons which I cannot explain, ended fatally in from a week to a month. In the latter long case the ultimate cause of death was heat prostration in the severe July of 1911. A fifth case died, so far as could be made out, from a terminal pneumonia, aided probably by unsuspected morphinism. Why pulmonary complications should have occurred in these cases and been absent in the non-toxic goitres, it is impossible for me to explain. Another death, similar to a case of Dr. Judd's in Rochester, I cannot understand. Autopsy showed purulent pericarditis, enlarged thymus and adrenals, as a cause of death thirty-six hours after a simple ligation with a clean wound. In another, a large infarct of the right lung, after a very satisfactory gas and oxygen thyroidectomy, caused death on the third day. In this series of 185 cases, there was one instance of tetany—the first post-operative case at the Massachusetts General Hospital. A young girl of eighteen, with severe Graves' disease, had double ligation under ether in 1911, followed by marked improvement which lasted for almost two years—then recurrence of symptoms. Under gas and oxygen in October, 1914, right hemithyroidectomy and partial left lobectomy were performed. The operation was apparently intracapsular. On the fourth day characteristic tetany developed, which was controlled with calcium lactate. In spite of parathyroid tablets and continued calcium lactate, severe convulsions occurred, followed in January, 1915, by acute mania and death. It seems to me necessary to believe that in this instance the blood supply of the upper parathyroids was cut off by the ligation, or the unfortunate patient had an abnormally small number, for the operation was performed in such a manner that some of the parathyroids under normal circumstances must have been conserved. Thus this analysis seems to show that the danger of toxemia is diminished by preliminary ligation a short time previous to thyroidectomy; that perfect control of these patients is necessary, lacking which operation should not be performed. There has been no death from pulmonary complications following local anesthesia. The case of pul-

monary infarct and purulent pericarditis seems to me to be impossible to avoid.

In endeavoring to analyze the good results of operation, it is most difficult to make satisfactory statistics. Those who have seen the very marked relief within a few days which follows lobectomy, are absolutely convinced that the thyroid *has to do* with Graves' disease. On the other hand, there are occasional cases which, in spite of ligature and lobectomy, disappoint us with their lack of improvement. Some surgeons believe that the thymus gland is in part responsible for the symptoms, and state that improvement has finally taken place after thymectomy. With this operation I have had no personal experience. I am inclined to believe, however, that the failure distinctly to improve or cure a majority of these cases is due to delay in operating until permanent degenerations have taken place, or some not understood action of the sympathetic nerves or other glands in the body.

I have operated upon some quite desperate cases with marked improvement. One, a young girl, with congenital heart murmurs who, it was decided, could not survive the cardiac lesion plus the toxemia of Graves' disease, has been practically cured, as far as any toxemia is concerned, by preliminary ligation, followed later by thyroidectomy. Another woman who was bedridden with coincident mitral insufficiency and severe Graves' disease, is now entirely cured of all her Graves' symptoms, except slight exophthalmos. Another middle-aged woman, bedridden, with marked edema and rapid, irregular pulse, has, after operation, improved so that she can do her work, though some tachycardia and irregularity of the heart persist. Another woman of fifty-five, confined in an asylum, much to my surprise became almost normal after operation; was able to look after herself and go to the theatre alone; dying suddenly of pneumonia a year afterwards. In my experience, the cases which have had exophthalmos for any length of time never again have normal eyes, and not a few of the cases apparently cured under normal circumstances show, under excessive fatigue or worry, slight tachycardia and tremor. Of the 85 cases, 20 are absolutely cured, 18 much improved, 13 improved, 4 not improved and require further operation; 16 have not reported, or done so in too general terms to be valuable for statistics. If then, after a patient has had a reasonable course of rest, medical treatment and x-ray treatment (if this proves to be really and permanently satisfactory), the symptoms persist, operation should be advised. Among wage-earners, who cannot give up the time, and in the chronic cases, in which the disease has advanced too far to allow of delay, after preliminary rest, operation should be performed. Whether this be a preliminary ligation, or lobectomy under local or general anesthesia, must be determined in each

case. Finally, in chronic cases in which there is doubt as to whether the patient is suffering from existing toxemia or has been poisoned irreparably by the disease, modern tests, of which basal metabolism is probably the most important, will aid the surgeon much in deciding for or against operation.

II.

RECENT ADVANCES IN OUR KNOWLEDGE OF THE ACTIVE CONSTITUENT IN THE THYROID; ITS CHEMICAL NATURE AND FUNCTION.

BY EDWARD C. KENDALL, PH.D.,

The Mayo Foundation, Rochester, Minn.

ABOUT eighteen months ago I reported the isolation from the thyroid of a crystalline compound containing 60% of iodine. Since that time I have perfected the method for its isolation, and to some extent have studied its chemical properties and functions. The work may be briefly summarized as follows:

The compound was first isolated in crystalline form in 1914. The work just prior to the isolation had been accomplished with such smoothness and success that no difficulty was anticipated in repeating it, but despite most persistent efforts, no more crystals were isolated until August, 1915. Again it seemed that the isolation had been perfected, but try as we would, no more crystals were seen until February, 1916. This failure occurred in spite of the fact that

FIG. 1.—Crystals of the alpha iodine compound, which is the active constituent of the thyroid.