IX.—*A Chemical Study of Vegetable Albinism.*

By A. H. CHURCH.

THE phenomenon of vegetable albinism or albication here considered, must not be confounded with *chlorosis*, the pallor of imperfect chlorophyllation, and the usual accompaniment of etiolation; nor is it identical with *icterus*, a defect of coloration due to absence of iron, and promptly remedied by the administration of traces of a ferric salt.

In the autumn of 1877 I began some experiments upon the white leaves and white parts of leaves of certain variegated plants. It might be thought that there was nothing easier than to obtain abundant supplies of these materials for analysis; but such is not the case. Very few plants can be employed for this study: for either the quantities of material obtainable in an isolated state are inadequate, or it is found to be impracticable to secure, from the same plants, both white and green portions of precisely the same age, and grown under the same conditions. The common ribbon grass, *Phalaris arundinacea*, affords an example of the first difficulty; several kinds of variegated *Pelargonium* illustrate the second. My earlier set of experiments was made upon three species of plants, all admirably adapted for the purpose, for in each the white and green parts were at once abundant and strictly comparable. The three species laid under contribution last year were—

Acer Negundo Gathered September 17th
Ilex Aquifolium „ „ 24th
Hedera Helix „ October 4th

In every instance the leaves were gathered without the intervention of knife or scissors, and were at once transferred to well-closed bottles previously weighed; thus no loss of water could occur during transit to the laboratory.

The water, the organic or combustible matter, and the ash were first of all determined; the results, translated into percentages, are here given:—

	<i>Acer Negundo.</i>		<i>Ilex Aquifolium.</i>		<i>Hedera Helix.</i>	
	White.	Green.	White.	Green.	White.	Green.
Water.....	82·83	72·70	74·14	62·83	78·88	66·13
Organic matter.....	15·15	24·22	23·66	35·41	18·74	31·63
Ash.....	2·02	3·08	2·20	2·47	2·38	2·24

Thus the white leaves of these three plants showed an average of 10 per cent. more water than the green leaves, while the dry organic matter in the former was not quite two-thirds of that present in the latter. The ash from the dry matter was more abundant in the white than in the green leaves.

	Percentages of ash.	
	White.	Green.
<i>Acer Negundo</i>	11·76	11·28
<i>Ilex Aquifolium</i>	8·51	6·52
<i>Hedera Helix</i>	11·27	6·61

The nitrogen and the matters soluble in ether were determined in the dry matter of holly and ivy with the following results:—

	Per cent. nitrogen.		Per cent. ether extract.	
	White.	Green.	White.	Green.
<i>Ilex</i>	2·28	·85	6·77	6·83
<i>Hedera</i>	2·76	2·35	6·12	9·65

These figures may best be studied in connection with the more complete data derived from the second series of experiments.

It is in the ash analysis that the most remarkable differences in composition between the white and green leaves were noticed. More potash and more phosphorus pentoxide were invariably found in the white parts than in the green, while they were almost always poorer in lime:—

	Percentage composition of Ash of—					
	Maple.		Holly.		Ivy.	
	White.	Green.	White.	Green.	White.	Green.
K ₂ O	45·05	12·61	35·30	16·22	47·20	17·91
CaO	10·89	39·93	21·50	34·43	12·92	48·55
MgO	3·95	4·75	3·23	2·43	1·11	1·04
Fe ₂ O ₃	—	—	3·11	3·11	2·62	2·31
P ₂ O ₅	14·57	8·80	9·51	7·29	10·68	3·87

There is here no indication that the presence of chlorophyll in the green parts involves a higher percentage of iron.

We may now proceed to the description of the second series of experiments performed in the summer and autumn of the present year. Thanks to the permission of Sir J. W. Hooker, and to the obliging attention of the officials at Kew, I was enabled to make successful use of the Jodrell Physiological Laboratory in the Royal Gardens. My

experiments were further aided by a grant from the Government Research Fund.

Four exotic plants were the subjects of my renewed inquiry. The specimens of variegated plants submitted to analysis were gathered with every precaution :—

<i>Elæagnus pungens</i>	On July 17th.
<i>Plectogyne variegata</i>	„ 9th.
<i>Cyperus alternifolius</i>	„ 10th.
<i>Alocasia macrorrhiza</i>	In October and November.

Of the *Elæagnus* several sprays of wholly white and of wholly green foliage were used : the white and green parts were separated by rending from five leaves of *Plectogyne* : part of the *Cyperus* specimens were entire white or green leaves, part similarly coloured portions of leaves.

I give the results of my experiments on these leaves in the same order as that previously followed in recounting the results of the 1877 series—

	<i>Elæagnus pun- gens.</i>		<i>Plectogyne variegata.</i>		<i>Cyperus alterni- folius.</i>	
	White.	Green.	White.	Green.	White.	Green.
Water	69·49	63·24	74·60	69·72	77·35	71·72
Organic matter	28·98	34·65	22·25	27·42	20·02	25·53
Ash	1·53	2·11	3·15	2·86	2·63	2·75

It will be seen that the white leaves and parts of leaves of all three plants contain on an average 5·58 per cent. more water than the green leaves and parts of leaves. The percentages of ash in the dry substance of the leaves of each of the three plants were as follows :—

	White.	Green.
<i>Elæagnus pungens</i>	5·01	5·74
<i>Plectogyne variegata</i>	12·39	9·44
<i>Cyperus alternifolius</i>	11·63	9·76

The dry leaves of these plants were not obtainable in sufficient quantities for an exhaustive analysis of each, but a more complete study of the *Elæagnus* than of the others was made. Still the nitrogen in each specimen was determined, the results being here given :—

	Nitrogen in dry substance.	
	White.	Green.
<i>Elæagnus pungens</i>	4·24	2·82
<i>Plectogyne variegata</i>	2·23	2·39
<i>Cyperus alternifolius</i>	1·94	2·04

Something more will have to be said presently about the nitrogen in *Elæagnus*, but as a whole the preceding figures do not indicate any definite divergence between white and green leaves, so far as their percentages of nitrogen are concerned.

The ashes of the three plants were analysed with care, although the quantities of materials at my disposal did not admit of a complete study of each specimen.

The following table gives the percentage composition of the ash of —

	<i>Elæagnus pun- gens.</i>		<i>Plectogyne variegata.</i>		<i>Cyperus alterni- folius.</i>	
	White.	Green.	White.	Green.	White.	Green.
Potash.....	60·93	31·31	39·54	34·46	48·54	24·20
Lime.....	6·35	31·50	19·95	24·00	6·05	10·81
Magnesia.....	?	?	2·35	2·94	?	2·89
Ferric oxide.....	2·23	·77	?	?	?	?
Phosphorus pentoxide ..	14·02	10·20	7·62	4·59	7·09	7·34
Silica.....	4·56	2·25	·35	?	36·75	46·03

On comparing these figures with those obtained in 1877 in the analysis of the ash of white and green leaves of maple, holly, and ivy, observe the following three features in common:—

1. The ash of the white leaves contains much more potash than that of the green leaves.
2. The ash of the white leaves contains much less lime than that of the green leaves.
3. No deficiency of iron in the white leaves has yet been detected.

Further evidence, corroborating these and previous statements as to the main chemical differences between white and green leaves, is afforded by a second series of analyses just made of white and green maple leaves, and by an analysis of a wholly white leaf of *Alocasia macrorrhiza*, and also of a wholly green leaf from the same species, though not from the same individual plant, both the latter specimens having been grown in the water-lily house at Kew. I give partial analyses of the dry matter of these two gigantic Aroid leaves, with more complete examinations of their ash and of the ash of the maple leaves gathered in 1878:—

In dry matter.	<i>Alocasia macrorrhiza.</i>	
	White.	Green.
Total nitrogen	3·45	2·98
Nitrogen in albuminoids	1·16	2·10
Nitrogen, as nitrates, &c.	2·29	0·88
Albuminoids	7·35	13·38
Ash.	23·89	15·01

In 100 parts of ash of—

	<i>Acer Negundo</i> (1878).		<i>Alocasia</i> <i>macrorrhiza.</i>	
	White.	Green.	White.	Green.
Potash	54·44	13·03	60·48	38·07
Lime.	8·50	40·16	19·37	20·37
Magnesia	3·22	—	—	2·77
Ferric oxide	—	—	—	—
Phosphorus pentoxide	14·87	7·38	13·08	6·89
Silica	·68	1·31	—	—

The most striking results here relate to the nitrogen in the white *Alocasia* leaf, nearly 70 per cent. of this element existing in non-albuminoid and but partially elaborated forms, whereas in the green leaf less than 30 per cent. is in a similar condition.

I will now describe some additional experiments made with *Elæagnus pungens*, the only plant besides the maple of which I was able to obtain a supply at once fairly abundant, and yet entirely fulfilling the necessary conditions of comparison.

From physiological considerations, I was led to inquire more particularly into these three points:—

1. Amount of matters soluble in ether in the two kinds of leaves.
2. Proportion of albuminoid to total nitrogen.
3. Presence of calcium oxalate and carbonate. I throw the results of this further inquiry into a tabular form:—

Percentages of	In dry matter of leaves of <i>Elæagnus pungens.</i>	
	White.	Green.
Matters extracted by ether	2·63	6·05
Albuminoid nitrogen	1·83	1·81
Non-albuminoid nitrogen.	2·41	1·01
Calcium oxalate	·41	2·10
Calcium carbonate	traces	decided traces

(1.) It was natural to expect that the absence of chlorophyll from the white leaves would lower their per cent. of matters soluble in ether; in fact, they yield to this solvent less than half the amount that is given by the green leaves. This result was obtained only when pure

ether, free from both alcohol and water, was employed in the extraction-apparatus. When ordinary ether, neither dry nor free from alcohol, was used, the percentages of dissolved matters were 4.62 from the dry green leaves, and 3.28 from the dry white. But it was clear that some chlorophyll still remained in the green leaves, while some substances, not soluble in pure ether, had been extracted from the white. Unless we could assume that the amounts of resin, wax, oil, &c., are the same in the dry matter of both sorts of leaves, it would be hazardous to assert that the difference between the two ether extracts ($6.05 - 2.63 = 3.42$) represents the percentage of chlorophyll in green *Elæagnus* leaves when dry.

(2.) In determining albuminoid nitrogen in white and green leaves, I used the method, which I originated some five years ago, of curdling the proteid matter by means of carbolic acid solution. In some plants leaves dried in the water-oven give precisely the same percentages of albuminoid nitrogen in the residue when washed with water, as the fresh leaves of the same plant furnish when the carbolic process is adopted. In such cases the soluble albumin of the leaf has been rendered insoluble in water by the heat of the water, just as effectually as by the carbolic solution aided by metaphosphoric acid. The following determinations of nitrogen are in close agreement with those in the *Alocasia* leaves:—

	Percentage in dry matter of <i>Elæagnus</i> leaves.	
	White.	Green.
Total nitrogen	4.23	2.82
Nitrogen by carbolic process	1.83	1.81
Albuminoid nitrogen in residue insoluble in water.....	1.82	1.83
Non-albuminoid nitrogen	2.40	1.01

Thus, nearly 60 per cent. of the nitrogen in the white leaves is non-albuminoid, while just over 30 per cent. is the proportion in the green leaves.

(3.) The third special question to which I have been endeavouring to obtain an answer is that concerning the form or forms of combination in which the lime is present in the two sorts of leaves. With one doubtful exception, lime is decidedly less abundant in the white leaves than in the green. Now I found, in the case of *Elæagnus*, that the percentages of lime salts soluble in water, cold and hot, were practically equal in the dry matter of the white and of the green leaves, but that both carbonate and oxalate were present in notable quantities in the latter, and almost absent from the former. Analysis gave:—

	Percentages in dry matter of Elaëagnus leaves.	
	White.	Green.
Lime in soluble salts	·28	·31
Lime as oxalate	·18	·92
Lime as carbonate	trace	·50

From these numbers it is clear that the great excess of lime found in the green leaves, when compared with the white, is mainly due to the physiological process, of which calcium oxalate is the final product, although a part of that excess must be attributed to the simple deposition of calcium carbonate drawn up by the roots in solution in the soil-water, and left behind on the transpiration of moisture from the leaf. From the white leaves, albeit they contain more water, the transpiration is probably less, since, through the absence of chlorophyll, they can hardly be the seat of the production of carbon-compounds, and of the various changes attendant upon that process.

Before summing up the chief results of the present inquiry and endeavouring to correlate them with the phenomena of vegetable albinism, it will be useful to cite an analysis of two other plants, namely, of a parasite and its host. I chose for this purpose the common dodder (*Cuscuta Trifolii*) and the red clover, on which it was living. Like the white parts of variegated leaves, the dodder is destitute of chlorophyll. Moreover, by the aid of its haustoria, it brings its several tissues into intimate connection with the corresponding structures of the supporting plant, and so draws its nourishment almost wholly from the latter. The samples of the two plants, both in flower, were gathered simultaneously on the 15th September, 1878, care being taken to collect representative specimens. The analyses gave :—

	In 100 parts of—	
	<i>Cuscuta Trifolii</i> .	<i>Trifolium pratense</i> .
Water	87·39	81·10
Organic matter	11·45	17·59
Ash	1·16	1·31

	In 100 parts of the ash of—	
	<i>Cuscuta Trifolii</i> .	<i>Trifolium pratense</i> .
Potash.....	48·00	26·43
Lime	3·47	38·40
Magnesia	2·88	13·07
Phosphorus pentoxide	9·59	9·61

It will be observed that in the more aqueous character of the parasitic plant, and in its greater richness of potash, as well as in its low percentage of lime, we have points of agreement with albino foliage, while the vegetable host may be compared with green leaves.

40 CHURCH'S CHEMICAL STUDY OF VEGETABLE ALBINISM.

If we further compare with white leaves those parts or organs of plants which live at the expense of other parts, we shall find close analogies between them. In short, if we determine the water, the organic substance, and the constituents of the ash in each of the following cases, we shall be able to affirm generally that white leaves are related to green pretty much as immature leaves to mature, tubers to foliage, petals to green bracts, vegetable parasites to their hosts. The white leaf may in fact be said to be parasitic on the green: it may be a warehouse, but it is not a factory. There seem to be present therein all the materials by the aid of which the organic compounds of the leaf should be constructed, but the constructive faculty or impetus is lacking. Lime compounds are, it is true, found, but sparingly, in most white leaves, but then the lime compounds in mature foliage are in great part a sign and relic of past activities. Perhaps it may be found that some minor ingredient other than those to which I have referred in the present paper, may be deficient or wanting in the white leaves. Further analyses are requisite to decide this point. But, after all, some structural or histological defect may be the real cause of the want of constructive power in the white leaves—of their lack of chlorophyll. Comparative microscopic examination and minute measurements of white and green portions have been made, but no real insight into the matter has thus been obtained. Often, as in the maple, the white portion of a leaf has but two-thirds of the thickness of the green; often a fluted or constricted aspect is caused by the imperfect development of the white tissues, as in the white borders of some *Pelargonium* leaves and the white stripes of some grasses. It has been further observed that the variegated and albino plants and shoots are usually smaller, more susceptible of injury by frost, and generally weaker than the normal green foliage. And when a variegated plant does throw out a wholly green shoot it grows with greater vigour than the rest of the plants. It has long been known that an albino cutting cannot be struck. Endeavours to stimulate the development of albino foliage in variegated plants, and to cause such plants to revert to the normal forms by chemical treatment of the soil, have not yet been attended with decisive results. Yet no decided structural defects accompany these slight peculiarities. But I hope another season to study this subject with further appliances and by new methods. I hope to measure the exact relations of equal areas of white and green foliage to carbon dioxide, oxygen, and moisture, and shall endeavour to see what results the injection into the vessels of white foliage of solutions of various organic and inorganic bodies may achieve. This summer I tried a few experiments of this kind with a measure of success, using the needle and syringe of hypodermic injections, and instantly closing the puncture with a drop of collodion. Thus several cubic centimeters

of the desired solutions were introduced beneath the epidermis of some *Echeveria* leaves, forming beneath the epidermis a gigantic blister, the contents of which were absorbed in a few minutes on exposing the plant to sunshine. Various colourless compounds becoming coloured by oxidation or reduction were employed; so also was lithium citrate, the course of which could easily be followed by subsequent dissection, drying, and spectroscopic examination of the various parts thus prepared. This use of lithium citrate, which I had the pleasure of introducing to Dr. McNab's notice some years ago, has yielded some valuable results in his hands and in those of certain foreign observers. I now find it quite possible to employ it on living plants without injuring their development. It is evident that it may enable us to learn something of the relations between the various parts of plants, and may aid in solving the question of vegetable albinism. But for all this physical work I must await the sunshine of another year.

A word about the ash analyses in the present paper. In a good many instances the figures are derived from a single determination, often necessarily made with a very small quantity of material. If a formula were to be established, such analyses, though performed in many cases with microscopic crucibles and other vessels, and with the aid of an exquisite assay balance, would be inadmissible. But they at least admit of certain general deductions being drawn, for they point without an exception in the same direction.

The nitrogen and ash analyses of the second series have nearly all been made by my assistant, Mr. C. Rawson.