scious. The temperature kept still rising, so that by 12 o'clock midnight it was 107.6° . He was now (Sunday 11th) cold-sponged every half hour, being covered with a cold damp sheet in the intervals. The temperature soon fell to 104.4° and remained at that during the early morning hours, and between 6 and 7 o'clock a.m. he again became conscious but the temperature was again 104° , the pulse very rapid, uncountable and weak. He soon again became unconscious and died 12.20 p.m.

Post-mortem examination confirmed the conditions made out during life. The lower six inches of the ileum showed large and congested Peyer's patches and solitary follicles, those near the cæcum having commencing superficial ulceration.

The case is of special interest in several points :—

(1) The acute onset with continuous high temperature— 103° - 104° —from the evening of admission, which was apparently the sixth day of disease, till it terminated; with a fatal exacerbation on the tenth day to 107.6° F.

(2) The diagnosis of enteric positively supported by the atropine test without delay and with great ease.

(3) The appearance of "delirium ferox," which appears to be a very fatal symptom in this disease.

(4) The confirmation of the diagnosis by *post-mortem* examination, where the intestinal lesions were seen in an unusually early stage.

ART. XVII.—*The Place of Curare in the Treatment of Tetanus.* By JOHN S. MCARDLE, M.Ch. (*Hon. Causâ*), R.U.I., F.R.U.I., F.R.C.S.I.; Professor of Surgery, National University of Ireland; Senior Surgeon, St. Vincent's Hospital; Consulting Surgeon, National Hospital, Holles Street, and Children's Hospital, Temple Street, Dublin.

I WAS driven to the study of tetanus by the loss of one very dear to me and for whom in the then state of our knowledge I could do absolutely nothing. That chloral, morphine,

belladonna, and other drugs of the kind gave relief no one doubts, but that they were in any way curative I never believed, and when face to face with acute traumatic tetanus they failed utterly.

The case now decided in the English Criminal Courts tempts me to address you on this subject. I notice that Sir F. E. Smith said in opening his case that "few people knew anything of the lethal substance curare." Now in 1884 I experimented with the crude wourali after reading the account of the effects of this poison when employed by the South American Indians. The first strange thing I learned from the writings of the traveller who wrote that account was that the flesh of animals killed by wourali may be eaten with impunity; second, that the poison itself produces no effect whatever when taken internally.

In the June number of the DUBLIN JOURNAL OF MEDICAL SCIENCE, 1885, there is a note of a case of mine which reads as follows:—

"Mr. McArdle read the notes of a case of acute traumatic tetanus, in which two-third grain doses of urari every fifth hour resulted in a cure, the more remarkable effects produced by the above named doses being relaxation of the contracted muscles in from six to ten minutes after administration, very rapid and tumultuous action of the heart, cyanosis, laboured breathing and dilatation of the pupils. Once the patient was sufficiently under the influence of urari, the evacuations from the bowels were regular. Mr. McArdle suggested the combination of urari and pilocarpine, in the hope that the cardiac and respiratory trouble produced by the former might be prevented by the latter.

"He also showed that urari, to be of service, must be used in large doses, and that the drug is cumulative."

First wourali engaged my attention, but there are many impurities in this crude drug, and many of the animals I tried it on (chiefly horses) succumbed as much to the medicinal remedy as to the tetanic virus. The first human

subject on whom I tried the crude drug escaped only after very vigorous treatment. I was only guessing at the dosage, and so gave this patient as much of the drug as would represent half a grain of the alkaloid "curare," which we now use.

Within a half minute of the injection the patient collapsed completely. He became deeply cyanosed with irregular breathing, pulse small and almost uncountable, cold clammy skin. The pupils, which had been contracted, became momentarily widely dilated, the whole aspect of the patient being appalling.

Artificial respiration, elevation of the end of the bed, enema of ammonia, and hypodermic injection of nitrate of pilocarpine turned defeat into victory. In this case impure catgut was the cause of the trouble, as several minor cases occurred before we discovered the origin of the most distressing disease with which we are surgically acquainted, save and except malignant œdema.

This patient made a quick recovery, and I had the pleasure of dining with him when last in America, where he is a prosperous banker.

Later on urari was isolated, and at the time we thought the *ultima Thule* had been reached in regard to this substance, which had now become a well established drug. This substance is the arrow poison of the Macusi Indians of the Amazon Valley.

The dosage of this drug is well established, one-tenth to one-sixth of a grain being perfectly safe and often effectual for the relief of the spasm of tetanus.

Before the serum treatment of tetanus was thought of very many cases even of the acute form recovered under the influence of the derivatives of wourali. The specific action of these drugs is to temporarily destroy the function of the motor end plates and thus put the voluntary muscles out of action. Under their influence the unstriped muscles function quite normally, and so the patient's strength can be

maintained while the tissues destroy the tetanus toxin already in the system.

The following case will illustrate the rôle of these so-called paralyzants in treatment of the dread disease, which usually shows a mortality of 80 to 90 per cent. even under favourable circumstances :—

J.M., aged 38, was admitted to St. Vincent's Hospital on June 20th, 1910, with a compound fracture dislocation at the ankle. The tibia had gone through his boot and entered the soil on the road, many particles of which became embedded in the periosteum and raw bone surface. When I saw him his jaws were already clenched, his abdomen hard as a board, and he had well-marked arching of the spine. He had had some attacks of laryngeal spasm which threatened his life. Fortunately I had in my case a tube of "curare," a drug nearer to perfection than urari. Within three minutes of a hypodermic injection of one-twelfth grain the spasms ceased. I injected into the sciatic the contents of one ampoula equal to 9 c.c. of antitetanic serum. After six hours spasms began to reappear, and so a further one-twelfth grain of curare was administered, and so for three days, every sixth hour the dose was repeated.

Owing to septic conditions I was obliged to amputate the leg at the seat of election, but to-day this man walks about as if nothing had happened to him. The Rowley leg which is now being supplied to the French and English well-to-do deprives amputation of most of its terrors, and any surgeon who does not know fully about this device should in justice to himself and his patients make himself acquainted with its perfections.

The reading of the case above referred to is very simple. If perchance we had not been in possession of a drug to control the laryngeal spasm this patient would have died. Anti-toxin would not have had time to act. By keeping the patient alive, so as to let the serum do its duty, the toxins were neutralised and recovery made possible.

It is important to note the difference between curare and curarine. The latter alkaloid is more certain in its effects, and its dosage is more certain. Its chemical composition is $C_{18}H_{35}N$.

The dose of curare is $\frac{1}{32}$ to $\frac{1}{6}$ grain.

The dose of curarine is $\frac{1}{200}$ to $\frac{1}{40}$ grain.

Their action is similar. The following notes kindly furnished me by Mr. M. J. O'Connor gives an account of the physiological properties of the derivatives of wourali :

“The characteristic action of curare—general paralysis—is due to the alkaloid ‘curarine.’ Curarine is not crystalline, and it is not known if it is quite pure. It was prepared by Boehm by extracting the crude ‘curare’ with various solvents until he obtained a material which gave a maximal poisonousness. This curarine in doses of 0.0002 gram per kilo intravenously causes complete paralysis of a rabbit. This would mean that if a man has the same sensitiveness as a rabbit a dose of one-sixth grain (intravenously) would be fatal or nearly so. The dose of ordinary curare cannot be given, especially as it tends to deteriorate with time. It would be much greater than one-sixth grain.

“When given by the mouth curare is usually ineffective owing to slow absorption and rapid excretion through the kidneys. This excretion is so rapid that if artificial respiration be given a rabbit recovers from ten times the fatal dose in 30-90 minutes.

“Curare produces its action by preventing the nerve impulse from passing from the nerve to the muscle—paralysing the neuromuscular junction. The muscles are not all affected at once. The respiratory muscles are the last to be affected. But as the effect of the drug passes off rapidly—owing to excretion of the curarine by the urine—it would be difficult to keep an individual under the correct dosage, though perhaps not more so than to maintain anæsthesia by intravenous transfusion of an narcotic (like hedonal

narcosis). It can to some extent be antagonised by eserine. *i.e.*, an injection of eserine may cause the reappearance of spontaneous respiratory movements.

“Owing to its power of putting a block on the junction of nerve and muscle, and thereby preventing convulsions arising from the central nervous system producing any effect, curare has to be tried in tetanus strychnine poisoning and hydrophobia, and would perhaps be still more used if reliable preparations of the drug were available. It has, however, no direct antagonistic effect; in fact, if applied to the central nervous system of the frog it causes convulsions and consequently its apparent antagonism would be of use only in allowing other methods or natural defences time to act.”

Since I have treated cases of tetanus by early serum injection with curarine hypodermically on the slightest sign of spasm, local or otherwise, I have had many gratifying successes. I never see a case of soil-infected wound that I fail to inject with antitoxin, and I always have at hand tablets of curare. The antitoxin should be injected into the nerve supplying the affected part, so as to intercept the toxin on its way to the central nervous system.

ART. XVIII.—*Acute Alcoholic Pancreatitis.* By WILLIAM ST. C. SYMMERS, M.B., Musgrave Professor of Pathology, Queen's University, Belfast.

AMONG the *post-mortem* examinations performed by me in Belfast during the past eight years there is a remarkable series of thirty-one cases in which the suddenness of death and the absence of any obvious cause necessitated a coroner's inquest. Most of these cases ended with startling suddenness. An apparently robust man, while chatting with a friend, collapsed dead on the pavement. Another man, while drinking on one of the cross-Channel boats, fell to the floor, and was dead by the time his boon companions could lift him up. Others, taken suddenly ill, died on the way to the hospital, or soon after admission.