Writers on hygiene and sanitation have paid but little attention to the subject of air vitiation by sulphur dioxide gas, consequently their information is very limited and it is no wonder, then, that one,⁶ among the rest, has made the careless statement in his otherwise very excellent work on hygiene that its effects "are exerted more on the digestive (?) than on the respiratory function."

From the presentation of all these facts it will appear evident that the weight of evidence concerning the relation of this gas to public health indicates that its effects are really "serious if not lasting" on certain susceptible persons, and that its presence in atmospheric air is a positive menace to the hygienic welfare of beings.

REFERENCES.

- ¹ Kionka and Ebstein: *Zeitschr. für Hygiene und Infectious Krankheiten*, 1902, vol. 14, p. 123.
- ² *A Manual of Hygiene and Sanitation*. By Seneca Egbert, A.M. M.D., 1900, p. 74.
- ³ *A Manual of Inorganic Chemistry*, vol. i, Non-Metals. By T. E. Thorpe, p. 215.
- ⁴ *Air and Rain*. By R. Angus Smith. London, 1872, p. 287.
- ⁵ *Chemistry, Inorganic and Organic*. By Charles Loudon Bloxam, 1895, p. 68.
- ⁶ *A Manual of Practical Hygiene*. By Charles Harrington, M.D., 1902, p. 251.
- ⁷ *A Text Book of Pharmacology and Therapeutics*. By Arthur R. Cushny, M.A., M.D., 1901, p. 549.
- ⁸ *A Text Book of Pharmacology*. By Torald Sollmann, M.D., 1906, p. 461.
- ⁹ *Grundriss der Toxikologie*. Von Dr. Heinrich Kionka, 1901, pp. 407-410.
- ¹⁰ *Lehrbuch der Intoxikationen*. Von Dr. Rudolph Kobert, 1904, vol. ii, 1st half, pp. 25-27.
- ¹¹ *Die menschlichen Nahrungs- und Genussmittel*. Von Dr. J. König, 1904, 11 Band, 1433-1434.
- ¹² *The Principles of Hygiene*. By D. H. Bergey, A.M., M.D., 1901, p. 100.

A CASE OF EXTENSIVE RESECTION OF THE STOMACH FOR CARCINOMA.*

BY FARRAR COBB, M.D.,

Assistant Visiting Surgeon, Massachusetts General Hospital.

No accurate knowledge of the extent of malignant disease of the stomach and the chance of its complete removal can be obtained by any other than operative methods. The case reported below is primarily of interest because it presents a strong argument in favor of operation in all cases of gastric cancer, even those which are apparently hopeless. The emaciation, severe pain and persistent vomiting, associated with the exceptionally large, palpable, epigastric mass in this case discouraged the thought of anything but a palliative gastro-jejunostomy, yet on opening the abdomen it was found possible to do a radical operation and by resecting about two-thirds of the stomach to cleanly remove the diseased area, leaving a good margin of healthy tissue. The case is interesting also from a pathological and etiological point of view, inasmuch as it furnishes a striking illustration of the ingrafting of carcinoma upon a previous gastric ulcer. The co-existence of a benign and a malignant ulcer side by side is shown in Fig. 1.

It has been estimated that fully 50% of the cases of gastric carcinoma will give a history suggestive of previous ulcer of the stomach. A majority of the cases of cancer of the stomach show

*This case was shown at the clinical meeting of the staff of the Massachusetts General Hospital on Jan. 11, 1907, and a brief report appeared in the *Boston Medical and Surgical Journal*, Feb. 7, 1907.

direct evidence of malignant disease developing on a previous ulcer. It is thought that in this case both ulcers were benign originally and that the cancerous degeneration set in not long before the time of operation. It is interesting, however, that it was not possible to obtain a history of long standing symptoms of indigestion. Undoubtedly much of the large mass felt in the epigastrium was due to chronic inflammatory thickening of the stomach wall; if the entire mass felt had been due to cancer there probably would have been no possibility of a complete removal of the disease, because of extensive glandular metastasis. The symptoms and signs, however, before operation were so markedly those of advanced

FIG. 1. The reconstructed stomach cut open from behind, showing the disease on the anterior wall.

Note the carcinomatous area near the pyloric end and the distinct benign ulcer toward the cardiac portion. Note also the marked thickening and infiltration of the anterior wall in general.

gastric carcinoma that one was justified in the belief that the case was one of hopeless cancer, demanding only a palliative operation. It is fair to state that a case presenting such a typical picture of inoperable carcinoma under the care of some physicians would not have received the benefit of a surgical consultation.

The fact that at the end of seventeen months after the operation the patient is alive and in apparently perfect health and engaged in hard manual labor justifies some hope of lasting cure. Such gratifying operative results ought to be reported, for only by so doing will practising physicians as a whole be convinced that it is their duty to send all cases of gastric carcinoma to the surgeon at the very first stages, even when the diagnosis is suspected only. As Mayo says: "In the history of medicine there is no recorded example of a case of cancer of the stomach cured by medical means. Of all diseases cancer of the stomach is most surgical, yet for some reason cases are sent to medical men, are entered in medical wards and subjected to treatment, which must result in 100% mortality."

The various steps in the technique of the operation as shown may be worthy of note.

REPORT OF CASE.

History. R. J. McD., fifty-two years old, white, married, teamster by occupation, entered the South Surgical Service of the Massachusetts General Hospital on Dec. 19, 1905. By the kindness of Dr. Beach, the chief of the service, the writer was given charge of the case and allowed to operate. Family history unimportant. He had always been in good health until seven months previous, when he first began to have indigestion and pain in the epigastrium, worse after eating. When first noticed the pain would come from an hour to an hour and a half after taking food. The severity and duration of the pain depended somewhat on the kind and amount of food taken. Later there was almost continual pain in the epigastric region, gradually increasing in severity and shooting up and down the right side and into the chest and back. He had steadily lost weight. He began to vomit about six weeks before entrance and continued to vomit three or four times a day, without any definite relation to eating. The vomitus for the most part was undigested food, but dark brown material was vomited occasionally. For two weeks before entrance he had been able to retain no food and had endured great pain.

Examination. — A pale and emaciated man; weight, 108 lbs. Heart and lungs normal; liver not enlarged. A large mass, slightly movable, dull on percussion, readily palpated in the epigastrium. The muscles in the epigastrium and the right hypochondrium were held somewhat rigidly. No fluid was made out in the abdominal cavity. No glands could be felt above the clavicles. By the stomach tube 50 cm. of thick brownish fluid mixed with mucus was obtained. An examination of this showed free hydrochloric acid, 0.18%; the total acidity was 0.365%; the guaiac test showed blood. The fasting stomach contained 30 cm. Inflation showed a small and contracted stomach. Examination of the urine showed nothing abnormal. Pain and vomiting continued after entering the hospital, but was not so frequent or severe as before.

Operation. — Twenty minutes before anesthesia an injection of $\frac{1}{4}$ gr. of morphine and 1-150 gr. of atropin was given. Ether was the anesthetic used. During much of the suturing of the stomach wall it was found possible to keep the patient only lightly under ether. An incision 4 in. long was made in the median line. Upon exploration it was found that a large portion of the anterior wall of the stomach was involved in an irregular, nodular mass, which extended from the middle line well toward the pylorus and was nearer to the greater than the lesser curvature, and in addition to this the major part of the anterior wall of the stomach seemed thickened and infiltrated with disease; the whole stomach seemed somewhat contracted. The extent of the diseased area is shown in Fig. 1, and is also shown, as well as the portion of the stomach resected, diagrammatically in Fig. 2. The stomach was not fixed by adhesions, except that in the gastrocolic omentum near the median line there were several enlarged glands adherent to the omentum and transverse colon. On microscopic examination these glands were pronounced simply inflammatory.

The steps of the operation are suggested in the diagrammatic illustrations accompanying this paper. Because of the presence of the diseased glands the gastrocolic omentum was tied off in sections first, care being taken to avoid the mid-colic artery. The enlarged glands and omentum were dissected off with care. The left gastro-epiploic artery was tied far out on the patient's left. After ligating the gastric artery

as high up as possible the lesser curvature of the stomach was freed and the superior pyloric artery tied. The stomach and pylorus could then be lifted well, so that hot gauze pads could be placed in the lesser omen-

FIG. 2. Showing application of clamps and portion of stomach removed. The lines of division were between the clamps.

tal cavity. Two straight narrow bladed clamps were then applied at about the juncture of the pylorus and the duodenum and two similar clamps on the left end of the stomach. The position of the clamps and their relation to the diseased area is shown in Fig. 2. After

FIG. 3. Redundant portion of the stomach wound closed by continuous catgut suture through all the coats preliminary to the continuous Cushing stitch to the peritoneum.

the abdominal cavity was carefully protected with gauze pads the duodenum was divided between the clamps and the pylorus and stomach with considerable omental fat and glands attached were lifted up. The stomach was then divided between the remaining

FIG. 4. The upper portion of the stomach wound has been finally closed by a continuous Cushing suture. Note that approximation of the stomach and duodenum has been begun with the continuous through-and-through suture of catgut.

FIG. 5. Approximation completed by a continuous Cushing suture with the addition of an occasional interrupted Lembert stitch.

clamps. The cut edges of the stomach and duodenum were carefully wiped clean with sponges wet in salt solution. It was found that the stumps of the stomach and duodenum were sufficiently mobile, after freeing the duodenum as much as possible to allow approximation end to end, as shown in Figs. 3, 4 and 5. Accordingly, all idea of complete closure of the stump of the stomach and the duodenum followed by gastro-intestinal anastomosis was abandoned and the end-to-end suture method adopted, as shown. Only very rarely will the stomach be sufficiently movable to allow the use of this method without undue tension on the sutures. In the great majority of cases where such extensive resections are done it will be necessary to resort to gastro-intestinal anastomosis.

After the introduction of the guide stitches the upper or redundant portion of the stomach wound was closed by using first a continuous suture of chromicized catgut, followed by a continuous Cushing suture of Pagenstecher thread. Approximation of the duodenum to the lower portion of the stomach wound was made by a continuous suture of chromicized catgut through all the coats, followed by a continuous Cushing stitch of Pagenstecher thread. It was found necessary to use a few additional interrupted stitches of linen thread. After the removal of the pads the abdominal wound was closed in layers, without drainage, leaving a quart of hot salt solution in the peritoneal cavity.

At the end of the operation the patient's pulse rose to 120, but dropped to 70 on the third day, where it remained. The temperature did not rise above 99.2° F. after the operation and remained normal after the third day. In other words, there was no serious reaction from the operation.

After-treatment.—The after-treatment consisted of salt solution by rectum 6 oz. every four hours, together with nutrient enemata of white of egg, beef juice, peptonized milk and salt solution every eight hours. Within twenty-four hours hot water and albumen water in teaspoonful amounts were given by mouth and on the third day the patient was taking $\frac{1}{2}$ oz. of water or albumen water every hour and retaining the enemata well. There was no vomiting.

The wound united by first intention and the stitches were removed on the eighth day. At this time he was taking full liquid diet freely. On the fourteenth day he sat up in bed and was up out of bed on the seventeenth day, when he was eating soft solids. Convalescence was uneventful and he was discharged one month from the day of the operation. He was then eating the regular hospital diet, with care as to amount. His weight on leaving the hospital was 118 lbs., or 10 lbs. over his weight on entrance.

Result.—This patient has been seen by me at intervals for the last sixteen months, and as was stated, he is apparently in perfect health. His present weight is 140 lbs. He eats anything and everything without any distress. He does not vomit and has no symptoms of indigestion. There is no evidence of a recurrence of the disease.

Pathological report.—The two ulcerated areas and extent of the disease are shown in Fig. 1. The large ulceration nearer the pylorus was 5.5 cm. in greatest diameter; its edges were raised and infiltrated. Microscopic examination of the indurated border showed masses of irregular tubular glands, with little fibrous or cellular tissue between them infiltrating the sub-mucous layers of the stomach wall, — *adeno-carcinoma*. Close to this malignant ulceration was a circular depressed cicatrized area, evidently the seat of a former ulceration. Microscopic examination showed simply granulation tissue, without any evidence of new growth.