operation was often as high as 50 or 75 per cent. Indeed, we have heard it positively affirmed that every fibromyoma of the uterus which could be recognized at the examining table was to be regarded as practically an indication for operation. In spite of the improved statistics of hysteromyomectomy, we would do well to heed the warning that there will always be unavoidable deaths after the operation.—ED.]

Ultimate Results of Myoma Operations.—BURCKHARD (*Zeitschrift für Geb. u. Gynäkologie*, Band xliii., Heft 1) concludes an extended article on this subject as follows: The operator need not be deterred from leaving the cervix by any fear of its subsequent malignant degeneration.

With few exceptions the ultimate results are good, such psychical disturbances as follow being transient. After entire removal of the ovaries hemorrhage ceases, and the tumor ceases to grow; it usually diminishes.

Menstrual molimina are rarely noted after the removal of both ovaries; when one or both is left after extirpation of the uterus these may occur.

In short, since the patient's health is not injured, but, on the contrary, the usual nervous disturbances are sensibly diminished by leaving the healthy ovaries after removing the uterus, the writer believes that this should be done whenever it is possible.

Epithelial Ingrowths in the Myometrium.—MEYER (*Zeitschrift für Geb. u. Gynäkologie*, Band xliii., Heft 1) describes the microscopical appearances in sections made through seven uteri, which were extirpated on account of obstinate hemorrhages, which had not been relieved by frequent curettement and cauterization. In five nothing abnormal was noted to account for the bleeding except glandular processes which dipped down into the submucous muscular layer. The direct cause of the hemorrhages was inferred to be certain degenerative changes in the bloodvessels, which were more or less constant. The writer believes that the cell-processes were the result of local irritation, and that the hypertrophy of the muscle in this vicinity was sufficient to cause a marked increase in the blood-pressure.

If this supposition is correct, total extirpation is the only cure for such hemorrhage, and is fully justified in view of the tendency of these benign glandular in-growths to become malignant. The curette, however vigorously used, does not reach the seat of the trouble.

The positive diagnosis of this condition is seldom possible clinically, but it should be suspected when no improvement is noted after repeated curettings, while the microscope shows no evidence of malignant disease. On the contrary, it should not be forgotten that the hemorrhage may be due to a small polypus at the fundus or at the cornua.

[This paper is exceedingly suggestive in that it throws new light upon a class of cases which often cause the surgeon great anxiety, lest he may, on the one hand, advise a radical operation for a condition in which it is not justified, or, on the other hand, may counsel delay in a case of incipient carcinoma when early intervention is important. Doubtless the most conservative will agree with the writer, that the persistence of hemorrhage after repeated curetting, especially in a woman above forty years of age, amply justifies total extirpation.—ED.]