promptly by the appropriate remedies—venæsection, leeching, calomel and opium, &c.

In other respects, if the patient go on well, she must be treated as after natural labour.

ART. XVI.—*Medical Observations, &c.* By SIR FRANCIS SMITH, M.D., of Paris, Fellow of the College of Physicians, Dublin.

IRREGULARITY OF THE DURATION OF THE FIRST STAGE OF
SCARLATINA.

WE are sometimes struck in practice with the very great non-accordance of symptoms, both as to duration and degree, with what is laid down by systematic writers; and junior practitioners must always be glad to have opportunities afforded them for storing their memory with facts irregular in their occurrence, and dissonant from the usual or routine rules laid down by authors. Among such facts may be placed the following:

S. S. a child of three years and a half old, robust and healthy, was exposed to the full infection of scarlatina by communicating freely with his sister, aged six months, who was carried off on the third day after she was attacked, and without throwing out any thing like eruption. The latter was brought to town ill from the country, and died on the third day, not having been previously in communication with her brother. On the day following her death,—that is to say, on the fourth day from the period when the infected child first was brought in contact with the healthy one,—the latter became sick, pale, with a shrunk and quickened pulse, and a disordered stomach; and this state continued for seventeen days, during which period the most gentle possible interference of a medical nature was adopted, with attention to diet, &c., but avoiding all such medication or disturbing influences as could interfere with the

march of the suspected malady; exercise according to strength, and amusements of a childish nature, were not suspended. The long continuance of this state, with disturbed functions, began to give rise to uneasiness; and I consulted with my esteemed and learned friend Doctor Mott, of New York, upon the case, and he advised a continuance in the expectant course which had been adopted. On the seventeenth day the eruption of scarlatina made its appearance and came freely out, the latter stages of the disease running a severe but regular course.

It would be manifestly improper to call the period of seventeen days which elapsed between the appearance of the eruption and the exposure to contagion, the latent period; for from the very commencement there existed the phenomena of fever, and we must therefore consider that period to have constituted the first stage of the malady, and its occurrence as one of the irregularities of disease which it is well to bear in mind as a possible contingency.

This case contrasts strongly with two others which I have lately seen, in which the first stage either did not exist at all, or at least was so faint as to elude the observation, either of the individuals who were the subjects of the disease or of those immediately about them.

Miss E. J., a blooming girl of thirteen years of age, whilst apparently in perfect health, and having, on the morning of the same day, eaten her usual breakfast with excellent appetite, complained of sore throat, and looking very much flushed, with a hot skin, I was asked to see her. When I visited her I found her with the rash of scarlatina already slightly developed, and ordered her to bed,—an order which she very reluctantly obeyed. The case ran a favourable course, and the eruptive stage lasted but the usual period, counting from the day on which I first saw her. Dr. Scott of Bologne saw the case with me.

Miss C., Rue de l'Observation, aged twelve, of a delicate strumous constitution, complained on the 12th of the month of sore throat and pains in the limbs, with headach, &c., having,

up to that day, been in her usual state of health. She was given some slight aperient medicine, and I was asked to see her on the 13th, on which day I found her in bed, with the eruption of scarlatina well developed on the whole of her limbs and body. Her mother, a lady much versed in sick attendance, and a most watchful observer of the health of her children, was quite persuaded that up to the 12th there had been no rigors, no fever, no diminution of appetite or derangement of other functions; the eruptive stage ran a normal course.

I shall but add one more case to the foregoing. Master E., son of Commissary-General E., was suddenly attacked with a violent convulsive fit, in which I was called to see him; no previous history could be obtained in the least calculated to shed light upon the case; the child, about seven years of age, had been in good health till the invasion of the convulsive attack, for which I was called to treat him; under these circumstances I ordered a small bleeding to be made at the arm, and a stimulant injection and warm bath to be administered; and as I was not the family attendant, advised the parents to summon their usual medical adviser. I did not again see the patient, but Mr. Cusack was subsequently in attendance with Dr. M'Munn, and the child, I understood, sunk rapidly without further development of the nature of the disease, but a light was but too soon shed upon the case by the appearance, in other children of the family, of a virulent scarlatina, which carried off, I believe, four children.

The last case is particularly worthy of observation, as it shews an instance of one of the most irregular, and at the same time formidable, shapes in which an eruptive fever can commence; and was, probably, a hopeless case from the beginning, had even its nature been clearly foreseen. A blow had been struck upon the nervous system, under which it was unable to rally; and though the other cases in the family were doomed to end equally fatally, in these, however, no obscurity existed as to their nature.

The above cases by no means include all the anomalies to be met with in the first stages of scarlatina; they will, perhaps, however, tend to bring the young practitioner's mind to appreciate the value of storing up facts, and being ready to apply them extemporaneously in the course of his practice.

THE USE OF POULTICES IN INFLAMMATIONS OF THE GREAT CAVITIES.

Nothing is more remarkable in the practice of physic in France, than the constant use of cataplasms in the treatment of inflammations of the great cavities. The moment inflammation is supposed to exist in the chest or abdomen, the first order given is to apply cataplasms. Where a practice prevails so generally, it is worth while to inquire into its value, and to examine whether its advantages might not entitle it to a favourable consideration, where its use has hitherto not prevailed.

The benefits to be derived from the application of poultices over inflamed cavities, are analogous to those sought from, in their application to other inflamed parts, and may be stated to consist, first, in maintaining the temperature of the parts over which they are applied, and thus giving nature a better chance of effecting the resolution of the existing inflammation; secondly, by their moisture mechanically aiding in the relaxation of injured tissues; and thirdly, by drawing the capillary circulation towards the surface, acting the part of derivatives or revulsives.

That poultices are capable of influencing the contained organs in the great cavities, some have affected to doubt; but yet, few will be found to deny the influence of cold upon the same organs, when applied externally. A moment's consideration would convince any one that organs such as the lungs and pleuræ, the abdominal viscera, and peritoneal coverings, sympathising as they do with their superjacent integuments, and frequently suffering from the accidental application of cold thereto, would be susceptible of considerable modification from the soothing, relaxing, and derivative effects of emollient poultices, applied over

their investing parietes ; and that much good effects may be derived from the careful and dexterous application of poultices, I am daily more and more convinced ; but I am also persuaded, that upon the mode in which they are applied, much of their good effects depend ; and I am free to confess, that a careless or slovenly application of them may, so far from proving serviceable, be productive of great mischief.

The usual material for poultices is the linseed meal ; and they are always included within linen cloths. The poultices are generally so large as to cover the seat of inflammation. They should be removed and renewed every three or four hours. Particular attention to the latter circumstance is desirable, as the edges are liable to grow cold, and a disagreeable and dangerous dissipation of animal temperature is the result ; and in this and in their weight consist, I think, the principal disadvantages of the system.

In the bronchial irritation attendant upon the exanthematous fevers, poultices to the chest are universally employed ; and where any want of energy appears to exist in the system, preventing the coming out of the eruption, cataplasms, consisting of the usual linseed meal, with the addition of a very small quantity, say a twelfth or a fifteenth part of mustard meal, (not flour,) are applied to the feet, and maintained for days together : thus keeping up both the temperature of the extremities, and gently stimulating the skin. The same mixture is also, when necessary, applied to the calves of the legs and thighs—the practice in the latter instances being similar in intention to that adopted in Great Britain ; but from the mustard meal being less irritating than the flour, and being used in so small a proportion, the sinapized cataplasms can be longer retained in their situation.

Pleurodynia, diaphragmitis, and many other affections of a painful nature, are likewise treated by poultices ; and it is certain, that their effects are often, in such cases, very satisfactory.

NEW PREPARATION OF IRON.

A new and valuable preparation of iron has lately been introduced into practice ; and in most cases, where chalybeate medicines are indicated, it possesses advantages over all other forms in which iron can be administered ; I allude to the Lactate of Iron. Its superiority would appear to depend partly upon its very great solubility, and partly, perhaps, upon its constituents, and its being more grateful than others to the digestive organs. Its use in private practice here is already very considerable ; and from the trials which I have given it in some cases, I am disposed to think highly of it ; but its price, which is considerable, renders it as yet unattainable by the poor. The forms in which it is generally administered are the fluid form, in distilled water, or in troches or pastilles ; and in many cases, combined with chocolate, in the form of lozenge.

CHOCOLATE.

Though so little used in England, chocolate enjoys a high and increasing reputation on the continent, and is not only made the vehicle of many different medicines, but is very extensively prescribed for children and delicate persons, as an article of nourishment. It is generally eaten dry, sweetened with sugar, and flavoured with many different matters, and formed into sticks or lozenges. Its manufacture is particularly attended to, and the consumption is enormous. Its medicinal introduction into Great Britain is, I think, much to be desired.

SALIVATION FROM IODINE CONTRASTED WITH THAT FROM MERCURY.

In the July number of this Journal I made mention of a case in which I had met with salivation from the use of hydriodate of potass. The fact of salivation occurring from the use of iodine or its preparations is not new : I have, since the publication of the article referred to, met with it in two instances, in one that of a young gentleman using the hydriodate of potass, for a scrofulous necrosis in the bones of the legs ; the other

that of a young lady, who, for a glandular disease of the neck, was put under the influence of iodine, administered in the watery solution as recommended by Dr. Osborne, of Dublin, at the rate of a grain to the pint; both had been under the influence of the mineral during nearly four months, and had been copiously salivated, and required mercurial purgations, and the administration of bark for the relief of the ptyalism. In each of these cases the entire absence of fetor of the breath was remarked, as was mentioned to have been the case in that which was recorded in the July number. Now, it is worthy of inquiry, why, when fetor is so constantly found to attend salivation by mercury, none should accompany salivation by iodine or its salts? I believe the difference to depend upon the fact, that in the action of mercury upon the mouth, there is not only an action induced upon the salivary glands, but also upon the mucous membrane of the mouth, from the alteration in the structure of which latter tissue arises the fetor. If the mucous membrane of the lips, cheeks, or gums be examined with a powerful lens, at the period when mercurial action is commencing in them, a species of ulcerative absorption will be observed to be in progress, which, being increased by the continuance of the cause, soon ends in ulcers of greater or lesser size, perceptible to the naked eye. On the contrary, in the salivation by iodine, the principal, if not the only action of the medicine, is directed to the salivary glands; and though I have in one case observed, with great surprise, the teeth to be loosened, I have never been able to discover any thing like ulcerative action to exist in the mucous membrane of the mouth, and upon this peculiar difference in their relative actions depends, I believe, the presence or absence of fetor of the breath during salivation induced by mercury, and iodine or its compounds.

SECRETION OF AIR FROM THE HUMAN SKIN.

I shall now state a case illustrative of a phenomenon which, as far as I am aware, has never before been observed or recorded;

and as I do not attempt to found any theory upon it, more especially as it, as yet, stands alone, I shall be very brief in what I have to say.

To the majority of my readers it requires no argument to prove, that various tissues in the human body, when in a state of disease, may secrete air. Hunter, J. P. Frank, and Professor Graves, have made statements more or less directly illustrative of my position. The latter extensive and enlightened observer, to whom I am proud to acknowledge myself indebted for much valuable information upon the subject upon which I now write, has seen cellular emphysema occur as a consequence of excessive hemorrhages. He has also seen air accumulated in the chest, as the result of pleuritis; and the ephemeral annals of medicine furnish several instances of the secretion of air in closed cavities.

Whenever animal decomposition or the fermentative process can be considered the cause of the production of the gaseous fluid; it, of course, may be put out of the question; but unless we admit that the mucous membrane of the stomach may occasionally take on the function of secreting air, I know not how we shall be able to account for those large and sudden disengagements of gas from that organ in certain hypochondriacal and hysterical subjects. In the class of persons I allude to, the disengagement of gas is but an occasional symptom, and is generally accompanied with other signs of disordered functions. It generally occurs when digestion has proceeded about one half, or to express it differently, about two or three hours after meals; but it also sometimes occurs when the stomach has been long empty. It is, sometimes, accompanied with a copious secretion of acid; but in, I believe, by far the larger number of cases, when a deficiency of acid in the stomach is observable, and in subjects who are liable to it, it may, and often is, brought on by any disagreeable mental emotion. Its early appearance after fresh food has been taken into the stomach, would forbid the idea, even were we unaware of the antiseptic properties of

the gastric secretions, that putrefaction of the contents of that viscus could have any thing to say to the phenomenon in question. And the occasional occurrence of the symptom when the stomach has long been empty, equally forbids the idea that it owes its origin to fermentation; and besides the extreme rapidity of the disengagement, and the immense amount of the gas evolved, tend to make us reject the idea that fermentation in the comparatively inconsiderable mass of the ingesta could satisfactorily account for the almost alarming amount to which the disengagement in some instances extends. Persons afflicted with the symptom I allude to, in any thing like an exalted degree, may be observed during the existence of the fit to hold their hands pressed upon the epigastrium, whilst discharge after discharge takes place; and should any suspension of the eructations be induced, either by a feeling of shame, hurry or other causes, they complain of a sense of faintness, and even occasionally go off in syncope. These phenomena, I submit, cannot be satisfactorily accounted for by any other supposition than that of a sudden taking on by the stomach of an abnormal secretory process.

In general, pathological physiologists readily admit the faculty of certain tissues to absorb air, with which they have, for a long time, been in contact, and occasionally this absorption occurs with vast rapidity; examples of which are every day to be found in the absorption of effused or rather injected air in traumatic and interlobular emphysema; and it requires but little stretch of imagination to conceive, that where absorbents or veins may take on a process of absorbing a gaseous fluid with which they may find themselves accidentally in contact, so likewise may exhalants be capable of eliminating or secreting a similar fluid, although they, in so doing, take on for the moment an unusual or abnormal function. But the case I am about to communicate was so unexpected and so novel, that I have hitherto abstained from publishing it, in the hope of either being

enabled to make fresh observations on it, or of meeting with others of a similar nature, in both of which hopes, however, I find myself as yet disappointed ; and I, therefore, give it to the Profession without comment, wishing that others may supply an explanation for phenomena, which have deeply interested me, but for the explanation of which I feel myself unable to offer any plausible hypothesis.

On the 12th May, 1840, I was consulted by an hypochondriac gentleman, Mr. G. æt. 35, who was labouring under what he chose to term gout, but whose symptoms I recognized for those of biliary derangement ; and I prescribed for him accordingly. Amongst other statements, he informed me that he was liable to immense disengagements of gas from the stomach, which he voided by eructation ; and he complained of enduring much distress whilst the fit was upon him ; that he was also liable to an occasional discharge of air from the urinary bladder, and besides, to an escape of air from his skin, and which latter symptoms he only observed whilst in the bath, of which he made frequent use. I confess I paid but little attention to the latter statement of symptoms ; and it was only upon his requiring to know from me the cause of those circumstances, that I thought fit to tell him that I thought that perhaps he might be in error upon both points ; and I explained to him how easily he might be misled in his observations. He replied, that he did not despair of convincing me of the facts being as he related them ; and thus the matter rested. On the morning of the 15th, I was hastily summoned to see my patient in a bath. He told me, on my arrival, that the reason of his sending for me, was for the purpose of convincing me of the truth of his statements as to the exudation of air or gas from his skin ; and as he only occasionally was able to demonstrate it, he lost no time in informing me of its occurrence. I found him in a bath at 79° of common river water (Seine) ; and upon examining his chest, shoulders, abdomen, and hands, I found him literally covered with small gas bubbles. I inquired of the attendant of the es-

tablishment, if he could account for the circumstance, and he said he had never observed anything of the kind before. I then caused my patient to lift his hands alternately from the water, when, of course, the air bubbles disappeared from those extremities; and on his returning them to the water, and retaining them near the surface, I could plainly perceive bubbles make their appearance, at first very minute, but gradually increasing in size, till they reached that of a pin's head, and the palms of his hands again become coated with them. I caused him to wipe away the bubbles from his hands and chest repeatedly, and in every case they were gradually replaced by others; they ran together, when pushed with the finger, like globules of quicksilver, and could be moved about upon the surface of the hands or body, without quitting them, or becoming loose in the water. These experiments and observations continued nearly twenty minutes that I was present; and towards the end of the bath, the edges of the metallic reservoir were coated all round, but more particularly towards the upper end, where the shoulders had been, for a depth of from one to two inches, with small air bubbles, as if its contents had been charged more or less with carbonic acid gas. Mr. G. also expected to be able to exhibit to me an evolution of gas from the urinary bladder; but on his attempting to do so, was unsuccessful.

I may observe that, to use his own words, he had had a very bad fit of belching when I prescribed for him, and which had given way under the treatment, and much more rapidly than usual,—similar attacks generally continuing more or less during a week or ten days at a time; and on the morning of the 15th he had felt *unusually strong and well*, as he walked to the baths. The next day he started for Weisbaden, in order to make use of the waters of that place for his gout, which I failed in convincing him to have no existence except in his own mis-conceived notions. I have not since heard from him, and the hopes which he held out of letting me see him on his return from Germany, have not been fulfilled.

Upon the above case, I shall abstain from all comment, save to add, that I went to its investigation, fully convinced that its subject was one of that numerous class of persons who live amidst self delusions ; and that, upon the subject of exudation of air from his skin, he was deceived. I came away convinced that a real secretion of air had actually taken place in my presence, and to a considerable extent, and that from a membranewhich I did not previously believe could, under any circumstances, take on that function.