2. ENDOCRINOPATHIES.

Maisonnet. ACUTE ADDISON'S DISEASE FOLLOWING DESTRUCTION OF THE RIGHT ADRENAL BODY. [*Presse Médicale*, Jan. 2, 1919.]

The author reports the case of a man who had been hit in the right flank by a shell fragment. He soon developed slight hematuria, without abdominal symptoms. In the succeeding days a distinct Addisonian syndrome came on, with discoloration of the mucous membranes, ochre color of the skin surface, Sergeant's white line, low blood pressure, greenish brown urine, pronounced asthenia, and a small, frequent pulse. The patient died on the tenth day. The autopsy showed an oblique tear of the lower pole of the right kidney, the vessels of the hilum being, however, intact. The right adrenal was completely destroyed. The shell fragment was found embedded in the body of the tenth thoracic vertebra. The left adrenal body was very small, measuring one half by one centimeter; it contained no tuberculous lesion. The clinical picture and ultimate death are doubtless to be ascribed to this anatomical peculiarity, Sergeant and others asserting that Addison's disease does not result from unilateral adrenal disease.

Comby, J. SUPRARENAL HEMORRHAGE IN CHILDREN. [*Arch. d. Med.*, Dec., 1918.]

The author reviews the recent literature on hemorrhage in the adrenals in children. Actual apoplexy is inevitably fatal, but in minor grades of hemorrhage low blood pressure, intermittent cyanosis and asthenia develop. Large and repeated doses of epinephrin are indicated. Vollbrach's case of purpura a year before developing Addison's disease is quoted.

Zueblin, Ernest. PITUITRIN AND ADRENALIN INJECTIONS IN HAY-FEVER. Cincinnati, Ohio. [*New York Medical Journal*, Vol. CVIII, p. 51, 1918.]

Report of a series of 6 cases with enlarged heart, chiefly of the right ventricle, with indistinct heartsounds, low systolic and high diastolic bloodpressure, increased or decreased cardiac strain pro minute, marked signs of vasomotor depression. The dose of adrenalin varied from 0.5 to 0.2 c.c., the dose of pituitrin from 1 c.c. to 0.25 c.c. At first the dose of adrenalin was high, then gradually diminished whilst the pituitrin dose was gradually increased, depending upon the heart, bloodpressure findings. The injections were repeated whenever any tendency of cardiac weakness was noticeable, first every other day, then twice a week and gradually discontinued. The subjective improvement after these injections were favorable as regards the severity and the duration of these attacks, in a number of cases the hay-fever symptoms disappeared without recurrence extending over an observation period of several years. Cases of hay-fever can be favorably helped by these injections,