excited in all probability by the laxative given the night before, somewhat exhausted her. In a few hours uterine contractions began and the membrane soon ruptured. The rush of liquid that followed was something appalling. The patient was nearly inundated and everything upon the bed was saturated. The tumor disappeared, and coincident with the rush of the waters, a fetus of five and a half to six months was expelled alive, but soon perished. The amount of amniotic fluid was so great that the patient had to be removed to another mattress as the one upon which she was lying was soaking, and the liquid dripping through it upon the floor. In grasping the fundus with my left hand I noticed it was peculiar in outline, and a little larger than usual. Suspecting twins, I carefully introduced my right hand, and encountered a foot. A second fetus, a little smaller than the first, which had evidently been dead for some days, was promptly delivered. The placenta soon followed, and a good cricket-ball contraction was obtained. An examination of the placenta, cord, and fetuses were next carefully made. The first fetus was well-developed and healthy. The second was smaller, less-developed, bones of skull collapsed, and body bloodless. They were both males. There was one placenta and two cords. The cord leading to the first fetus was healthy, full of blood, and pulsating at birth; that leading to the second was about half as large and bloodless. The placenta was divided into two unequal parts by a sharp line of demarcation. The larger part was that through which the living fetus received its nourishment, and was healthy in appearance. The smaller part was of much lighter color, bloodless, and had the appearance of being shrivelled; in its central part were several dark clots, more or less adherent to its surface, which had evidently been there for some time and been subjected to pressure.

There was nothing specific in this case, and the parents had a good hereditary history. The patient made an uninterrupted recovery.

The appearance of the placenta would indicate that there had been an accidental hemorrhage, which had caused the death of the second fetus. The sudden disturbance of the placental circulation, incident to its death, may explain the rapidity of the accumulation of liquid which resulted in an hydramnios. From the time this liquid began to accumulate until the premature birth, was but seven days, and the case is remarkable, not only for the rapidity and enormity of the distention, but for the very little constitutional disturbance which accompanied it.

A CASE OF URETHRO-VAGINAL FISTULA AND OTHER INJURIES FOLLOWING CHILD-BIRTH, WITH OPERATIONS.¹

BY EDWARD JACOB FORSTER, M.D.,
Visiting Physician at the Boston City Hospital.

I REMEMBER that, at one of our meetings, it was stated that vesico-vaginal fistulæ were now more uncommon than formerly for two reasons: first, that the recent advance and greater interest in gynecology had caused all the old cases to have been operated upon; and second, that, with a more intelligent practice of obstetrics, such lesions were less likely to occur.

Such cases being now comparatively so uncommon,

¹ Read before the Obstetrical Society of Boston, March 8, 1890.

I report the following as of interest on account of the rather unusual seat of the fistula and the other injuries which attended the patient's delivery.

The following history is condensed from the hospital records:

Boston City Hospital, First Medical Service, October 1, 1889. M. P., twenty years, married, housewife.

Family History.—Negative.

Previous History.—Always well. Menstruation regular; quite painful; flow moderate. One child born September 4th, which died in two hours.

Present Illness.—Labor was quite long, lasting about thirty-six hours. It was not instrumental. Got up upon the eighth day, but felt so badly that she went back to bed again. Has been in bed most of the time since. Complains that the urine comes away all the time; also of a feeling of weight. No bearing down pains. Has pain in back if she sits up long. Urine scalds, and there is soreness of the vulva. Does not pass urine, as it drains away all the time; and this is worse in the night. Considerable headache. No cough. Bowels have not moved for four days. Appetite good. Sleeps poorly. Temperature 99.2°, pulse 120, respiration 24. Urine acid; color normal; specific gravity 1.020; albumen trace; slight sediment, pus, little blood, bladder and vaginal cells.

Physical Examination.—Well developed and nourished. Heart and lungs negative. Tenderness in lower portion of abdomen, in median line, and extending into left iliac fossa.

Vaginal Examination, October 3d, shows the presence of a large and deep tear in cervix, extending posteriorly. Above the arch of the pubis, about three-quarters of an inch within the vagina, beyond the meatus urinarius, there is a tear through vesico- or urethro-vaginal septum, one inch or more wide, measured at right angles to urethral canal. This is just where urethra enters the bladder. Large tear of perineum. All the ulcerated surfaces are red, and bleed easily. The uterus is well contracted. A tube is placed in bladder, supported by a pledget of iodoform gauze in the vagina, and cleansed and rearranged, p. r. n.

October 8th. Lacerated parts are healing.

October 15th. Vesical tube omitted. Douche of clear water given three times a day, for cleanliness. Very great improvement and contraction of lacerated parts since entrance.

October 22d. Comfortable. Douches continued.

October 31st. Under ether, Dr. Forster operating: A denudation made of a strip about one-third of an inch wide about the vaginal side of the margin of the fistulous opening at the neck of the bladder. The edges drawn together by six silver-wire and three silk sutures. A soft rubber catheter retained. Recovery from ether accompanied by more than the usual vomiting. Antiseptic pad. Patient comfortable in evening.

November 3d. Douche: hydronaphthol, 1:5,000, twice a day.

November 7th. Bowels moved a little to-day of their own accord. Catheter causes discomfort.

November 10th. Wire and silk sutures removed to-day. Good union, except a small slough size of pea, at about middle of wound. Patient kept clean by douching three times a day. Cervix contains a considerable amount of glairy mucus. Painted with iodine.

November 14th. Under ether, Dr. Forster freshened

edges of the present fistula, and closed by interrupted silk sutures. Recovery from ether satisfactory. Sim's catheter retained constantly.

November 20th. Slight chill to-day. Temperature rose to 102°. Patient thought some urine which escaped around the catheter had leaked through old place. Examination found result of operation good. Sutures removed.

November 23d. Under ether, Dr. Forster operating, edges of the tear of cervix, which is in median line, and posterior, and extends to vaginal wall, are freshened, and united with four wire sutures. Patient then turned on her back. Butterfly-shaped denudation made on perineum. United by six wire sutures. Antiseptic pad. Recovery from ether protracted: intoxication violent for a time. Catheterized every four hours. Champagne, one ounce every three hours.

November 24th. Patient comfortable. Impossible for any of the nurses to pass catheter, so catheter retained to-day.

November 26th. Catheterized every four hours. Rose spray, antiseptic pad, etc. Yesterday a douche of corrosive sublimate (1:5,000) given; repeated daily.

November 28th. This morning, at five o'clock, a hemorrhage from the vagina occurred. House officer called. Hottest douches that could be borne, for one and a half hours proved futile to stop the flow (110°-120° temperature of douche). Patient put on her back. Infantile speculum introduced towards arch of pubis, and vagina emptied of fresh forming clots, and by delicate manipulation packed with small bits of glycerine and cotton tampon.

November 30th. Packing removed. Douche given. Perineum looks all right.

December 3d. Comfortable, hungry; douche daily.

December 4th. Stitches from perineum removed by Dr. Forster.

December 12th. Pain this morning. Examination finds tenderness in posterior cul-de-sac, about wires which remain in cervix. Pain relieved by tampon soaked in solution of morphia.

December 10th. Tinct. nucis vom., ten minims three times a day.

December 15th. Three stitches removed from cervix to-day by Dr. Forster. Cervix looked well. Uterus in normal position.

December 19th. Has been troubled by incontinence of urine the last few days. Pad applied over the vulva.

December 24th. Pad enables patient to retain urine well. Feels perfectly well. Wishes to go out. Discharged well, at own request.

Since the patient left the hospital, I have seen her several times. At first, she had difficulty in holding her urine for more than an hour, or when bending over about her work. A cotton pessary overcame these difficulties, and now she reports herself as quite well.

—There are to be various congresses of women engaged in different pursuits, in Chicago at the time of the World's Fair. In describing what would be done in this direction, one of the committee referred to the members of the different professions as women doctors, female nurses, and lady lawyers. The *Medical Record* asks, "What is the difference?"

THE CONSERVATIVE PHASE OF HYPNOTISM IN THERAPEUTICS.

BY CHARLES H. HARWOOD, A.B.

In a recent article in this journal by Dr. Hamilton Osgood,¹ attention was called to the therapeutic value of hypnotism. Dr. Osgood defined hypnotism by saying that, "In every-day English it means the act of putting a person to sleep"; adding, with italic emphasis, "The character of the sleep is normal." He even declared, "It is absurd to consider the sleep neurotic, pathological, and especially hysterical." Such statements deserve most careful consideration from practising physicians, for upon their substantiation or disproval largely depends the practical value of hypnotism in therapeutics.

The etiology of hypnotism shows that it is not simply "putting a person to sleep." A frightful clash does not put one asleep, and Charcot hypnotized his epileptoids at the Salpêtrière² by the stroke of a gong. Hypnotizing is done by engaging the attention on a stimulus, sometimes subjective, but usually a monotonous sensory stimulus. This stimulus is seen alike in the lancet case of Braid, the blackboard chalk-marks used by Professor James, the ticking of a watch used by Heidenhain, and the passes used by Dr. Liébault. As would be expected, the application of this stimulus is followed by a diffuse nervous discharge, varying in its effect from an almost imperceptible acceleration of respiration and pulse, to a most violent hysterical attack. As in the use of hypnotic drugs, however, the nervous disturbances produced by the irritation of first contact are lessened in successive applications, so in hypnotism this discharge often decreases with successive sittings, and "old subjects" may show very little of it, the eyes perhaps being the only evidence of its existence. Yet the production of a state of apparent unconsciousness, preceded by a state of nervous excitement, does not suggest the simple process of going to sleep.

The character of the sleep is not normal. In normal sleep, the blood-circulation in the brain is reduced; but Heidenhain writes that he hypnotized his brother after giving him nitrite of amyl. Not only has hypnotism been called "artificial catalepsy," but most of the arbitrary lists of "degrees" or "stages" of hypnosis (of which each writer on the subject has furnished his own) have one degree called "cataleptic." It is a free use of technical terms to call a sleep "cataleptic" and "normal" too. Unexpected noises usually awaken a person from normal sleep, but suggest to a hypnotized patient that all is quiet, and the shriek of a whistle will not awaken him. Strümpell³ says: "There is no hypnotic phenomenon which can be artificially produced in healthy persons but has its perfect analogue among the symptoms of hysteria, and is referable to the same main conditions." Paul Richer agrees with Strümpell in the analogy of hypnotism and hysteria, declaring their etiology to be the means of distinction. He says that they are nearly related. The best evidences there are of an analogy between sleep and hypnosis, somnambulic phenomena, not only are abnormal, but usually point to a pathological state of the brain resulting from overwork or ill health. Moreover, somnambulism has yet to show all the phe-

¹ The Therapeutic Value of Suggestion during the Hypnotic State, etc. Boston Medical and Surgical Journal of May 1st and 8th, 1890.

² Etudes Cliniques sur la Grande Hysterie. P. Richer.

³ Text-book of Medicine, p. 763.