tuberculosis as a lever in the broader campaign for better medical supervision of all persons, for no disease is more dependent upon general living conditions than is tuberculosis.

As health educators, finally, anti-tuberculosis associations must strive to stop the waste arising from the development of more and more specialties. I have spoken of over-specialization as one of the dangers to which the tuberculosis campaign has led. A string of "public health movements" has followed in the wake of the anti-tuberculosis campaign. There are now societies to prevent infant mortality; to promote better obstetrics and establish pre-natal clinics; societies for school hygiene, for mental hygiene, etc. Each of these agencies is pushing forward special health needs, is advancing a special technique, and is putting doctors, clinics and visiting nurses into the field. Even in smaller communities often two or three, or perhaps four, of these special health interests will be actively engaged in field work. The anti-tuberculosis association may well strive to pool the health interests of a locality and coördinate the special clinics or nurses in one coöperative group, whether they are maintained by a city or by private agencies. Clinics may be combined in one building, the work of different nurses may be coördinated or generalized; and this without the impairment of the integrity of the different organizations supporting the branches of the work. Such pooling of health interests will mean financial economy and gain to public health. Inasmuch as no movement is more dependent for its success on the general advancement of public hygiene than is the anti-tuberculosis movement, none will profit more by this coördination of special activities.

The time has come when tuberculosis associations must be health educators, because this is the more effective way to be tuberculosis educators. The anti-tuberculosis campaign must be a public health campaign, if it is to be most effective. Let public health be on the banner borne on the front of the car, while the public interest in preventing tuberculosis, already aroused, will furnish the gasoline.

VIII.

THE RESPONSIBILITY AND OPPORTUNITY OF THE TEACHER IN PREVENTING TUBERCULOSIS.

By THOMAS F. HARRINGTON, M.D., BOSTON,

Director of School Hygiene, Boston.

MUCH of the apparent disagreement among authorities relative to the prevalence of tuberculosis among children is due to the fact that a distinction is not made between a tuberculous infection and a tuberculous disease. It is now admitted generally that all persons are infected with tuberculosis at some period of life. The great majority recover without ever knowing

that they had been infected. In childhood, it is very difficult to detect tuberculosis either by physical examinations or by recognized tests. Reliable data at hand demonstrate that relatively few infected children go on into the tuberculosis state. All infected in the first year or two of life die. Only ten per cent. of all deaths from tuberculosis occur during the ages of five to twenty years, inclusive. Even in families of the tuberculous, the infection of the exposed children is hard to demonstrate. In Boston, there are 7000 cases of positive tuberculosis on record at the office of the Health Department; these people live in 1340 families in which there are 2131 children of school age. A recent study of these children, by which the school nurses secured a report from the family physician or from a hospital clinic, showed that only 304 of the children had tuberculosis in a form that could be diagnosed. A later study of 196 children diagnosed as positively tuberculous confirmed this diagnosis in only eleven cases. Fourteen children of the group were diagnosed as doubtful; forty were deferred for subsequent observation, while 92 children of the group were returned to school as negative cases. A study of over 90,000 pupils in our elementary grades shows that five per cent. of the children were anemic, glandular, or under-sized. In other words, they were physically below par to a degree requiring preventive measures if they were to be saved from genuine tuberculosis. All reliable evidence today demonstrates that the education of the public in the cultivation of hygienic habits of living is the most effective means of combating tuberculosis. Obviously this leads us at once to the classroom. Here are grouped by compulsion, for a long period of the formative age of children, about one-fifth of the population. The opportunity to establish right habits of living is far greater in the school than at any other period or place. This the state of Massachusetts recognized early in the enactment into law that "schools shall give instruction in physiology and hygiene," and further that "in each of the subjects of physiology and hygiene, special instruction as to tuberculosis and its prevention shall be taught as a regular branch of study to *all pupils in all schools* supported wholly or partly by public money." Other subjects *may* be taught, e.g. algebra, geometry, language, natural science, kindergarten training, manual training, etc., if the school committee thinks such teaching expedient. Furthermore, the statute provides that every city and town of 10,000 or more inhabitants shall maintain a school in which the subjects of physiology and hygiene shall be taught, and towns which do not maintain a school as required by the statute shall not be entitled to its portion of the appropriation allowed from the "Massachusetts school fund." Nor does the responsibility of the school end here. A further act is mandatory in its provision that "the school committee of each city and town shall appoint one or more school

physicians, who shall make a prompt examination and diagnosis of all children referred to him, and such further examination of teachers, janitors, and school buildings as in his opinion the protection of the health of the pupils may require." Also that every child in the public schools shall be separately and carefully tested and examined at least once in every school year; and further, the school committee is required to keep a physical record of each child and to give notice to the parent or guardian concerning the defects or disability requiring treatment. The statute also places upon the State Board of Education and the State Board of Health the responsibility for carrying out the provisions of the act, including the giving of adequate instruction on these subjects to the pupils in the normal schools. Later legislation requires that all children under sixteen years of age, seeking a working certificate, shall be in good health, sufficient to undertake the work for which the application is made. We might go even further and point out the dangerous trades and occupations which minors, eighteen years of age and less, cannot enter; these trades and occupations require that teacher and school physician shall have some knowledge of the processes involved if they are to act as guides to the pupils entering vocational life. The limited time at my disposal necessitates the restriction of the whole problem to two points only, namely, first, in what manner is physiology and hygiene teaching being carried out; and secondly, what can this state anti-tuberculosis association do to aid in efforts to meet the requirements of the statutes on these subjects. Let us divide the problem into:—

1. The teaching of physiology and hygiene.
2. Medical inspection.

THE TEACHING OF PHYSIOLOGY AND HYGIENE.

Data collected recently from most of the cities and from many of the towns justify only one conclusion, namely, that no subject in the curriculum is so neglected or so inadequately taught as the subject of physiology and hygiene. Many superintendents write candidly that nothing is being done in their communities to carry out this law. Others present a course of study in which less time is allotted weekly to this subject than to any other subject in the daily program. In only a few cities is any special instruction given in the prevention of tuberculosis. There is little or no definite policy carried out, other than in notable exceptions, covering classroom temperature and open windows. Only two courses of study show any correlation between such habits as cleanliness, good teeth, food, sleep, fresh air and sunshine, and the daily home life and school life of the child. The opinion of the superintendents seems to be unanimous that the cause for this state of affairs lies in the lack of preparation of the teacher for the responsibility placed upon her by the statute. In the normal school, where any attention is given to this spe-

cial subject, the course of study is so scientific and so overloaded with anatomy often that it defeats its own object,—namely, to give the teacher-pupil a preparation in methods of teaching the subject in the grades.

The prevention of tuberculosis is not given serious consideration in many of the normal schools.

MEDICAL INSPECTION.

Here the situation is akin to that of physiology and hygiene teaching—namely, inadequate preparation for the work and the lack of a just appreciation of the responsibility involved. Many physicians here and there are making great personal sacrifices to do this work well. On paper, many cities have a very attractive organization; too many communities fail to recognize the tremendous power of health supervision in raising the physical and mental standard of the group as well as of the individual. In only a few cities and towns is there any team work between the health and the educational departments. Little or no medical inspection can be effective unless the teacher is prepared to select the children needing medical care. Normal schools and medical schools are not giving their pupils special preparation. Medical inspection without a "follow up" by school nurses is in the vast majority of cases (more than 90%) a waste of time and statistical gathering only. No remedial or corrective results follow such an inspection. On the other hand, there is a decided trend to rely upon the teacher and nurse combination without the guidance of a school physician. No school principal nor superintendent should take the responsibility of this problem unaided by some medical authority. In summing up, therefore, it can be said that an organization such as this association can do a great good in cities and towns, first, by urging school and health authorities to get together to carry out the statutes quoted above; second, by requesting school authorities to legislate that windows in class-rooms be kept open and that room temperature be kept between 60 and 65° F; third, by establishing open-air classes for the debilitated children who are not frankly tuberculous; fourth, by providing literature on tuberculosis prevention for distribution in the schools; fifth, by demanding that teachers and physicians be prepared, in normal schools and in medical schools, for this special work.

IX.

MEDICAL INSPECTION OF SCHOOL CHILDREN.

By FRANCIS LEE DUNHAM, M.D., BROOKLINE, MASS.

INVESTIGATIONS in the department of school hygiene comprehend as many factors of social as of medical or educational importance. Consequently an intimate coöperation between the