ART. LXVII.—*Post-Glacial Faults at St. John, N. B.*;
By G. F. MATTHEW, D.Sc., F.R.S.C. (With Plate XI.)

NOWHERE, so far as the writer is aware, has there been put on record any account of disturbances in post-glacial times, of the rocks along the lines of faulting, in this North-eastern part of America. There are numerous places in all parts of this region where the removal of drift deposits allows of the inspection of the glaciated surfaces of the rocks, and had such faults been common it would seem highly probable that their presence would long ere this have been noticed. This thought induces the writer to give a brief sketch of the numerous small displacements, produced since the glacial period that have been observed in the rocks in and near St. John, N. B., Canada.

Numerous proofs of post-glacial disturbances of the earth's crust are visible in the slate rocks which form a large part of the rocky foundation on which St. John is built. These are true *post-glacial faults*. They are well shown along the hillside south of "The Valley" in that city. This valley is an east to west depression in the northern section of the thickly inhabited part of the town.

The rocks along the southern slope of this valley consist of the fine, dark-colored shales or slates of the Bretonian or Upper Division of the St. John Group; which by their fine grain and uniform texture, are well adapted to preserve the sharp grooves and striæ impressed upon them in glacial times, by the moving ice.

The faults visible along this hillside belong chiefly to a system having approximately a N.E. to S.W. course, and a hade of

* Univ. of Chicago, 1893-4. Results not yet published.

† Phil. Magazine, vol. xxxviii, p. 379, October, 1894.

from 60° to 80° to the S.E. But these faults are traversed by diagonal ones, some with a course approximately N. to S. and others E. to W. The principal movement of faulting in post-glacial times has been along the fault planes that have a N.E. to S.W. course.

The place where these faults were first observed was on the side of City Road, north of the City Hospital. Here a glaciated surface of slate about twelve feet wide has been exposed by the removal of the surface soil. There are nine faults visible here having a course nearly parallel to that of the street and a hade to the S.E. They vary in downthrow from one quarter of an inch to four inches, the downthrow being in all cases on the *north side*.

Beside these faults of the northeast and southwest system, two faults diagonal to these were seen, these have a course approximately N.E. to S.W. and the downthrow in both cases, though small, is on the *north side*.

On the steep hillside east of the end of Charles Street, numerous small faults of the N. E.-S. W. system were seen. In all these the downthrows were on the *north side*. Here the diagonal faults are of more importance as regards the downthrow that at the exposure on City Road; the hade of these faults is also less, being about 50° ; the course of these is different from that of the diagonal faults at City Road being approximately E.-W. The downthrows here are on the *north side* of the diagonal faults and on the *north side* of the faults that have (approximately) a northeast to southwest course.

The best display of the S.E.-N.W. post-glacial faults that has been observed is that on Rock street behind Fowler's Mill where a space of about one hundred and seventy feet of slate ledges has been laid bare. In this space sixty-one faults were observed which had been attended with displacement of the measures since their glaciation. In all of these faults, with one exception the *downthrow was on the north side*, and the hade of the faults was to the southwest. The sum of the displacement at these faults was five feet eight inches; the average throw for each fault being thus somewhat over one inch.

In one case only, a fault of half an inch had a downthrow on the *south side*. The diagonal faults which are not very conspicuous at this place, were not measured.

The heaviest throw seen at this place was five inches and very often an individual fault when traced along its course was found to vary in the amount of displacement. This was especially the case when the fault was complicated with a diagonal displacement, so that wedges of the slate would be raised above the general contour, or depressed below it; and occasionally a

rhombic block could be observed, lifted above the surrounding slate.

At almost every place in the city and east and west of it, where the glaciated ledges of the slates of the St. John Group could be observed, some displacement was seen, though often very slight. At the N.E. corner of the Church of England burying-ground, east of the city, there is a ledge which shows two slight faults ($\frac{1}{4}$ inch each) with a downthrow *on the south*, eastward of Courtney Bay, at its head, the slates at the shore and on the road, have about a dozen faults with throws varying from one half an inch to an inch; these in almost all cases have the downthrow *on the north side*.

A fuller account of these faults will be given in the next Bulletin of the Natural History Society of New Brunswick, with some conjectures as to the causes which have produced them.

St. John, N. B., Canada, November 1, 1894.

EXPLANATION OF THE PLATE.

A ledge of clay-slate on City Road, St. John, N. E., Canada, showing a series of post-glacial faults. The view is taken looking southward up the hillside. When exposed to the weather the sharp edge of the slate along the fault crumbles away from the effect of frost, and in the larger faults is only found perfect when the fault has been freshly exposed. In most of the faults here figured the projecting edge has thus suffered from exposure, except where they approach the covering of soil. The course of the glacial stria is S. 15 E.