This article was downloaded by: [University of Arizona]
On: 17 December 2014, At: 17:13
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 7

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/tnah13>

XIX.—On a second collection of reptiles made by Mr. E. Lort-Phillips in Somaliland

G.A. Boulenger F.R.S.

Published online: 22 Sep 2009.

To cite this article: G.A. Boulenger F.R.S. (1898) XIX.—On a second collection of reptiles made by Mr. E. Lort-Phillips in Somaliland , Annals and Magazine of Natural History: Series 7, 2:8, 130-133, DOI: [10.1080/00222939808678026](https://doi.org/10.1080/00222939808678026)

To link to this article: <http://dx.doi.org/10.1080/00222939808678026>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content

should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

EXPLANATION OF PLATE VII.

- Fig. 1. *Trachycystis lignicola*.
 Fig. 2. *Ennea elliptica*.
 Fig. 3. — *formosa*.
 Fig. 4. *Succinea piscinalis*.
 Fig. 5. *Buliminus zuluenensis*.
 Fig. 6. *Hapalus globosus*.
 Fig. 7. *Pupa Farquhari*.
 Fig. 8. *Physa gradata*.
 Fig. 9. *Cyclophorus minimus*.

XIX.—*On a Second Collection of Reptiles made by Mr. E. Lort-Phillips in Somaliland.* By G. A. BOULENGER, F.R.S.

SINCE the publication in these 'Annals'* of a report on a small collection of Reptiles made by Mr. Lort-Phillips at Berbera and in the Goolis Mountains the British Museum has received a second collection made by the same gentleman in the same district in 1896-97. It contains examples of most of the Reptiles mentioned in the first report, and, in addition, of eight species, two of which are new, enumerated in the following list.

LIZARDS.

1. *Hemidactylus mabuia*, Mor.

Goolis Mountains.

2. *Agama Rueppellii*, Vaill.

Goolis Mountains.

3. *Agama (Xenagama) batillifera*, Vaill.

Found in holes on the Sogsoda plain, Goolis Mountains, at an elevation of about 2000 feet.

4. *Latastia Hardeggeri*, Steind.

Latastia Hardeggeri, Steindachner, Ann. Hofmus. Wien, vi. 1891, p. 371, pl. xi.

Eremias heterolepis, Boettger, Zool. Anz. 1893, p. 115.

This species is very closely allied to *L. Neumanni*, Matschie, from Aden, recently described by J. Anderson (P. Z. S. 1895,

* Ser. 6, vol. xvi. 1895, p. 165.

Downloaded by [University of Arizona] at 17:13 17 December 2014

Downloaded by [University of Arizona] at 17:13 17 December 2014

Downloaded by [University of Arizona] at 17:13 17 December 2014

Downloaded by [University of Arizona] at 17:13 17 December 2014

Two specimens from Berbera.

This species is very closely allied to *L. spinalis*, Ptrs. Apart from the coloration, fundamentally different in the light stripes being in odd number, it may be distinguished by the longer foot, the fourth toe measured from the base of the fifth exceeding the distance between the fore limb and the anterior border of the orbit.

6. *Chamæleon basiliscus*, Cope.

Goolis Mountains.

SNAKES.

7. *Typhlops cuneirostris*, Ptrs.

Goolis Mountains.

ÆLUROGLENA, gen. nov.

Maxillary teeth 11, increasing in size posteriorly, forming an uninterrupted series; mandibular teeth subequal. Head elongate, slightly distinct from neck; eye moderate, with vertically elliptical pupil; nostril pierced between two nasals and the supranasal. Body slender, cylindrical; scales smooth, with apical pits, in 21 rows. Ventrals rounded. Tail moderate; subcaudals in 2 rows.

This genus should be placed not far from *Zamenis*.

8. *Æluroglena cucullata*, sp. n.

Snout rather long, slightly prominent. Rostral much broader than deep, not visible from above; internasals as long as the præfrontals; frontal bell-shaped, once and a half as long as broad, broader than the supraocular, as long as its distance from the end of the snout, shorter than the parietals; loreal twice as long as deep; a single large præocular, narrowly separated from the frontal; two postoculars; temporals 1+2; eight upper labials, fourth and fifth entering the eye; five lower labials in contact with the anterior chin-shields, which are shorter than the posterior; latter separated from each other by scales. Scales in 21 rows. Ventrals 216; anal divided; subcaudals 67. Pale buff above; head dark brown above, speckled with whitish, shading into a blackish blotch on the nape; two large cream-coloured spots on the upper lip, in front of and behind the eye; lower surface of head dark brown, with a cream-coloured spot on each side of the chin, and a band of the same colour below

each mandibular ramus, sending up a process to meet the præocular labial spot; the brown is prolonged as a median stripe along the throat; lower parts cream-colour.

Total length 375 millim.; tail 75.

A single female specimen from the Goolis Mountains.

XX.—*Descriptions of Two new Fishes from the Coast of Sind.*

By G. A. BOULENGER, F.R.S.

THE fishes here described were found among a collection made at Kurrachee by Mr. F. W. Townsend, and presented by him to the British Museum.

Epinephelus Townsendi.

Teeth in rather broad bands, in three series on the sides of the mandible; canines small. Depth of body equal to length of head, $2\frac{3}{4}$ to 3 times in total length. Snout rather pointed, as long as diameter of eye, which is $4\frac{1}{2}$ to 5 times in length of head; interorbital width 6 to 7 times in length of head; lower jaw strongly projecting; maxillary extending to below posterior border of eye or a little beyond, the width of its distal extremity $\frac{2}{3}$ to $\frac{3}{4}$ diameter of eye; præopercular border rounded, finely serrated behind, the serræ slightly enlarged towards the angle; opercular spines equidistant, lower further back than upper; opercular flap pointed, upper border feebly curved; head covered with cycloid scales, maxillary naked. Gill-rakers moderate, 16 or 17 on lower part of anterior arch. Dorsal XI 16–17, originating above base of pectoral; spines increasing in length to the third or fourth, which is $\frac{1}{3}$ length of head and not, or but slightly, longer than the last, and a little shorter than longest soft rays. Pectoral $\frac{2}{3}$ length of head. Ventral shorter, not reaching vent. Anal spines strong, second and third equal, nearly $\frac{1}{3}$ length of head. Caudal rounded. Scales strongly ciliated, 95–110 $\frac{13-15}{36-40}$; l. l. 67–76. Olive-brown above, whitish beneath; one or two more or less distinct dark streaks on the cheek and gill-cover; dorsal, anal, and caudal edged with blackish.

Total length 250 millim.

Four specimens.

This species is nearest allied to *E. merra*, Bl.