ward from adult to child. This is a retrograde means of acquiring knowledge which has proved eminently unsuccessful. In the Old World, as well as in the New, these false methods have been pursued. What little has, by more rational methods of study, been accomplished in the investigation of infants and children has as large a place in America as in Europe. Here, then, is our opportunity for original research, for we have a branch of medicine which is universally new; we have the same advantages in clinical material, in well equipped laboratories, in especial hospitals, and in the intelligence which is bred and born in our New England climate throughout all of our six New England States, as is to be found anywhere. Let us endeavor to give as much, if not more, to the *savants* of Europe as we receive from them. It is not an exaggeration to state that a large number of sick infants and young children throughout the land are suffering from the vigorous treatment of their over-zealous medical attendants rather than from the disease with which they started. This should be stopped.

I ask you to carefully consider this vital question. I ask you to attack it with iconoclasm and original thought, for I assure you it will repay you by opening up vast fields of intensely interesting brain work. It is not difficult to pick out instance after instance of the truth of what I have just said. The therapeutics of infancy and childhood, as understood by thousands of practitioners in Germany and France, in England and in America, is a wonderful exhibition of what vagaries the human mind, sound on other subjects, can be induced to indulge in.

A total disregard of natural processes, an over-exaggeration of symptoms, which in the adult mean danger and require active treatment, hold too great prominence in the mind of the general practitioner of medicine, since in the child they are but simple and harmless manifestations of an over-stimulated nervous system. Such is the "*ignis fatuus*" so misleading, not only to those who have not especially studied the early stages of life, but also to those who have seen much of it, but with eyes blinded by traditions of the past.

It would seem as though in this age of rapid discovery and quick change and interchange of ideas, — in an age when the precision of our investigations in all branches of research has almost placed the much-abused profession of medicine on a level with the exact sciences, — that we who are endeavoring to place our especial branch of study on firm ground should consider, with unusual care, the opinions which we express to the profession at large and to the laity. What we say in this society should be founded on principles as accurate as the science of the day can make them.

There is one especial subject which I hope may be brought before this society during the course of the coming year, and that is, the endeavor to reform the nomenclature of the pediatric medicine of to-day. Some effort should be made to first determine upon the classification of certain important subjects, such as diseases of the mouth

and of the gastro-enteric tract, and then, as the exponent of some accepted classification, we should endeavor, through our influence as a society, to bring the various large teaching centers of our country to join with us in introducing and adopting such classification. Much of the present medical nomenclature is unintelligent and misleading. Many old terms, such as dysentery, phthisis, summer diarrhea, and many others should be dropped as mere relics of past ignorance.

What we need, and what the Pediatric Department of Harvard University has proved to be the best classification for practical purposes, especially in diseases of the gastro-enteric tract, is to place our classification of these diseases on an etiological basis. This we find can be done, and this we believe will enable physicians in different parts of the world to understand which particular disease their colleagues at a distance are talking and writing about. At present, many names are used for one disease, and again, many diseases are at times classified under one name.

This is only one of many subjects which we should investigate and promulgate in the future, and I trust that you will enter into work of this kind with enthusiasm, for I assure you it will not only aid you to appreciate the sick children who come under your care, but will help to place this especial branch of clinical medicine on a more intelligent basis. Let us all work shoulder to shoulder to advance the subject of the clinical medicine of early life which has grown so rapidly in the last twenty-five years, and which we can but feel will continue to have its horizon expand, until it equals that of any of the useful sciences in the world.

I welcome you, gentlemen, to the first regular meeting of the New England Pediatric Society.

Original Articles.

THE SERUM TREATMENT OF EPIDEMIC CEREBROSPINAL MENINGITIS.*

BY CHARLES HUNTER DUNN, M.D., BOSTON.

At the present day it is coming to be generally recognized by the medical world that in the study of the problem of immunity lies the key to further advance in the knowledge of the infectious diseases. To this study we must look for the results which are to be of the greatest future practical benefit to humanity. The complexity of the problem is so much greater than can possibly be realized by one not actually engaged in its study, that the progress of scientific research is necessarily slow, and the practitioner, while eagerly awaiting disclosures which shall be of value to him in his practice, has no time to follow each successive step, each little addition to our knowledge, which the unceasing patient efforts of investigators are reaching. He has difficulty, too, in choosing, from the complex

* Read before the first meeting of the New England Pediatric Society, Feb. 15, 1908.

mass of the results of scientific research, those which may be of practical value, and waits, perhaps too long, for proof of such value before contributing his part to the progress of knowledge. I believe the practitioner should play a greater part in this progress. The investigator on his side has neither the time nor the opportunity to prove the practical value of the results following his conclusions. He can only point the way. It is only by the earnest co-operation of the man engaged in the actual treatment of the disease, that valuable resources can be proved or disproved and that theoretical resources can be made practical and brought without excessive loss of time to the aid of the community.

I am presenting this paper at this time, not because the value of the serum treatment of epidemic cerebrospinal meningitis is proved, but because it is not proved. The number of cases in which it has been used is still too small to permit the drawing of final conclusions as to its value. My object is to enlist the co-operation of as many as possible of those engaged in actual practice, in the testing, confirming, or rejecting of its value, by showing how great, even at this early stage of the testing, is the hope of practical benefit held out by serum therapy in this disease, a disease occurring principally in childhood, and one of the most fatal.

During the last few years, several methods of using specific therapy in epidemic cerebrospinal meningitis have been suggested and applied in practice. In the winter and spring of 1906 the prevalence of this disease in various parts of the country attracted unusual attention, and a number of publications appeared, reporting favorable results from the use of diphtheria antitoxin in meningitis. In this case the practitioner outran the course of scientific research, for the use of diphtheria antitoxin was not based on any proper scientific foundation, no evidence being offered of its containing any antibody specific toward meningitis, but was based either upon pure empiricism or the vague hope that among its antibodies might be one of efficacy against the meningococcus. This treatment was thoroughly tested by Dr. Rotch at the Children's Hospital, with no further result than the showing that the giving of frequently repeated large doses of diphtheria antitoxin to cases of meningitis is harmless. And the final verdict from other localities also is that no specific benefit is to be derived from its use.

In 1907, great interest had been awakened by the ingenious theory of the opsonin as brought forward by Wright, which attracted especial attention to the aspect of a primary artificial immunity and to the possibility of treating infections by means of vaccines. During this year Dr. Rotch treated his cases of meningitis with vaccines prepared from the meningococcus, given according to the indications shown by the determination of the opsonic index, as suggested by Wright. While the results showed some evidence of favorable action, they were totally inconclusive, too inconclusive to afford much basis

for hope that this would be the solution of the practical problem.

In the meanwhile investigators have been working upon the production of an antiserum specific toward the meningococcus. The recent work of Flexner, at the Rockefeller Institute for Medical Research, has brought such an antiserum into the domain of practical realization, and it now remains to confirm the value of this antiserum in actual practice.

The Flexner serum is prepared from horses inoculated with the meningococcus in a similar way to diphtheria antitoxin, and, like diphtheria antitoxin, contains specific immune bodies. It differs from diphtheria antitoxin in that it does not, as in diphtheria, neutralize the poison formed by the specific organism, but hastens disintegration of specific organism, and prepares it for phagocytosis. It is bactericidal rather than antitoxic in its nature, wherefore it should not be called meningitis antitoxin, but meningitis antiserum.

The antiserum was proved by Flexner's experiments to have protective power in laboratory animals, monkeys being used. During the past year it has been tested upon human beings in several series of cases of epidemic cerebrospinal meningitis. The report of these cases has just been published by Flexner and Jobling in the *Journal of Experimental Medicine* in January, 1908. The results reported are so favorable that they hold out the greatest possible hope for practical benefit, though the number of cases is still too few for the drawing of final conclusions.

The method of using the serum recommended by Flexner is to inject it directly into the spinal canal. This is done by means of lumbar puncture and just as much fluid as will run freely is allowed to escape and then a syringe filled with the antiserum is connected with the needle through which the fluid has escaped and the serum is injected through this same needle. The questions of the dose and the frequency of the injections, are, of course, still to be determined. Flexner suggests 30 ccm. as the maximum dose, to be repeated daily for three or four days, and that at least as much cerebrospinal fluid as this should be first withdrawn, in order to avoid the risk of an undue increase of cerebral pressure.

Since last November, I have used Flexner's antiserum in 15 cases of epidemic cerebrospinal meningitis, in all but one of which the diagnosis was confirmed by the finding of the diplococcus intracellularis in the cerebrospinal fluid. Of these cases, 8 have resulted in complete recovery, 2 have resulted in death and 5 are still pending. The 8 cases which recovered are all perfectly well, having been left with no sequelæ of any kind, an unusually favorable result in this disease. The 2 fatal cases were both chronic cases, in which the disease had run considerable time before coming under my observation. Of the 5 cases which are still pending, 4 are now convalescent and will undoubtedly recover; the other is a chronic case, in which the outcome is dubious.

Every one of the 8 cases to which I have given the serum in the first week of the disease has resulted in complete recovery, and of the pending cases, two in which the antiserum was given early, are convalescent.

the only early case which did not show a marked immediate improvement after the giving of the antiserum. The doses given early in this case were exceptionally small, 5 ccm. to 15 ccm., as very little fluid could be obtained by lumbar

Case.	Day of disease.	No. of doses.	Amount of dose, ccm.	Total serum given.	Immediate effect.	Duration of fever.	Duration of symptoms.	Result.
1	2	6	5-30 ccm.	105 ccm.	Temporary fall in temperature.	23 days. 99-105	20 days	Complete recovery
2	8	8	15-30 ccm.	200 ccm.	Temporary fall in temperature.	16 days 99-103	16 days	Complete recovery
3	14	2	15 ccm.	30 ccm.	None.			Death
4	3	4	30 ccm.	120 ccm.	Marked relief of symptoms.	6 days 99-103.6	3 days	Complete recovery
5	21	8	25-40 ccm.	325 ccm.	Permanent fall in temperature.	1 day 101-102		Death
6	2	3	30 ccm.	90 ccm.	Critical fall in temperature to normal. Relief of symptoms.	5 days 98-101	2 days	Complete recovery
7	1	1	30 ccm.	30 ccm.	Permanent fall in temperature to 99.	1 day 98.4-99.6	4 days	Complete recovery
8	4	4	15-30 ccm.	100 ccm.	Marked relief of symptoms.	5 days 98-100	1 day	Complete recovery
9	1	2	30 ccm.	60 ccm.	Fall in temperature to normal. Marked relief of severe symptoms.	12 days. 99-100	2 days.	Complete recovery
10	2	1	30 ccm.	30 ccm.	Fall in temperature to normal. Complete relief of symptoms.	12 hours	1 day	Complete recovery
11	21	4	45 ccm.	180 ccm.	Slight fall in temperature. No relief of symptoms.	2 days		Unfavorable chronic case
12	5 (?)	3	15-30 ccm.	75 ccm.	Fall in temperature.	15 days 98-100	2 days	Convalescent.
13	8	2	30 ccm.	60 ccm.	None.	4 days 99-102	4 days	Convalescent.
14	5	4	30 ccm.	120 ccm.	Fall in temperature. Marked relief of symptoms.	6 days 99-103	6 days	Convalescent.
15	5	4	25-45 ccm.	125 ccm.	Fall in temperature. Marked relief of symptoms.	99-101	5 days	Convalescent. Deaf.

The two fatal cases were in children who had had the disease for several weeks and were already in the chronic stage when first seen. Both were unconscious, with normal temperatures, although in both there were a few organisms to be found in the cerebrospinal fluid.

The other unfavorable case was an adult in which the disease had existed for three weeks. There was some fever and the organisms were still present in the fluid, which was obtained in unusually large quantity. Forty-five cubic centimeters of antiserum were given on Jan. 23, 24, 28 and Feb. 3. The temperature came down to normal after the first dose and has been normal since but there has been no improvement in the general condition. No organisms were found in the fluid after the first puncture. I believe the continuance of symptoms in this case was due to the organic lesions in the brain, possibly an internal hydrocephalus resulting from the infectious process, which itself has ceased, leaving extensive damage behind. The patient is still alive, but is daily growing weaker.

Two cases, which ended in complete recovery, showed no immediate effect from the giving of the serum. In both a temporary fall of temperature followed the giving of the first dose, but both ran a prolonged course of fever, with gradual but steady improvement of the symptoms. One of these was an early case, and was

puncture and the failure of immediate improvement may be due to this fact.

Two cases, neither very early, showed no immediate effect after receiving the antiserum, but recovery was rapid, both as to the reaching of a permanent normal temperature and the disappearance of symptoms.

Eight cases, in which the serum was given early in the disease, showed a very marked, even startling, improvement immediately following the giving of the serum. In four of these cases there was an immediate and permanent fall of temperature, exactly resembling the crisis of a pneumonia, which was accompanied by a complete and permanent return of the mental condition to normal, complete and permanent disappearance of headache, and followed by rapid disappearance of rigidity of the neck and all other signs. Two of these cases, in twelve hours after the first dose, returned from a condition of complete unconsciousness to one of absolutely normal mental condition. In the other four cases, the temperature fell to the normal by a fairly rapid lysis, accompanied by rapid improvement in the mental condition and disappearance of symptoms and signs. In two of these, there was a rapid change from a condition of active, almost violent, delirium to one of normal mentality.

It is difficult to convey a full impression of

the remarkable behavior of these cases by mere statistics. Actual contact with them impressed upon me the contrast between their course and the course of similar cases of cerebrospinal meningitis untreated by the Flexner antiserum.

This series of cases is too small to afford a statistical basis for drawing any definite conclusions as to the final value of the Flexner antiserum in cerebrospinal meningitis. Much remains to be proven, and it is possible that my series included cases of exceptional character. I believe, however, that I am justified in urging a widespread trial of this treatment.

In the first place, I think we can conclude that the use of the antiserum does no harm. I understand from Dr. Flexner that I have used it in larger doses than any of his other investigators. In no case was there any sign of bad effect, and in two cases I injected it into the spinal canal without the previous withdrawal of fluid without any sign of bad result from increased intradural pressure. I would not recommend the use of this procedure without great caution.

While the percentage of recoveries in cerebrospinal meningitis is very variable, in my experience of this disease recovery almost invariably occurs after a more or less prolonged period of illness, often running into actual chronicity. A very large proportion of cases which do not die at an early stage pass into a chronic stage, the final outcome of which is doubtful. I have never seen nor even heard of a case proved by lumbar puncture to be true epidemic meningitis, which aborted, or terminated by sudden crisis at an early stage. The recovery by crisis and the rapid improvement followed by recovery immediately following the giving of the antiserum, in so large a proportion of this series of cases, is, I believe, strong evidence of the favorable specific effect of this treatment.

The completeness of the recovery of the cases in this series is another very important feature. It is well known how frequently permanent sequelæ occur in those cases of epidemic cerebrospinal meningitis which survive. Permanent deafness, blindness, paralysis, mental impairment and idiocy are common enough to make the prognosis dubious in all cases which survive the initial storm of the disease. In the 13 cases in this series now alive, 8 are now and have been for many weeks absolutely well, and 3 are fully convalescent without evidence of any unfavorable sequelæ. In one case now convalescent, there is marked deafness, the other being the chronic case in which there is probably hydrocephalus. I believe this number of complete recoveries is significantly large.

The most important point suggested by the results of the use of the Flexner serum, in this series of cases of epidemic cerebrospinal meningitis, is the advantage to be gained by giving it early in the course of the disease. Not only did all the cases in which it was given early recover, but all those cases in which its use was followed by a marked immediate improvement were cases in which it was given within the first few

days of the disease; and in two of these earlier cases, the disease was apparently completely aborted by one dose of the antiserum. I believe that the importance of early diagnosis and treatment cannot be overestimated.

I think the results of the Flexner antiserum in these cases are sufficiently good to afford a very strong basis of hope that this treatment will prove of great value in the treatment of cerebrospinal meningitis, a value commensurate with that of antitoxin in diphtheria. I believe, even though it requires further testing, the hope of good result is so strong that the antiserum should be used in every case of this very dangerous and fatal disease and as early as possible. In every case of suspected meningitis or even of possible meningitis, lumbar puncture should be made as soon as the disease is suspected and the physician should be prepared to give the antiserum at once. If the fluid obtained by the lumbar puncture is notably cloudy, the antiserum should be injected at once through the same needle, without incurring the loss of time caused by waiting for the result of the bacteriological examination of the fluid. The great majority of all cases with distinctly cloudy fluid, especially in children, are caused by the meningococcus; and in those rare cases in which cloudy fluid is found in tubercular meningitis, or in occasional cases of pneumococcus meningitis, no harm will be done. If, after one dose, the temperature falls to the normal and the symptoms show rapid and progressive improvement no further dose of antiserum may be necessary, but the physician should be ready at any time to repeat the treatment, if the temperature begins to rise or if the symptoms show a tendency to recur. Relapses do occur and should be treated exactly as the original attack.

If the temperature does not come down to the normal, or if the symptoms do not show progressive improvement, the treatment by lumbar puncture and [the injection of antiserum should be repeated daily for three days, making four doses in all. After this time further injections may be made, as indicated by increase at any time of fever or symptoms.

In those rare cases in which no fluid can be withdrawn by lumbar puncture, it is questionable how safe it is to inject antiserum into the canal. I have done it without bad result. It would be worth while in these cases to try subcutaneous injections.

Late cases afford a prospect of benefit, if there is still fever and evidences of an active infectious process or if there are still organisms to be found in the cerebrospinal fluid. The effect is less immediately noticeable and recovery much more prolonged.

Cases which have passed into the chronic stage without fever and with no organisms to be found in the fluid are not influenced by the antiserum.

CASE I. Charles Clausen, two years ten months. Admitted Nov. 6, 1907.

F. H. — Negative.

P. H. — Osteomyelitis at eighteen months.

P. I.—On Nov. 5, one day previously, patient began to vomit and vomited everything taken. Since, has been sleepy and feverish all the time.

P. E.—Well developed and nourished. Child is unconscious; eyes are closed; restless and tossing about at times; slight rigidity of neck; bending head forward causes cry and restlessness. Pupils equal and react normally. Strabismus at times; knee jerks normal; no Kernig. Babinski's sign doubtful. Temperature, 100° to 103°. Lumbar puncture, 25 ccm. turbid fluid withdrawn. Examination of fluid,—many polynuclear leucocytes; many Gram-decolorizing diplococci, both within and without the cells. Thirty cubic centimeters antiserum given.

Nov. 7. Restless night; fall in temperature this morning. Stiffness of neck more marked; Kernig's sign present. White count, 49,200. Lumbar puncture, 5 ccm. turbid fluid obtained. Examination same. Culture positive. Ten cubic centimeters antiserum given.

Nov. 8. Child is now conscious and much improved; not restless. Kernig and stiff neck still present. Pupils unequal at times. Lumbar puncture, no fluid obtained. Five cubic centimeters antiserum given.

Nov. 9. Condition about the same. Vomits at times. Lumbar puncture, no fluid obtained. Ten cubic centimeters antiserum given.

Nov. 10–25. Temperature has been elevated. Kernig and rigidity have persisted; but general condition has been good and no nervous symptoms; no vomiting or headache.

Nov. 25. Lumbar puncture, 50 ccm. fluid obtained. Very slightly cloudy. Examination: Centrifugalized fluid showed polynuclear leucocytes and few Gram-decolorizing diplococci, both within and without the cells. Culture positive for meningococcus. Thirty cubic centimeters antiserum given.

Nov. 26. Tenderness and rigidity of neck have disappeared. Lumbar puncture, 20 ccm. Very slightly cloudy fluid obtained. Examination showed polynuclear leucocytes and few Gram-decolorizing diplococci, both extra- and intra-cellular. Culture: no growth. Thirty cubic centimeters antiserum given.

Nov. 27. Temperature coming down. Lumbar puncture, 50 ccm. fluid obtained. Centrifugalized fluid showed a few leucocytes, but no micro-organisms. Culture: no growth.

Nov. 28. Lumbar puncture, 20 ccm. clear fluid obtained. Examination as on Nov. 27.

Nov. 29. Temperature reached the normal. Child is quiet; shows no symptoms except persistent Kernig's sign.

Nov. 30 to Dec. 4. No symptoms except occasional attack of vomiting. Child perfectly normal mentally, playing with playthings; Kernig has disappeared.

Dec. 4–31. Temperature normal except for two brief rises, which were unaccompanied by depression or symptoms of any kind. Nothing found to account for them. Child bright; talks and plays.

Jan. 1, 1907. Patient discharged to Convalescent Home.

CASE II. Bessie Koplin, three and one-half years. Admitted Nov. 13, 1904.

F. H.—Negative.

P. H.—Measles at one and a half years.

P. I.—Seven days ago, on Nov. 6, the child had a pain in the stomach and went to bed feeling dull and appearing quite sick. On the same day had several convulsions, vomiting and fever. The fever has persisted. The child has been in bed; is very dull, but recognizes parents.

P. E.—Poorly developed and nourished. Child

lies with eyes closed and head markedly retracted. When disturbed, it cries and resists with arms. Head retracted; neck rigid. Attempt to move the head seems to cause great pain. Knees held drawn up. Pupils slightly unequal at times; react to light. Knee jerks lively. Double Kernig's sign present. Babinski's sign absent. Temperature 103°. White count, 26,000.

Nov. 14. Condition the same. Temperature 104°. Lumbar puncture, 50 ccm. slightly turbid fluid obtained. Examination of fluid: sediment consists chiefly of polynuclear leucocytes; few Gram-decolorizing diplococci, both intra- and extra-cellular. Culture positive for meningococci. Thirty cubic centimeters antiserum given.

Nov. 15. Fall in temperature. Condition otherwise the same. Lumbar puncture, 50 ccm. slightly turbid fluid obtained. Examination of fluid, as on Nov. 14. Culture positive. Thirty cubic centimeters antiserum given.

Nov. 16. Temperature 101°. Condition unchanged. Lumbar puncture, 50 ccm. of slightly cloudy fluid obtained. Examination of fluid showed fewer leucocytes; very few diplococci and none found outside the cells. Culture negative for meningococci. Thirty cubic centimeters antiserum given.

Nov. 17. Child seems somewhat brighter. Notices surroundings, but lies quiet. Kernig's sign present. Retraction of head and rigidity of neck persist. Lumbar puncture, 15 ccm. of very slightly cloudy fluid obtained. Examination of fluid: few leucocytes; no diplococci found. Culture negative for meningococci. Fifteen cubic centimeters antiserum given.

Nov. 18–24. Temperature 100° to 102°. General condition good. Child quiet; conscious; restless when disturbed. Retraction of head has gradually disappeared. Rigidity of neck and Kernig's sign persist.

Nov. 25. Lumbar puncture. No fluid obtained. Fifteen cubic centimeters antiserum given.

Nov. 26. Lumbar puncture. No fluid obtained. Twenty cubic centimeters antiserum given.

Nov. 27. Lumbar puncture, 50 ccm. clear fluid obtained. Centrifugalized sediment showed occasional polynuclear leucocytes and lymphocytes in about equal proportions. No micro-organisms. Culture: no growth. Thirty cubic centimeters antiserum given.

Nov. 28. Lumbar puncture, 30 ccm. fluid obtained. Examination of fluid as on Nov. 27. Thirty cubic centimeters antiserum given.

Nov. 29 to Dec. 1. Temperature fell to the normal. No change in general condition. Child has no symptoms except slight rigidity of neck and double Kernig's sign.

Dec. 2. Temperature went up again, without any attendant symptoms.

Dec. 3–21. Constant irregular temperature occurs, varying between occasional remissions to the normal and 105°. Rigidity of neck disappeared about Dec. 4. Kernig's sign persists. During this time the child has had a constant succession of furuncles.

Dec. 21–28. No meningeal symptoms except the Kernig's sign, which disappeared about Dec. 27. Temperature normal. The child lies quietly, smiles occasionally, and talks volubly in Jewish tongue to parents.

Jan. 4. Discharged well to Convalescent Home.

CASE III. Jeffrey Morrison, sixteen months. Admitted Nov. 16, 1907.

F. H.—Negative.

P. H.—Negative.

P. I.—On Nov. 6, fever was present and child refused food; constipation; vomiting after feeding since onset. History unsatisfactory.

P. E. — Child very dull. Seems quite ill, but no definite signs found. No rigidity of neck, tenderness, nor retraction of head. Reflexes normal. No Kernig.

Nov. 18. Photophobia has become marked. This morning the left pupil is dilated and does not react to light. Vomiting has persisted, the child having been supported on nutrient enemata.

This case was diagnosed as *probable* TUBERCULAR MENINGITIS and was not reported to me. I happened to see it on Nov. 20, and suggested lumbar puncture.

Nov. 20. Lumbar puncture, 50 ccm. slightly cloudy fluid obtained. Fibrinous clot found on standing. The cells were polynuclear and mononuclear in about equal proportions. A few extra-cellular Gram-decolorizing diplococci were seen, morphologically resembling the meningococcus. Culture gave no growth. Thirty cubic centimeters antiserum given.

Nov. 21. Child has gradually become completely unconscious, with closed eyes and unequal pupils. No other signs. Lumbar puncture, 20 ccm. fluid obtained. Examination of fluid as on Nov. 20. Thirty cubic centimeters antiserum given.

Nov. 22. Died.

CASE IV. Joseph Hazelton, six and one-half years. Admitted Nov. 27, 1907.

F. H. — Negative.

P. H. — Negative.

P. I. — Yesterday, Nov. 26, in the afternoon, the child came in from his play with headache. Ate his supper, but vomited right afterwards. Had a comfortable night and seemed fairly well this morning; he was about the house until 1 p.m., when he suddenly became unconscious, and could not be aroused. Was brought at once to the hospital.

P. E. — Child unconscious; face flushed; head retracted; neck rigid; eyes closed, opening at times. Pupils are dilated; react to light. Signs of pain on extending lower extremities. Knee jerks lively. Double Kernig's sign present. Babinski's sign absent. Temperature 103°. Lumbar puncture, 15 ccm. very turbid fluid obtained. Examination of fluid showed sediment to consist chiefly of polynuclear leucocytes; many Gram-decolorizing diplococci, both extra- and intra-cellular. The sediment presents very much the same appearance as that from a fresh case of gonorrhea. Culture positive for the meningococcus. No antiserum given, as supply had not arrived.

Nov. 28. Temperature came down to normal during the night, but in the afternoon it went up to 104°. No change in signs. Lumbar puncture, 50 ccm. turbid fluid obtained. Examination of fluid as on Nov. 27. Thirty cubic centimeters antiserum given.

Nov. 29. Temperature again fell to normal during the night. Child returned completely to consciousness, complaining only of headache. Lumbar puncture, 30 ccm. fluid obtained; much less turbid. Examina-

tion as on Nov. 28. Culture positive for meningococcus. Thirty cubic centimeters antiserum given.

Nov. 30. General condition continues to improve. Moderate retraction of head and stiffness of neck. Pupils moderately dilated. Boy says his headache is almost gone. Double Kernig persists. Lumbar puncture, 30 ccm. fluid obtained. Examination of fluid showed fewer leucocytes and very few diplococci. None extra-cellular. Culture negative for meningococcus. Thirty cubic centimeters antiserum given.

Dec. 1. Very slight headache; slight retraction of head and stiffness of neck. Child is bright and takes an active interest in his surroundings. Temperature, 100°. Lumbar puncture, 30 ccm. fluid obtained; very slightly cloudy; no diplococci found. Culture gave no growth. Thirty cubic centimeters antiserum given.

Dec. 2-4. Temperature, 99° to 100°. Rigidity, retraction and Kernig's sign have all disappeared. No headache. Child seems perfectly bright and well.

Dec. 4-18. Temperature normal. There were no further rise and no further symptoms. Convalescence was very rapid, it being difficult to keep the boy quiet.

Dec. 18. Discharged well to Convalescent Home.

Jan. 1. Boy is perfectly well and very active.

CASE V. Emily Smith, one year. Admitted Dec. 10, 1907.

F. H. — Negative.

P. H. — Nine weeks ago had bronchitis followed by pneumonia, lasting four weeks. Cough has persisted.

P. I. — For three weeks the child has been vomiting two or three times a day and has wanted to sleep most of the time. Has taken food well. History very unsatisfactory.

P. E. — Emaciated; eyes sunken, with dusky rings under them. Head retracted; neck rigid; child is unconscious. Pupils dilated, equal, react to light. Anterior fontanelle is bulging. No Kernig's sign. Reflexes normal. Temperature, 102°. Lumbar puncture, 30 ccm. of turbid fluid with a thick fibrinous clot. Examination showed numerous polynuclear leucocytes and frequent Gram-decolorizing diplococci, both intra- and extra-cellular. Culture gave no growth. Twenty-five cubic centimeters antiserum given.

Dec. 11. After the giving of the serum, the temperature fell to normal.

Dec. 12 to Jan. 1. The temperature has been normal; the general condition has grown steadily worse. The child has gradually fallen into the condition of post-meningeal idiocy frequently seen. Vomiting has been constant and nutrient enemata have not been well retained. The record of lumbar punctures and antiserum is as follows:

	Fluid obtained.	Antiserum given.	Diplococci.
Dec. 10,	30 ccm.	25 ccm.	+
" 11,	40 "	30 "	+
" 12,	50 "	30 "	0
" 13,	30 "	30 "	0
" 18,	70 "	45 "	0
" 19,	60 "	45 "	0
" 20,	60 "	45 "	0
" 21,	30 "	45 "	0

Jan. 5. Died.

CASE VI. Albert Munroe, six years. Seen in consultation with Dr. F. W. Rice, Dec. 17, 1907.

F. H. — Negative.

P. H. — On Dec. 10, the little niece of the maid who took care of the patient had cerebrospinal meningitis and died on the fourth day. On Dec. 14, the maid was taken with the same disease. Albert Munroe had slept with this maid, but not after Dec.

10. He was, however, in more or less intimate contact with her up to the day she was taken ill.

Raymond Abbott, a playmate of Albert's living across the street, was taken with cerebrospinal meningitis on Dec. 21.

P. I.—The first symptoms began on Dec. 16. The boy was well and went to school in the morning, but in the afternoon had a chill. After the chill the temperature was 103° axilla and the boy complained of severe headache and was very apathetic. He vomited everything given. During the night the temperature fell and was normal at 8 A.M. This fall in temperature was accompanied by signs of collapse. The boy became pulseless, with cyanosis and cold extremities. He reacted to stimulation. After the collapse the patient was seen in consultation by Dr. John Lovett Morse, who advised the use of antiserum. I saw him at 3.30 P.M. He was still retaining nothing by mouth and was being given nutrient enemata.

P. E.—Well developed and nourished. Face pale; mucous membranes and finger tips slightly cyanotic. Expression dull and suggests pain. Mental condition very apathetic. Noticeable tenderness and rigidity at the back of the neck. Pupils equal and react to

light; not dilated. Knee jerks somewhat exaggerated. Babinski's sign present on the right. Otherwise, nothing abnormal. Temperature, 101° axilla. Lumbar puncture was done and about 90 ccm. of fluid withdrawn, not under pressure. The fluid was very cloudy, a thick layer of pus settling on standing. Examination of fluid: Numerous polynuclear leucocytes; frequent Gram-negative biscuit-shaped diplococci, both within and without the leucocytes. Culture gave a positive culture of a Gram-decolorizing diplococcus. Thirty cubic centimeters of meningococcus antiserum were injected into the spinal canal.

Dec. 18. Within a few hours after the injection of the antiserum, the boy's general condition began to improve. Vomiting ceased at once. The temperature fell to the normal. When seen, at 3.30 P.M., the change in the general condition was very marked. Temperature, 99°; pulse, 110 and of good quality; respiration, 28. Color good. Pallor and cyanosis have disappeared. Mental condition much brighter. Complains only of slight headache. Tenderness and rigidity of the neck have disappeared. Kernig present, but not marked. Plantar reflexes normal. Lumbar

puncture, about 20 ccm. of cloudy fluid obtained. Fluid was slightly less cloudy than on the previous day. Examination of the fluid showed fewer organisms. Culture positive. Twenty-five cubic centimeters antiserum injected into the canal.

Dec. 19. Improvement has been steady. There is a slight afternoon rise of temperature, it being normal in the morning. Boy was found playing with a Teddy bear. No headache; no Kernig; knee jerks and plantar reflexes normal. No lumbar puncture.

Dec. 20. Improvement continues. Boy seems thoroughly convalescent though there is still a slight rise of afternoon temperature. Lumbar puncture, 100 ccm. of clear fluid obtained. On examination a few leucocytes were found, but no micro-organisms. Culture sterile. Thirty cubic centimeters of antiserum injected into canal.

Dec. 22. No symptoms. Temperature, 99° at 4 P.M.

Dec. 26. No rise of temperature for 2 days. Boy seems perfectly well. Discharged well.

CASE VII. Raymond Abbott, seven years. Seen in consultation with Dr. F. W. Rice, Dec. 22, 1907.

F. H.—Negative.

P. H.—This boy was the chum of Albert Munroe (see case 6) who came down with cerebrospinal meningitis on Dec. 16. He played with him until that time.

P. I.—This morning, about 3 A.M., the boy became very feverish, restless and complained of headache. He vomited several times during the morning. Headache is referred to the back of the head. Temperature, 102°. Pulse, 100. Respiration, 28.

P. E.—Well developed, but thin. Good color. Expression dull and apathetic and boy is conscious. Slight tenderness and rigidity of the back of the neck. No retraction. Slight Kernig. Knee jerks lively. Plantar reflexes normal. An eruption on the chest and abdomen, somewhat resembling typhoid rose spots, but darker in color, although not petechial. Lumbar puncture, about 90 ccm. of fluid obtained. Fluid runs almost perfectly clear, a very faint cloud being detected. No clot found on standing. Examination showed very few leucocytes; a few red blood corpuscles and frequent Gram-negative biscuit-shaped diplococci, none being found in cells. Culture was positive for the meningococcus. Thirty cubic centimeters antiserum injected into the spinal canal.

Dec. 23. Temperature fell at once to normal after the injection. No more vomiting. Boy complains only of slight headache. Slight retraction of head. Tenderness and rigidity slightly more marked. Kernig's sign more marked. Knee jerks somewhat exaggerated. Plantar reflexes normal. No antiserum given.

Dec. 24. Dr. Rice telephones that boy is about the same as on the preceding day.

Dec. 25. In the afternoon Dr. Rice telephones that the patient is not so well. Had two or three attacks of vomiting during the day. He complains slightly of headache; seems more listless and the Kernig and rigidity of the neck have increased slightly. No rise of temperature with these symptoms.

Dec. 26. Seen at Brighton. Boy is better. Headache only on sitting up. Still somewhat listless as if tired. Vomiting has ceased. No tenderness or rigidity of neck and no Kernig. Reflexes normal. The symptoms of the previous day were probably due to the over-excitement of Christmas.

Dec. 27. Dr. Rice telephones that the boy is much better, with normal temperature. Much brighter.

Jan. 1. Discharged well.

CASE VIII. Adelaide Walker, ten years. Negress. Admitted Dec. 27, 1907.

P. H. — Well until present illness.

P. I. — On the night of Dec. 23, child came home bent over and complaining of abdominal pain. During the night she had a chill, followed by high fever and headache. Vomited the next day. Has been very feverish and restless, sleeping only a few minutes at a time, crying with pain in the back of the head.

P. E. — Well developed and nourished. Conscious; lies with eyes closed; expression of pain and constant cry. Head retracted; neck rigid, the slightest movement causing pain. Knees drawn up. Pupils equal, moderately dilated, react to light. Knee jerks lively. Double Kernig's sign present. Temperature, 100°. Lumbar puncture on admission gave 20 ccm. cloudy fluid. Examination of fluid showed sediment to consist chiefly of polynuclear leucocytes; few diplococci, Gram-decolorizing; intra- and extra-cellular. Culture positive for meningococcus. Thirty cubic centimeters antiserum given.

Dec. 28. Child is much better and is brighter in every way, showing complete change in general condition. Notices surroundings. Complains only occasionally of headache. Head is not retracted so much; neck less rigid. Lumbar puncture, 30 ccm. very slightly cloudy fluid. Few leucocytes; no micro-organisms found. Culture, no growth. Thirty cubic centimeters antiserum given.

Dec. 29. Improvement continues. Child got hold of a storybook this morning and read a story through. Temperature, 99° to 100°. No headache. Lumbar puncture, 15 ccm. clear fluid obtained. Few leucocytes; no micro-organisms; culture, no growth. Fifteen cubic centimeters antiserum given.

Dec. 30. Patient sang a song this morning and talks and plays a good deal. No headache. Slight stiffness of neck. No retraction; Kernig persists. Temperature normal.

Jan. 1. Temperature up to 100°. Patient seems perfectly well. Kernig and all signs have disappeared. Urine shows signs of pyelitis. Rapid, uneventful convalescence.

CASE IX. W. W. Weeden, Jr., six years. Seen with Dr. Walker, of Providence, at 2 A.M., Jan. 6, 1908.

Onset at 2 A.M., Jan. 5, with fever and slight headache. Was better during the morning, but in the afternoon became delirious, with high fever and convulsive twitchings. In the evening a complete collapse occurred, the child becoming unconscious,

with subnormal temperature, cold extremities and no pulse at the wrist. He reacted a little to stimulation. When I saw him at 2 A.M., the temperature was 106°, the child was unconscious with no radial pulse and widely dilated pupils reacting very little to light. There was slight retraction and rigidity of the neck, normal plantar reflexes, double Kernig, no knee jerks, and a fine petechial rash on the chest and back. Lumbar puncture gave 50 ccm. very turbid fluid. Examination of the fluid showed frequent Gram-negative diplococci, mainly intracellular. Culture was positive for the meningococcus, but they died out on the first

transplantation. Thirty cubic centimeters antiserum given.

At 9 A.M., the patient was conscious and rational, with normal temperature and no headache. At 12 M., the boy was sleepy but rational with good pulse, normal temperature and no symptoms except the Kernig, retraction and rigidity. Lumbar puncture gave 30 ccm. of fluid. Examination of the fluid as before. Thirty cubic centimeters antiserum given. From this time on there were no symptoms except slight fever, about 100°, which persisted about 10 days. Convalescence was rapid, the boy being now perfectly well.

CASE X. Abraham Rush, four years. Seen Jan. 15, 1908.

Onset yesterday at noon, with sudden fever, headache, vomiting and delirium. When seen at 3 P.M., patient was semi-conscious, responding slightly when spoken to. Neck very rigid and head drawn far back. Pupils dilated, reacting to light. All the limbs were very rigid. The knee jerks were absent and the plantar reflexes were normal. Temperature was 104°.

Lumbar puncture gave 25 ccm. of cloudy fluid, examination of which showed many Gram-negative diplococci, both intra- and extra-cellular. I could not get a growth in culture. Thirty cubic centimeters antiserum given.

Jan. 16. The temperature began to fall soon after the giving of the antiserum and became normal this morning. The child was conscious and rational and did not complain of headache. Rigidity and retraction of the neck were still present, but the rigidity of the limbs had disappeared. I intended to give more antiserum when the temperature should begin to go up again, but from this time on the temperature never rose above 99° and there were no more symptoms. The rigidity and retraction disappeared in the course of a few days and the child made a rapid recovery.

CASE XI. The complete record of this case has not yet been received. Patient was an adult, who had been sick for three weeks when seen on Jan. 23, 1908.

At that time patient was semi-conscious with muttering delirium and picking at the bedclothes. Pupils were dilated and somewhat irregular. There was marked rigidity and tenderness of the neck and some retraction. Kernig's sign present on both sides. Knee jerks not obtained. Plantar reflexes normal. Temperature was about 101°. Lumbar puncture gave 90 ccm. of turbid fluid. Few Gram-negative

diplococci were found, both intra- and extra-cellular. Culture gave no growth. Forty-five cubic centimeters antiserum given.

Jan. 24. There was no notable change in patient's condition except a somewhat lower temperature. Lumbar puncture gave 60 ccm. of slightly turbid fluid. Few organisms were found in the fluid only in the cells. Culture gave no growth. Forty-five cubic centimeters antiserum given.

Jan. 28. Temperature has come down to normal, but there is no notable improvement in the patient's general condition. Lumbar puncture gave about 60 ccm. very slightly turbid fluid, in which no organisms were found. Forty-five cubic centimeters antiserum given.

Feb. 3. Temperature has remained normal, but the patient's condition has remained unchanged. I suspect a condition of internal hydrocephalus has developed. Lumbar puncture gave 40 ccm. of clear fluid. Forty-five cubic centimeters antiserum given. Since this time there has been no further change in patient's condition.

CASE XII. Esther Dickerman, age fourteen weeks. Admitted to the Infants' Hospital, Jan. 27, 1908.

The history was unsatisfactory, but the baby is said to have been sick one week. On admission the temperature was 102°, the sensorium was notably clouded, the fontanelle was bulging and tense, there were slight retraction and marked rigidity of the neck and Kernig's sign was present on both sides. Lumbar puncture gave only one or two fibrinous clots, examination of which showed Gram-negative diplococci, both intra- and extra-cellular. No antiserum was injected into the spinal canal, but 15 ccm. were given subcutaneously.

Jan. 28. There was no notable change in the patient's condition. Lumbar puncture gave no fluid. Fifteen cubic centimeters antiserum were injected into the spinal canal.

Jan. 29. The child seemed generally better and the temperature was somewhat lower. Lumbar puncture gave 15 ccm. turbid fluid. Gram-negative diplococci were found, both intra- and extra-cellular. Fifteen cubic centimeters antiserum given.

Feb. 2. The temperature has been normal until today, when there is a slight rise. Lumbar puncture gave 10 ccm. of slightly cloudy fluid which showed very rare intracellular diplococci. Culture gave no growth. Fifteen cubic centimeters antiserum given. Since Feb. 2, the temperature has been normal and there have been no symptoms.

CASE XIII. Adult, twenty-nine years old. Seen at the Newburyport Hospital, Feb. 6, 1908.

P. I. — One week ago patient began to have very severe headache and fever, which have persisted. Has never been unconscious and there have been no other symptoms. Well developed and nourished. Mind perfectly clear. Marked tenderness and rigidity of the neck with some retraction. Pupils normal. Kernig's sign marked on both sides. Knee jerks not obtained. Plantar reflexes normal. Temperature, 102°. Lumbar puncture was made in several different places. The needle apparently entered the canal, but no fluid was obtained. Thirty cubic centimeters antiserum were injected into the canal.

Feb. 7. There has been no change in condition. Patient was etherized and repeated attempts were made to obtain fluid with lumbar puncture, without success. Thirty cubic centimeters antiserum injected into the canal without unfavorable symptoms.

Feb. 8. Temperature has fallen to 99° and headache is less. There has been no further rise of temperature and the symptoms have gradually lessened and the patient is now reported convalescing.

CASE XIV. Girl, nine years old. Seen in consultation with Dr. Alexander at the New England Hospital for Women and Children, Feb. 7, 1908.

P. I. — Four days ago patient entered the hospital with fever and abdominal pain and was operated on for appendicitis. After recovery from ether, patient was noted to be feverish and still drowsy. To-day the patient was noted to have retraction and rigidity of the neck and lumbar puncture was made and Gram-negative diplococci were found in the cerebrospinal fluid.

P. E. — Temperature, 103°. Patient is conscious, but slightly delirious, complaining of headache. Skin and mucous membranes notably cyanotic. Respiration accelerated. Head somewhat retracted. Neck very rigid and tender to pressure. Kernig's sign marked on both sides. Lumbar puncture gave about

60 ccm. of purulent fluid in which Gram-negative diplococci were found, both intra- and extra-cellular. Thirty cubic centimeters antiserum given.

Feb. 8. Temperature began to fall immediately after the first injection and the general condition is much improved. Mind perfectly clear. Thirty cubic centimeters antiserum given.

Feb. 9-10. Improvement continued steadily. Thirty cubic centimeters antiserum given on each day.

Feb. 13. Child is practically convalescent. Temperature has come down by lysis. All symptoms have disappeared except a rigidity of the neck and the Kernig sign, which are slightly marked.

CASE XV. Jennie, eight years old. Admitted Jan. 13, 1908.

F. H. — Brother of patient two and one-half years old, was taken sick at the same time and died within twenty-four hours.

P. I. — Four days ago patient was taken with headache, vomiting and fever, with marked rigidity of neck. Lumbar puncture gave 50 ccm. of cloudy fluid, in which Gram-negative diplococci were found. Thirty cubic centimeters antiserum given. This was repeated on the three subsequent days, the fluid becoming rapidly clear and organisms disappearing from the fluid. The patient's mental condition became rapidly clearer and symptoms disappeared. She is now suffering from otitis media.

Since the reading of this paper patient has had a relapse.

DISCUSSION OF DR. DUNN'S PAPER.

DR. W. W. McKIBBEN, Worcester, Mass.: I was called, on the evening of Feb. 6, to see a boy of four

years. The parents said he had been sick three days with high fever, intense headache and irritability. He had had initial convulsions. I found a temperature of 105°, moderate opisthotonos and rigidity of the neck extreme; he objected to any noise or light in the room and to being handled. The next day he had the cephalic cry and internal strabismus; he also had stiffness of lower extremities and tache cerebrale. I saw him four days during which time the temperature remained 103° to 105°. I then persuaded them to allow the boy to be sent to the hospital as they would not permit lumbar puncture at home.

On the evening of Feb. 10, with a temperature of 103°, 20 ccm. of Flexner's antiserum was injected by Dr. E. L. Hunt, Asst. Pathologist at Worcester City Hospital. The white count was 18,400. On the morning of Feb. 11, temperature was 105°. At noon another 20 ccm. were injected; temperature in afternoon was 104.5°. On the morning of Feb. 12, temperature dropped to 98.5° and all symptoms except slight irritability disappeared. White count 9,000. Convalescence uninterrupted since that time.

Dr. Wentworth called attention to a paper on the serum treatment of cerebrospinal meningitis by Levy which appeared in a January number of the *Deutsche Medizinische Wochenschrift*. The "Kolle-Wassermann" serum was used. In this paper Levy gives the results of treatment of 55 cases of cerebrospinal meningitis which occurred in the town of Essen from January to November, 1907. Of these 55 cases, 15 were treated outside of the hospital without serum and showed a mortality of 80%; 14 cases were treated in the hospital without serum and showed a mortality of 78+%; in 17 cases the "Kolle-Wassermann" serum was injected intra-spinally and the mortality was 11+%. Levy lays stress on the value of early treatment with serum and the necessity for its intra-spinal injection to obtain good results.

THE USE OF FAT-FREE MILK IN INFANT FEEDING.*

BY CHARLES W. TOWNSEND, M.D., BOSTON.

ANY suggestion which simplifies infant feeding or renders the management of cases of indigestion and diarrhea in infancy easier, is certainly worthy of consideration, for it is a noteworthy fact that the general practitioner is prone to shirk the careful study which a thorough conception of scientific percentage feeding entails.

For a long time the proteids of cow's milk have absorbed much of the attention of specialists in infant feeding, and it has been thought that once these proteids were thoroughly understood and modified to resemble the proteids of mother's milk, the problem of infant feeding would be largely solved.

Curds in infant stools are generally regarded as evidence of faulty proteid digestion, for it is thought that the cheesy part of the milk, instead of being digested, is curdled and passed through the gastrointestinal tract. Whenever these curds occur, efforts are made to modify the proteids which are considered to blame. In working out this problem, whey, which lacks the caseinogen so largely in excess in cow's milk, has been used, as well as

the apparently more scientific but more complicated split proteids, where the same relative proportions of soluble lactalbumen and the caseinogen may be prescribed as in woman's milk.

While the proteids have been studied so carefully, the fat in the mixtures for infant feeding has, on the other hand, often received but scant attention. This, I believe, is due to the fact that the fat of cow's milk and of woman's milk both occur in about the same proportion to the total bulk, and no differences in the quality of the ingredients are recognized by the practitioner as is done in the case of the proteids. The quantity of the fat is borne in mind, but the quality is not considered, yet it has been shown, as by Stern,¹ "that the composition of the fat of cow's milk is greatly at variance with that of the fat of human milk, differing especially in its far greater contents of volatile fatty acids, among which butyric acid is the most important." He finds from six to eight times as much of these fatty acids as in normal woman's milk, a fact that easily explains some, at least, of the relative indigestibility of cow's milk.

Another interesting fact is the one discovered by Luzzatti and Biolchini, namely, that there is a more active fat-splitting ferment in woman's milk than in cow's milk.

Although the value of fat as a food is generally recognized by the practitioner, it is often regarded by him more in the light of a laxative, and, if the bowels are constipated, he is apt to increase the percentage of this ingredient without hesitation. That excess of fat is often a cause of constipation is rarely recognized. Such, however, is the case, and stools are passed that are hard, dry, large, and of a grayish white color, showing a greasy surface when rubbed smooth. In these stools there may or may not be found hard curdy masses like cheese which are often thought to be made up of proteids, but are in reality masses of fat. A diminution in the amount of fat in the food in these cases will often relieve the constipation, much to the surprise of the physician, whose only idea previously had been to increase the fat for its laxative action. Too little fat, it must be borne in mind, will also cause constipation, so that a happy mean for each case is desirable.

Excess of fat is often a cause of gastric disturbance as shown by vomiting, and of intestinal indigestion resulting in the fatty stools just described, and in a failure to gain properly in weight. The infant may even become atrophic. Infants fed with an excess of fat are often ravenous at their meals,—they are starving in the midst of plenty,—and the physician is tempted to increase the strength of the food, especially in fat, with increased trouble for the little patient. Grave general disturbances, as shown by eczema, rachitis, or even convulsions, may occur in infants thus overfed. It must always be borne in mind, however, that what is a proper amount of fat in the food in one case may be excessive and the cause of considerable disturbance in another. No cast-iron rule can be made for infant feeding.

* Read before the Suffolk District Medical Society, Dec. 18, 1907.

¹ Archives of Pediatrics, June, 1905.