facial nerve; and, lastly, the absence of a long, unsightly scar on the cheek.

FIG. 1.

FIG. 2.

FIG. 3.

The woodcuts show—(1) on a healthy jaw-bone, between the dotted lines, the extent of bone removed; (2) the appearance of the patient *before* operation; (3) the right side of the patient's face six weeks *after* operation.

WESTMINSTER HOSPITAL.

CASE OF CHANCERE ON RIGHT SIDE OF PENIS AND BUBO IN LEFT GROIN.

(Under the care of Mr. A. PEARCE GOULD.)

T. M—, aged thirty-two, attended as an out-patient on May 22nd, 1878, with a sore on the penis. It was sharply cut, circular, depressed, and had a greyish-yellow sloughy base. It was seated on the right side of the sheath of the penis; the lowest point of its circumference was three-

quarters of an inch to the right of the lower median line of the penis, and its edge was still further removed from the upper median line. In the left groin there was a suppurating bubo. There were no enlarged glands in the right groin. He had no gonorrhoea, nor any sore on the lower limb. There was no history of strain, nor of injury, nor other ascertainable cause of the bubo. The patient had noticed the sore on the penis for three weeks, and a week after the first appearance of the sore he felt pain in the groin, and discovered a lump there three or four days later, which gradually increased to its present size. The man was admitted into the hospital through the kindness of Mr. Cowell. The bubo was opened, and the chancre treated with iodoform ointment, and in a few days he left convalescent.

Remarks by Mr. GOULD.—There can be no doubt that the cause of the bubo in this case was the absorption of irritant matter from the sore on the penis, for there was an entire absence of any other efficient cause. The interest in the case lies in the fact of a sore distinctly on the *right* side of the penis producing a bubo in the *left* groin, showing a crossing of the lymphatics from the right to the left side, from which it may reasonably be inferred that there was a decussation of these fine channels (those of the left side passing over to the right). Such cases are not rare. While surgical registrar at University College Hospital I saw two such cases in that hospital. One was a man under the care of Mr. Marshall, admitted with a large bubo in the left groin from a soft chancre on the right side of the glans penis. The other was a woman, under the care of Mr. B. Hill, with a soft chancre on the right labium and a bubo in the left groin. In this case a bubo subsequently developed on the right side. And I have met with other cases of this "cross lymphatic infection" among the out-patients, but of these I have no notes. Each of the cases mentioned above had the sore on the right side and the bubo on the left. This is probably accidental, as might be shown by a larger number of cases. When the arrangement of the lymphatics in a fine network is borne in mind, the occasional occurrence of this "cross infection" will not be surprising.

PETERBOROUGH INFIRMARY.

COMPOUND COMMINUTED FRACTURE INTO THE ANKLE-JOINT; IRRIGATION WITH ALCOHOLIC SOLUTION OF COAL TAR; RAPID RECOVERY.

(Under the care of Dr. T. J. WALKER.)

FOR the following notes we are indebted to Mr. C. R. Illingworth, M.B.

Samuel M—, aged twenty-four, of unsteady habits, whilst riding, in a state of intoxication, on the shaft of a trolley loaded with two hundredweight of piping, was thrown upon the ground in consequence of the wheel rolling off the kerb-stone. The left wheel passed over his right foot from heel to ankle. On admission (March 30th, 1878) there was a longitudinal wound two inches in length in front of the ankle, opening into the joint, from which blood was oozing pretty freely. The ligaments in front of the joint were torn across, so that the articular surfaces, bare and comminuted, could be felt through the wound. The fibula was fractured a little above the malleolus, and the tibia was obliquely fractured, the articular surfaces being to a great extent separated and comminuted, although the inner malleolus remained continuous with the shaft and its ligamentous attachments to the bones of the foot were intact. There was no displacement of the foot. Irrigation was employed in the following manner:—The limb having been laid on its outer side, flexed at the knee and kept in position by sandbags, a gutta-percha dish, with a spout for the purpose of carrying off the liquid, was placed beneath the foot, and a tin containing the fluid for irrigation was hung above the limb. The fluid employed was an alcoholic solution of coal-tar, one part of the solution to sixty of water. The "drip" action was secured by several worsted threads tied to a piece of metal at the bottom of the tin, and brought over the edge so as to hang directly over the wound. A piece of linen, wet with the lotion, was laid upon the wound, with its ends cut so as to carry off the lotion into the dish beneath the foot.

On the second day after the injury a dusky redness of the skin, with œdema, extended up to the knee, and the foot was