of bactericidal amboceptors, agglutinins and specific precipitins in the serum. It is now widely believed that acquired immunity depends on the presence of the bactericidal amboceptors in the circulation and body fluids, although

Metchnikoff holds, on the other hand, that it depends to a large degree on an increase phagocytic power on the part of the leucocytes. According to Pfeiffer and Marx, the antibodies are produced in the blood-forming organs. An attack of cholera confers immunity of prolonged duration, although it is not always absolute.

Passive immunity is readily induced in animals by the injection of anticholera serum. As in other instances, it is of short duration, however. Doubtless the same condition may be induced in man. Besredka has proposed mixed active and passive immunization by way of protective inoculation, using killed bacteria which have been saturated with the specific amboceptors.

Serum therapy has been no more successful in cholera than in typhoid fever. The antitoxic serum of Roux and others has had no practical trial. According to Achard and Bensaude, the serum of cholera patients, on the third or fourth day of the disease, agglutinates the cholera vibrio. However, they used the serum in dilutions of 1:20, and in this strength even normal human serum may be agglutinating (Pfeiffer and Kolle, cited by Paltauf). Convalescents even after seven months may show an agglutinating power of from 1/100 to 1/120.

The bacteriologic examination of the stools is the most reliable means of early diagnosis (see above).

Clinical Reports

ERYTHEMA MULTIFORME FOLLOWING VACCINATION.

LOUIS M. WARFIELD, M.D.,

ST. LOUIS.

Considering the great number of persons vaccinated, the comparatively few complicating eruptions that occur is not a little surprising. As has been often remarked, many eruptions occurring in the course of a vaccination period are undoubtedly accidental, while others are undoubtedly due to the vaccine virus.

A condition sufficiently interesting to report came under my notice not long ago.

Patient.—Miss J. H., aged 23, a trained nurse, had never been vaccinated previously, and has always enjoyed exceptional health. Her appetite and digestion are good and menstruation regular. She had had the usual mild children's diseases when young, but had never had any eruptions. For about a year she has been very slightly troubled with constipation. Previous to her vaccination she had been in perfect health and had no gastrointestinal disturbance nor any symptoms of even a mild illness.

Vaccination and Result.—She was vaccinated July 29 on the upper left arm. On August 1, before she noticed anything at the site of the vaccinated area more than a slight itching, she saw irregularly shaped bright red spots on the forearm. These spots soon appeared on the lower legs, and shortly both arms from shoulder to wrist, and both legs from thigh to ankle, were thickly covered. These spots soon became firm, nodular and painful. There was no itching or burning. She felt sore all over. There were pains in the knee joints, and wherever the lesions were close to the bones they were very painful to the touch.

Examination.—August 4. She is a well-developed, well-nourished woman. The mucous membranes were a good color, tongue somewhat coated, pulse slightly accelerated, temperature 100 F. Over the extensor surfaces of the arms and legs there was an eruption consisting of rose-red irregularly-sized and shaped spots, level with the surface of the skin, bright red, somewhat infiltrated areas, varying in size from that of dimes to five cent pieces, and firm, deeply seated nodules,

perceptibly raised above the surface of the skin. There were no vesicles, nor was there umbilication of the nodules. The nodules situated over bony prominences were quite firm and exquisitely painful to the touch. Some were of a typical violaceous hue and appeared shiny. There was no definite line of demarcation seen between the lesions and the surrounding skin.

Subsequent History.—New crops came out, following the order of ripening described from day to day, until August 8. In the meantime, the older lesions grew softer, less painful and more purplish in color. On August 8 vesicles appeared at the site of the vaccination. These ran a normal course. On August 10, three small vesicles appeared on the right forearm; by the next day these had disappeared, leaving small red spots with slightly umbilicated centers. By August 14 the rash had entirely disappeared and only a few brownish stains were visible. During the period of the eruption there was slight fever but not more than could be attributed to the effects of the vaccine itself. The only inconvenience was the general soreness, especially in the knees and elbows.

Remarks.—That this was a typical case of erythema multiforme is certain. The whole course of the eruption was short, but the lesions and their appearance in crops were typical. Most curious were the great tenderness of the nodules and the feeling of soreness over the arms and legs, especially around the knee and elbow joints. There were no lesions on the backs of the hands or dorsal parts of the feet. As usual with erythema multiforme, the eruption showed itself before the vesiculation of the vaccinated area.

LAPAROTOMY UNDER LOCAL ANESTHESIA ON A PATIENT, AGED SEVENTY-TWO.

R. L. IGEL, M.D.

LEAVENWORTH, KAN.

I wish to report a very interesting case that came under my care July 31, 1905.

Patient.—A woman, aged 72, was suffering from an ovarian cyst of seven years' standing, which had grown to such an extent as to fill the whole abdominal cavity.

Examination.—I found her in terrible pain, resembling that of obstruction of the bowels. An enema was given, with no results, and I knew that cathartics would only increase the trouble. I therefore determined to relieve the bowel that I was sure was being obstructed from the pressure of the cyst by tapping the tumor.

Operation.—She was taken to the hospital, a few blocks from her home and was prepared for operation. Chloroform and ether were contraindicated by her age and exhausted condition, so I began by injecting a small amount of a 4 per cent. solution of cocain three inches below the umbilicus, in the median line, just under the skin, and followed with my knife, making a clean incision. The adipose tissue was then gone through and separated until the peritoneum was reached, and this tissue was cocainized in like manner in front of the knife. This incision brought the cyst into plain view. I then tapped it with a small needle, to allow shrinkage enough to grasp the walls with forceps, after which a trocar was inserted and the liquid, which measured four gallons, was allowed to run off slowly. I made a complete exploration of the abdomen and of the adhesions and withdrew as much as possible of the cyst from the walls of the abdomen, so as not to allow of any infection, washed the sac out with normal salt solution, replaced it and allowed one-half pint of the solution to remain. I also filled the abdomen with normal salt solution and did as thoroughly aseptic an operation as possible. The peritoneum was closed with four stitches and the abdominal wall with two stitches. The patient talked to me in the meantime and turned herself on one side to help me drain the cyst. Her pulse and heart were constantly watched and she was allowed to take continuous sips of cool water.

Result.—She was removed from the table to a ward. She did not complain during the operation, nor since. The bowels moved freely of themselves. That night she took two cups