worked too hard or eaten too much, as he often has, and goes to bed. If the drowsiness happens to be associated with pain in the eyes, or headache, an oculist will probably be consulted, and it will disappear after the eye defect has been corrected.

I was consulted a few years ago by a minister whose only symptom of eye-strain was extreme drowsiness on attempting to write his sermons. There was no pain in the eye or head, and he could see well enough in a good light. He was forty-five years of age, and had come to the conclusion that the trouble resulted from an effort in using the eyes, which proved to be the case. Another patient, a bookkeeper, who suffered still greater annoyance from this symptom, was seen last year. There was some headache and asthenopia, but his especial complaint was of feeling tired and sleepy most of the time. He often found it almost impossible, even during the morning hours, to keep awake when at work on his books. He had taken a vacation, but the trouble began again when he returned to his work. An examination showed a considerable degree of astigmatism, and correcting glasses were prescribed for constant use. He reported three or four weeks later that the drowsiness had almost entirely disappeared and that he was feeling much better generally. These two cases are not exceptional, and I have reported them simply to illustrate the degree of annoyance that may sometimes arise from this symptom.

Every practitioner is familiar with the patient who complains of feeling mean and played out most of the time, but has nothing especial the matter with him. He gets up feeling tired, and is glad when night comes. He sometimes improves under medical treatment, but is often overworked and needs a rest. Most of these patients get better, and an occasional bracing up keeps them in running order. Some of them feel well only occasionally; they go through life tired, and seem intended for a life of leisure rather than for hard work. Others break down and have "nervous prostration." All this misery cannot, of course, be attributed to eye-strain, but it is undoubtedly a fact (and one becoming more generally recognized) that many of these cases gain faster, and are less liable to "break down" in the future, if an existing eye-strain is relieved. With the class that may be called *eye-laborers*—students, bookkeepers, sewing women and the like—the ciliary is probably the hardest worked muscle in the body, even if the eye is normal. If, in addition, this muscle is called upon to correct hypermetropia or astigmatism, it is not surprising that the added strain is often the last straw in breaking down the overworked man or woman.

PROOF-READERS' MISTAKES—"Among the sickness of the plague in India are a colony of monkeys, near Hurdwar, and the authorities are now busily engaged in trapping and isolating deceased animals."—*Medical Press and Circular*.

Trapping would seem to be an unnecessarily difficult method of catching "deceased" animals. Under the heading, "A Bloodthirsty Physician," the same journal notes the following misprint: "Our attention is called to an advertisement of Aseptolin-Edson, which is appearing in the medical journals, containing the startling statement: 'Out of a number of cases of consumption treated therewith, only one proved unsatisfactory. Aseptolin has far surpassed our most sanguinary expectations.'"

Clinical Department.

TWO CASES OF ABSCESS OF THE LIVER.¹

BY HENRY JACKSON, M.D.,
Assistant Visiting Physician, Boston City Hospital.

CASE 1. Geo. McK., twenty-one, single, suffering with multiple small abscesses in the liver, entered my service on July 24th. The previous history was negative except that three years before his entrance to the hospital he was confined to bed by an acute appendicitis. At this time he was in Providence; according to the statement of the patient there was a tumor in the region of the appendix, with severe abdominal pain. No operation was performed. The pain lasted about one week, but the patient was confined to his bed for four weeks. Since then his health has been good.

For two weeks before entrance to the hospital his appetite was poor; he had headache and at times vertigo; there was also some palpitation and distress in the region of the heart. He continued at his work as a surveyor on the Boston & Albany Railroad at Newton until four days before entrance to the hospital; then after a hard day's work he had severe pain in the right iliac region which caused him to stoop forward when standing; nausea and vomiting soon followed. Two days later there was a chill followed by sweating, severe headache and delirium; the next day several chills, fever, vomiting and a good deal of pain in the right iliac region. The bowels were constipated.

Physical Examination.—Expression is somewhat anxious. Mind perfectly clear. Some tenderness in the back of the neck, a symptom noted as cerebro-spinal meningitis was considered as possible in that at the time there were several cases of this disease in the hospital. No enlargement of the heart; at the base a soft musical murmur heard over the pulmonic area. Lungs, liver, spleen, not abnormal. There was general abdominal tenderness, most marked in the right iliac region, where gurgling was felt; no rose-spots.

Examination of the blood showed no plasmodia malariae. Leucocytes, 7,800.

July 26th. No change in general condition. A positive Widal reaction was reported. I consider that this was probably an error due to some fault in the technique, as several subsequent examinations gave a negative result.

July 27th. Leucocytes, 8,000. Quinine was given, 20 grains after the temperature fell a degree and a half. The quinine produced no effect upon the fever or the chills.

July 30th. Mental condition bright. No complaint was made of abdominal tenderness. The spleen was found to be somewhat enlarged. No enlargement of the heart. There was some tenderness in the right side of the abdomen, but there was not sharply localized tenderness at McBurney's point. There was no rigidity of the muscles of the right side of the abdomen, and the legs were fully extended without causing pain. Severe chills recurred daily towards five o'clock and on the 27th, 29th, 30th there was also a chill in the morning. Leucocytes, 16,800.

July 31st. Leucocytes, 23,000. Slight jaundice appeared to-day.

August 1st. Leucocytes, 33,000.

Up to August 5th there was a gradual loss of flesh

¹ Read at the Clinical Section of the Suffolk District Medical Society November 17, 1897.

and strength. The jaundice became more marked, and the liver extended from the 20th interspace to just below the ribs. There was slight dulness in the lower part of the right back.

After consultation with Dr. Munro it was decided to aspirate the liver, on the supposition that we had to deal with an abscess of the liver. He was etherized, and three punctures made in the region of the liver anteriorly, and one posteriorly. All were negative.

From August 5th to August 11th there were no chills, and his general condition improved a little.

August 12th. Leucocytes, 19,000.

August 13th. Again a chill. I saw him on the surgical side, and suggested as a possible diagnosis malignant endocarditis, considering this diagnosis probable in view of the negative result of the puncture made, and because a new murmur had appeared in the mitral area. The abdomen was a little tympanitic, and there was some tenderness in the region of the appendix.

August 15th. There was a violent chill at 10.30 P. M., his temperature rising to 109.5°. His death occurred three hours later. Up to the last his mental condition was bright, modified only by mild delirium at the time of the intense fever which followed the chill.

In arriving at a diagnosis we considered typhoid fever, appendicitis, malignant endocarditis, general tuberculosis, pyemia from some source not determined, and abscess of the liver.

Autopsy. Two hundred c. c. brownish fluid in the peritoneal cavity. Old adhesions between the great omentum and the right iliac wall. The appendix and cecum were bound down by old adhesions. The interior surface of the appendix was roughened and the walls thickened. Heart and lungs not abnormal. Spleen, 475 gm. Liver enlarged. On the anterior surface many small white patches, especially on the left lobe; on the under surface multiple abscesses. On section many abscesses found, especially in the left lobe. The portal vein was occluded by yellow adhesive thrombi and pus.

Anatomical Diagnosis.—Chronic appendicitis and chronic peri-appendicitis, multiple abscess of the liver, abscess of the mesenteric glands, acute splenic tumor.

The origin of the abscesses in the liver was obscure,

as there was found aside from the acute suppurative process in the liver only the chronic appendicitis and the acute inflammation of the mesenteric glands. The duration of the disease was seven weeks, four before entrance to the hospital and three in the hospital; so it is possible that the primary trouble may have been an exacerbation of the chronic appendicitis, especially as there was found in the appendix a little inspissated pus, though there was at the time of the autopsy no evidence of a recent acute inflammation.

CASE II. The history of this case is very incomplete and I report it only because of the interest attaching to the rare disease found, namely, amebic abscess of the liver.

On July 28th an old man was brought to the hospital in a state of marked collapse. He had walked into the Boston Dispensary, and was at once sent to the hospital by ambulance. He was pale, covered with a cold sweat; pulse small, intermittent and irregular. There was abdominal pain, and a history of obstinate constipation for eight days and no movement for five days. It was considered indiscreet to operate at once, and he was placed in bed and stimulants given.

I saw him in consultation with Dr. Munro the next day. In the right hypochondrium was a large, tense, hard mass, extending about four inches below the edge of the ribs. The liver dulness began in the fourth interspace. Over the lower right front a friction sound was heard. The diagnosis appeared to be between an enlarged liver and subphrenic abscess with depression of the liver.

At the autopsy there was found an enormous amebic abscess of the liver; amebæ were found. In the upper part of the colon were two chronic ulcers.

There was during life no history of a preceding dysentery. His wife said he had had good health up to the last four years, though he had been subject to attacks of obstipation.

CANCEROUS STENOSIS OF THE PYLORUS.¹

BY G. LIEBMANN, M.D.

THE case I wish to report is that of a man forty-three years of age who had always been well up to the time of the advent of his fatal sickness, a little over a year ago; a man who did not offer symptoms of emaciation—he was thin before—but he had not a cachectic appearance at all. His appetite was good from the start of his sickness until he died. That was one of the most peculiar phases in this case, that his appetite was good from the start. He was taken about one year ago with what appeared to be paroxysms of spasmodic pains or cramps in the epigastrium and in the right hypochondrium. At first it was thought the paroxysms indicated gall-stones. The liver was enlarged and the region of the gall-bladder somewhat tender. These paroxysms soon lapsed into continuous suffering so that the man could not sleep and could not enjoy life at all unless under an anodyne; a small quantity of an anodyne would relieve him. His appetite, as stated, was good; tongue was clean; and the examination of the contents of the stomach showed that there was hydrochloric acid present in more or less normal quantities. The total acidity was at first 90, later 44. The pain complained

¹ Read at the Clinical Section of the Suffolk District Medical Society, November 17, 1894.