of tiny beasts. These sometimes staid still in one place and sometimes ran with inconceivable swiftness and hissing noises about his room from one corner to the other. If these disappeared after a time, almost immediately there appeared in their places naked women, and he thought it strange that the ladies of his family should regard them with any calmness. His memory for recent events had almost entirely disappeared, and his previously unusual intellect was brought down to a very low level. Being informed of the entirely subjective nature of his visions he began to observe them with interest, but still with a certain doubt if they were not, after all, real. Thus, besides the ordinary symptoms of a cerebral apoplexy in the optical territory, and apart from the frequent hallucinations, he had three unusual symptoms, cyanopia, macropsia, and metamorphopsia. All of these are so rare that it is the more extraordinary that they should all have occurred in the same patient. The variety of the symptoms is also remarkable: Paralysis, contracture, tremor, aphasia, agraphia, lessening of sight and hearing, intellectual decay, partial loss of memory, hallucinations of sight, hearing, and smell, and finally, besides all these, cyanopia, macropsia, and metamorphopsia.

MITCHELL.

76. AN ACUTE CASE OF LANDRY'S PARALYSIS. Henry H. Haward (*British Med. Jour.*, 1, 1898, p. 1654).

A man of fifty-four years had been for a week exposed to the weather more than usual. After having felt a tingling in the hands for two hours, he walked a mile to his home, felt disinclined for food, went to bed, could not sleep and three hours later found that food would not go down. After this he rapidly lost strength so that in twelve hours he was quite helpless, respiration was hurried and stridulous and both deglutition and expectoration were impossible. Deep and superficial reflexes were present and although he had no pain, tingling and numbness were very distressing. Anesthesia was absent throughout. The muscles were affected symmetrically, and approximately in the following order: First, the pharyngeal, laryngeal, spinal and cervical; then those of the hips, thighs, shoulders and arms; next, the lower intercostal and abdominal; followed by those of forearms, legs, hands, feet, and upper ribs; lastly, facial, lingual, labial, and diaphragmatic muscles. The patient died forty-three hours after the appearance of the first symptom.

PATRICK.

77. APHASIA IN RELATION TO TESTAMENTARY CAPACITY. (*British Medical Journal*, September 3d, 1898).

In a discussion on this subject at the last meeting of the British Medical Association, Gairdner said that no general or positive rules could be laid down. It was not a general question of law; not a thing that could be put into legal or physiological categories. It was a question of detail in each individual case and, we gather, to be decided pretty much without direct reference to the aphasia. Does the testator know what he wants? Does he form a clear conception of what he wants and does he succeed in giving effect to that conception in the particular document that is the "will?" Or is the document not really his will at all, but unduly inspired by others? These are the questions of detail to be decided.

William Elder in a much more systematic consideration of the