The Journal of the American Medical Association

Published under the Auspices of the Board of Trustees.

VOL. XLIX.

CHICAGO, ILLINOIS, SEPTEMBER 14, 1907.

No. 11.

Original Articles

AN EPIDEMIC PNEUMOCOCCIC CATARRHAL DISEASE.*

CLINICAL STUDY.

HARVEY G. BECK, M.D.

BACTERIOLOGIC STUDY.

WILLIAM ROYAL STOKES, M. D.
BALTIMORE.

CLINICAL STUDY.

INTRODUCTORY.

It may appear presumptuous to insert the word "pneumococcic" in the title of a paper which has to deal largely with a disease or condition with local manifestations involving the mucous membrane of the upper air passages and conjunctivæ, because recent investigations tend to show that these anatomic situations are a normal habitat of the pneumococcus. It is, therefore, with a considerable degree of reserve that this title has been adopted, even after coupling a definite clinical picture with the persistent presence of an overwhelming number of pneumococci characterizing this series of cases. It is, of course, impossible to make a numerical division as to what constitutes a normal or an abnormal invasion of pneumococci in the mucous membrane of the eye, nose and throat. Therefore, we do not regard the increase in the number of these organisms, even though it be enormous, of sufficient importance to warrant our conclusions, but when this is associated with a definite underlying pathologic basis and distinct clinical features we feel justified in reporting this series of cases under this caption.

REVIEW OF LITERATURE.

In a cursory review of the literature we find that epidemic pneumococcic conjunctivitis has been described in a special article by Axenfeld.¹ We also find a description of pneumococcic fibrinous pharyngitis by Bissell, and pneumococcic pseudomembranous bronchitis by Landrieux and Triboulet.² If the conditions these observers described correspond with our present epidemic then they failed to comprehend the entire clinical picture and merely presented single aspects of the disease.

Davis of Chicago³ intimates that acute catarrhal conditions resembling grippe might possibly be caused by the pneumococcus, but does not furnish clinical and bacteriologic proof of the assumption.

Samuel J. Darling, pathologist, Ancon Hospital, Pan-

ama,⁴ examined the accessory nasal sinuses in thirty-seven autopsies in which death was due to pneumococcus infection. Ninety-two per cent. of the cadavers showed involvement of one or more of these sinuses. Thirteen cases of pneumococcus meningitis were studied. All of these cadavers presented an inflammation of the sinuses, and in every one the middle ear and mastoid were normal. The inflammation was usually intense, fibrinopurulent in character, and apparently antedated the general infection. In all of his cases the pneumococci were found in abundance. He also found a pneumococcus sinusitis in 28 per cent. of the control cases in none of which death was due to pneumococcus infection.

Darling believes that the portal of entry of the pneumococcus is in most instances an accessory nasal sinus. the mucous membrane in the sinus becoming infected through a previous rhinitis.

I mention these investigations of Darling's because his observations and the study of our cases establish the fact that there is undoubtedly a direct line of invasion by the pneumococcus beginning in the upper air passages, frequently involving the accessory sinuses which may ultimately lead to the development of pneumonia, pneumococcus meningitis, endocarditis, septicemia, etc.

STUDY OF EPIDEMIC.

Two years ago this spring there occurred in my practice in Baltimore a curious catarrhal condition which I regarded at the time as epidemic in nature. It differed essentially from ordinary colds in that it was apparently not the result of exposure and it affected almost invariably more than one member in a family and sometimes entire families. Last year no such case came under my observation.

In March of this year my attention was again directed to the same condition. My interest was aroused, after being summoned within one week to three families, in which either the entire family or several members of a family were similarly affected. Since then many other cases have come under my care which have enabled me to make a careful study of the affection from a clinical standpoint, and have furnished Dr. Stokes, who kindly offered to make a bacteriologic investigation, an opportunity to study the epidemic from an etiologic standpoint. In this work Dr. Stokes was assisted by A. E. Winlack and D. G. Preston, students at the College of Physicians and Surgeons.

As the following notes will show, these cases may be looked on clinically as measles without a rash, as conjunctivitis with general catarrhal symptoms, as whooping cough without a whoop or as grippe without mental or physical depression. Since this group of cases presents a definite clinical picture—a symptom-complex not exactly corresponding to the familiar epidemic diseases—it appears to me not at all improbable that the epidemic is an affection having a distinct entity.

* Read in the Section on Pathology and Physiology of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

* From the Department of Medicine and Pathology of the College of Physicians and Surgeons, Baltimore, and the Bacteriologic Laboratory of the State and City Boards of Health.

1. Kölle and Wassermann: Bacteriology.

2. Rev. gen. de clin. et de therap., Paris, 1896, x, 97.

3. THE JOURNAL A. M. A., May 11, 1907.

4. THE JOURNAL A. M. A., Nov. 10, 1906.

This report is based on the clinical study of fifty-six cases, of which thirty-three were examined bacteriologically. I shall give the histories in full of a few of the most typical family groups of cases showing house infection, one case of the severest type of the disease, and follow these by a summary of the entire series:

GROUP 1.—My first experience was with the G. family, consisting of six members, all of whom were affected; father aged 42, mother 34, Pearl 18, Urban 6, Alfred 5, and baby 10 months. March 13, 1907, I was called in to see the baby, who was thought to be suffering from whooping-cough. The child had had none of the acute infectious diseases. Alfred and Urban had had whooping-cough one year ago. Alfred

and night sweats; this was associated with paroxysmal cough, discharge from the nose, watering of the eyes, slight soreness in the throat, slight pain in the chest and abdomen, and absolute loss of appetite. There was marked constipation. Vomiting frequently occurred after paroxysms of coughing. The cough was of a severe croupy character, and, at first, lasted half an hour. These attacks occurred more frequently during the night. There were four or more severe attacks at night and two or three lighter ones during the day. Occasionally during the night, these paroxysms terminated in a distinct whoop. When the child was first seen, the symptoms had subsided, except the cough and the discharge from the nose, which was at this time mucopurulent. He had a slight bronchitis and vesicles of chickenpox.

Urban, who was similarly affected, was also taken with chills and fever, which were followed by sweats. He had a peculiar dry paroxysmal cough, having three or four severe attacks at night, which lasted until he vomited. He had marked lachrymation without congestion, nasal discharge, sore throat, thoracic and abdominal pains, but no rash. The fever lasted about one week, during which he had several night sweats; anorexia was marked and he was very constipated.

Pearl's case corresponded closely to Alfred's and Urban's, except in the following particulars: Her eyes, at first watery, became mucopurulent and the lids were pasted together in the morning. In addition to the discharge from the nose, there was much sneezing. The cough, which was spasmodic

Fig. 1.—Dr. Friedenwald's patient, showing purulent conjunctivitis and swelling of upper eyelids.

Fig. 2.—Peritoneal exudate from mouse inoculated with fibrinous exudate from throat, showing many typical pneumococci with capsules. The photomicrographs were all made by Dr. T. M. Wright, of the City Health Department.

Fig. 3.—Peritoneal exudate from mouse inoculated with culture from throat of typical case, showing pneumococci, capsules and one short chain.

in character, produced gagging after each paroxysm. Her appetite was not impaired.

The mother had the same train of symptoms; namely, fever, sweat, cough, sneezing, rhinitis, mild conjunctivitis, pain in chest, slight abdominal cramps, anorexia, and vomiting after paroxysms of coughing. Her symptoms lasted three weeks.

The father had the same symptoms, but a more intense conjunctivitis. His eyes were painful, watery and itching at the beginning; the conjunctivæ soon became injected and the eyes felt dull and heavy; photophobia was quite marked. There was slight frontal headache. The discharge from the eyes, at first serous, became mucopurulent. The eye symptoms subsided in four or five days. The respiratory and gastrointestinal symptoms were less pronounced than in other members of the family. He had a sore throat, a moderately dry cough, not paroxysmal in character, no nausea, vomiting or anorexia, and no thoracic or abdominal pains.

The baby began with intestinal symptoms: viz., nausea, vomiting and diarrhea. The cough appeared two days later, accompanied by fever, sweats, conjunctivitis, rhinitis, pharyngitis and bronchitis. The cough, which was severe, appeared in paroxysms, simulating fully developed whooping-cough from the beginning. She had a mucopurulent conjunctivitis in both eyes, slight swelling of the eyelids and intense photo-

was the first one in the family affected with this catarrhal condition. The others, except Urban, were affected three days later. Alfred did not remember having played with children similarly affected, before he was taken sick. Later, two children in the M. family, living in the same block, developed similar symptoms. Alfred had been playing with these children during his illness.

The mother gave the following description of Alfred's illness: He began to complain of chills followed by slight fever

phobia, the child burying her face in her mother's lap to shut out the light. A rhinitis, which began with sneezing, followed by serous discharge, became mucopurulent in character. The throat was uniformly red; the tonsils were not enlarged. The mother felt alarmed because the child refused all nourishment for several days. It vomited considerable mucus and had diarrhea, passing, in twenty-four hours four to seven stools which contained much mucus and were highly offensive. The temperature was 102 F.; the pulse was rapid. Coarse bronchial râles were heard all over the chest. The liver and spleen were not enlarged. A slight heat rash was observed. Cover-slip preparations and cultures from the nose and the throat of the mother and the baby showed pneumococci.

GROUP 2.—May 10.—S. family. Four members were affected: Aunt K., aged 35, mother aged 40, and two children. Celeste aged 5 and Janette aged 7, both of whom had had measles and chickenpox. Mrs. M. and sister, living in a flat on the second floor of the same dwelling, were also affected. The first one affected was Mrs. M., who became ill about April 20. A week later aunt K. was affected; Janette three days later; Celeste three days after Janette, and the mother a few days later than Celeste.

Aunt K.'s illness began with chilliness and fever followed by sweats, associated with cough, sneezing, discharge from nose and lachrymation. Later, the nasal discharge became mucopurulent, but that from the eyes remained watery. The

remove them. There was complete anorexia. The bowels were constipated. The child suffered from headache, earache and abdominal pains and also had considerable mediastinal pain. She complained of dryness of the mouth, great thirst and hoarseness.

Physical examination of Janette.—The patient was up and about, but appeared somewhat nervous. Her eyelids were swollen, the upper more than the lower, and were of the peculiar rose color described by Axenfeld. There was a mild blepharitis and the eyelids were matted with dry, yellow masses of pus. The child presented a picture of adenoid facies and was breathing through her mouth. A profuse herpetic eruption was observed on the lips. There was a slight excoriation about the nares. The tongue was dry and slightly coated and showed several patches of aphthous stomatitis,

Fig. 4.—Many pneumococci in typical exudate from throat.

eyes became painful. She suffered with burning and itching and was not able to read without considerable difficulty. Epistaxis occurred three times during the attack. Her throat was sore and she complained of dryness and hoarseness. There was substernal pain. The cough, which was dry, occurred in paroxysms, and occasionally there were associated sensations of choking, as if a foreign particle were lodged in her trachea. This was sometimes accompanied by vomiting of intensely tenacious mucus. There was also loss of appetite, which lasted two days.

Janette was taken sick three days after aunt K. Her illness began with a distinct chill, which was followed by considerable fever and sweats, and by sneezing, nasal discharge, cough and mucopurulent conjunctivitis. The right eye was first affected; the left shortly after. The discharge from the nose, which was watery at first, became so thick and tenacious that it clogged the nostrils and the child was unable to breathe through her nose, especially in the morning. In order to relieve this obstruction, the mother resorted to the use of cold cream inserted into the nostrils, after which she was able to remove plugs of inspissated secretions by means of a hairpin. This procedure was carried out for five successive mornings. The cough, croupy in character, was worse at night than during the day and occurred in paroxysms. The tenacious secretions caused a choking sensation and vomiting, when an effort on the part of the child was made to

Fig. 5.—Pneumococci in a pus cell and in the nucleus of an epithelial cell from a case with a fibrinous membrane.

Fig. 6.—Many pneumococci within the protoplasm of a pus cell showing active phagocytosis.

which also appeared on the buccal mucous membrane. The mucous membrane of the mouth and pharynx were uniformly red. The tonsils were slightly enlarged; the uvula was swollen and edematous. The cervical glands were prominent. Coarse, dry râles were heard over the chest. Pulse and temperature were normal.

Celeste began to complain three days after Janette. She had played with Janette and had spent much time with the family, similarly affected, living in the upper flat. The subjective symptoms corresponded throughout with Janette's.

Physical examination of Celeste.—The child was sitting up, but not disposed to play. Pulse, accelerated; evening temperature, 101.5 F.; skin, dry. The child breathed through her

mouth. The eye symptoms were not as marked as Janette's, but were worse a few days before. She also had herpes and her tonsils were considerably enlarged. The mucous membrane was of a deep red color. While her throat was being examined, she was seized with a paroxysm of coughing, during which she gagged and expectorated a large amount of thick, tenacious, fibrinopurulent matter, portions of which were rusty in character, not unlike the sputum of pneumonia. This expectoration adhered to the bottom of the vessel containing it when inverted. Microscopic examination showed numerous pneumococci. The cervical glands were enlarged and painful. Coarse bronchial râles were heard all over the chest, otherwise the chest was negative. A blood examination showed a leucocytosis of 15,200, and the following differential count: Small mononuclear cells 20, large mononuclear 10.5, polymorphonuclear 69.5. Cultures and cover-slip preparations from the mucopurulent expectoration after vomiting, showed pure pneumococci.

The mother had a very mild attack; the most distressing symptom was the cough.

The disease ran a course of about two weeks in the case of the aunt and the two children. In the case of the mother it lasted only a few days.

One of the most interesting cases of this collection was treated in the Baltimore Eye, Ear and Throat Char-

Fig. 7.—Pus from eye showing phagocytosis.

ity Hospital. It was through the courtesy of Dr. Harry Friedenwald that I was invited to examine the patient and permitted to embody the history in my report. I am indebted to Dr. J. W. Downey, resident physician, for the history of the case, from which I have made the following abstract:

CASE 1.—The patient, a boy, aged 18 months, was admitted to the hospital April 25. The mother thought the child had measles a month previously, although she says the measles never "came out." The early symptoms of the illness were hoarseness, cough, fever and lassitude. Several brothers and sisters were affected in the same way; none of them had a rash. The eye condition had existed one week previous to the admission to the hospital, and the eyes had been slightly affected during the whole course of the so-called "measles."

Examination.—This showed a fairly well developed child, although rather poorly nourished. The lids of both eyes were swollen, so that the eyes were closed (Fig. 1). The lids were soft to the touch. There was a slight mucopurulent discharge from both eyes and a slight eczematous eruption around the eyelids. The palpebral conjunctivæ of both eyes were much swollen, and covered with a slight, loosely adherent membrane. The ocular conjunctivæ were not covered with a

membrane and were not edematous, but were very much congested. Examination of conjunctivæ caused bleeding. There was a cloudiness of the cornea of the right eye and a hazy appearance of the cornea of the left eye, but there were no ulcers. Impetigo pustules were observed on the face and body. There was a pale, stringy, nasal secretion. The voice was weak and hoarse, and his cries were crowing in character. The mucous membrane of the mouth and throat was slightly reddened, and there was a slight, dirty looking membrane, which appeared in patches, in the pharynx. The larynx also showed a few shreds of membrane. Cough was a distressing symptom. Moist bronchial râles were heard. The chest was normally developed and the heart was normal. Appetite was poor. The child passed from four to five diarrhetic stools daily; these were at times tinged with blood. The spleen was distinctly enlarged.

One thousand units of diphtheria antitoxin were at once administered, although cover smears and cultures on blood serum and agar later proved negative for diphtheria. The bacteriologic examination of the eyes showed almost pure pneumococci. A few staphylococci were also found, and staphylococci were obtained in cultures from pustules from the face and leg.

May 2, an ulcer developed on the right cornea. The left cornea was still cloudy, but no ulceration appeared. No improvement in general condition.

May 7, ulcer appeared on the left cornea.

May 17, the report showed that the child had had an almost continuous fever for three weeks previous, ranging from 99 to 100.3. The eyelids were still swollen and the child was unable to open them. The intestinal symptoms had improved and the membrane from the eyes and throat had disappeared.

May 25, Dr. Friedenwald reported the recovery of this patient with maculæ of both corneæ.

CASE 2.—A parallel case is that of Mr. S., aged 23, who had a severe purulent conjunctivitis with small ulcerations on each cornea and a yellow, loosely attached membrane on the left inferior turbinate and on the left side of the nasopharynx. The general appearance of the membrane was very suggestive of diphtheria and 3,000 units of antitoxin were at once administered. Cultural and inoculation methods of examination, however, demonstrated a pure pneumococcic membrane.

I shall not give a detailed statement of other cases which resemble greatly those above reported. In the following I shall give a brief summary, based on the observations made on my own cases and those of several colleagues.

ETIOLOGY.

That this disease is of an infectious nature is evident from the report of the two family groups in this paper. In all except ten cases there was a history of house infection. Forty-six cases occurred in thirteen families. In a family consisting of eight members, only two girls were affected. They slept in the same bed. Another family, in which six were affected, the disease was distinctly traced to a servant who had the symptoms for three days before the general outbreak. An interesting example of the epidemic character is an outbreak on board a bay steamer, in which 20 per cent. of the crew suffered the same symptoms, including the captain, chief engineer, second mate and watchman. A smear from the engineer's throat showed epithelial cells containing as many as sixty pairs of encapsulated diplococci. The epidemic character of pneumococcus conjunctivitis has been pointed out by Hauenschild⁵ and its invasion of families by Bänzinger and Silberschmidt and by Veasey.⁶ Most of the cases of pneumococcus conjunctivitis which have been reported have not occurred in epidemics.

5. Ztschr. f. Augenheilk., III, p. 200.

6. Arch. of Ophth., xxviii, 1899, p. 30.

From a single observation, it appears that animals are not immune. In a family in which three members were affected, one of the patients was fondling a pet cat. Three days later the animal began to sneeze, which was followed shortly by a severe cough and a purulent conjunctivitis. Unfortunately, the cultures taken from the eye were lost and no bacteriologic examination was made. Collica⁷ was able to produce pneumococcus conjunctivitis experimentally in young dogs, but not in rabbits.

AGE.

The disease is met with more frequently in children.⁸ In twenty-nine of the fifty-six cases, it occurred under 7 years of age, ten of these were under 1 year, the youngest patients were twins, 8 weeks old. It occurred four times in the second decade, eight in the third, nine in the fourth, and five in the fifth; the oldest patient was the engineer above referred to, 56 years of age.

PREVALENCE.

All of my cases occurred in the spring. Epidemics of pneumococcus conjunctivitis have likewise been observed in the spring or in the fall.⁹ Many new cases have been brought to my notice since I concluded the study of this series. A number of physicians have informed me of meeting similar cases in their practice in Baltimore. Physicians in other cities have also seen numerous cases which correspond to this disease. No reference has been made in any of the medical journals, although the lay press recently reported epidemics of "colds" in London, Philadelphia and Baltimore. These epidemics may possibly be of the same nature.

SYMPTOMATOLOGY.

The period of incubation ranges from two to seven days.¹⁰ The invasion is usually marked by chilliness, slight febrile disturbance, and occasionally night sweats. There is rarely a distinct chill. The temperature, usually from 99.5 F. to 102 F., continues from three or four days to one week. The highest temperature recorded is 103.5 F.; the pulse is not much accelerated. Sneezing, lachrymation and a mucous discharge from the nose are early symptoms. These are followed by a burning sensation in the nose, itching of the eyelids and slight soreness in the throat.

A spasmodic cough, croupy in character, usually worse at night than during the day and often associated with nausea, and even vomiting, is a fairly constant phenomenon. Occasionally the paroxysm terminates in a distinct whoop. In some instances the cough is less spasmodic and more hacking, like that of early tuberculosis. It is nearly always a dry cough, and, if expectoration is present, it occurs as an extremely tenacious, yellow mucopurulent excretion, which adheres to the bottom of a vessel when inverted, and rarely is it blood streaked. There is frequently an associated bronchitis, but the symptoms manifest a more profound inflammation of the upper air passages as indicated by a uniform redness of the mucous membrane of the nose, naso-

pharynx, soft palate, larynx and vocal cords and by hoarseness and a laryngeal or tracheal cough.¹¹

The eyes are also frequently affected; the usual symptoms are lachrymation and photophobia; the conjunctivæ often become injected, and in twenty-one of my cases (37.5 per cent.) a purulent conjunctivitis developed. Two of these cases were complicated with corneal ulcers.

Profuse herpetic eruptions frequently occur. The tongue is generally dry and somewhat coated; loss of appetite and often complete anorexia are distressing symptoms, especially in nursing children; nausea and vomiting of stringy, tenacious mucus, after paroxysms of coughing, is a quite constant feature. Constipation is the rule, although diarrhea may ensue, with blood and mucus in the stools. Intense thirst has been noted in many of the afebrile cases, as well as in those accompanied with slight fever. Polyuria was frequently present. Vesical irritability was observed in three of the cases. In one the irritation was doubtless due to hyperacidity of the urine (70 c.c. of deci-normal solution of sodium hydroxid were required to neutralize 100 c.c. of urine in the presence of an excess of potassium oxalate). The urine is generally pale, highly acid, low in specific gravity and free from sugar, albumin and casts.

A striking feature in the study of these cases is the absence of any marked nervous or mental symptoms. There is little or no mental or physical depression. Few of the patients were found in bed on my first visit, and practically all the children affected with the disease emphatically refused to stay in bed, even with a moderate fever of 101 F. or 102 F. Most of them did not feel ill or distressed except when seized with a paroxysm of coughing, during which they suffered from a severe choking sensation. In three instances I was hastily summoned, as the parents thought the children were dying from strangulation. During these attacks, which lasted only a few minutes, the patients became cyanosed and exhausted. Immediately after the attack they felt as well as usual. One patient, a girl of 20, had five such attacks in twenty-four hours. After a thorough cleansing of the larynx by means of an alkaline spray they promptly ceased.

Mild frontal headaches, substernal and epigastric pains are frequently complained of. Dr. H. C. Knapp furnished me with notes on an interesting case of pneumococcic bronchitis in which there was a pseudomembrane in the throat, and severe joint pains especially in both knees and ankles. With the exception of chickenpox vesicles in one case, sudamina vesicles in another and impetigo in a third, no eruption was observed. Many cases developed a mild adenitis, especially of the postcervical glands. The spleen was only enlarged in those cases associated with severe intestinal symptoms. A decided leucocytosis was observed in all cases in which the blood was examined, ranging from 12,500 to 21,000.

SPECIAL SYMPTOMS AND COMPLICATIONS.

A significant feature of this disease is the formation of a pseudomembrane. This occurred in 12.5 per cent. of all the cases. The membrane is of a light yellow color, usually situated in the nose, pharynx or nasopharynx and occasionally the conjunctiva. It is easily removed, but usually leaves a bleeding surface of the mucous membrane. Cover-slip preparations show an enormous amount of encapsulated pneumococci, and di-

7. Accordino, *Supplemento al Policlinico*, 1899.

8. The predisposition of children to pneumococcus conjunctivitis has been shown by Axenfeld in Külle and Wassermann's *Handbuch*.

9. Nicholas, C.: *Etude sur la conjunctivites à pneumonocoques*, Thèses, Paris, 1901.

10. Halle (*Annals d'Oculist*, cxxiii, p. 200) found the period of incubation of pneumococcus conjunctivitis very variable. An accidental infection in one case resulted in the development of an acute conjunctivitis on the third day; in another case on the seventh day.

11. Denig (*Ztschr. f. Augenheilk.*, iv, p. 213), has called attention to the association of pneumococcus conjunctivitis and pneumococcus angina.

rect inoculation into animals produces pneumococcic septicemia. Severe bronchitis was present in ten of the cases. Some of them which have been carefully studied by Dr. Stokes prove almost beyond a doubt to be due to a pneumococcus infection. Vomiting of clear or bile-stained mucus after coughing occurred in 37.5 per cent. of the cases. This appeared to affect small children more constantly than adults. Vomiting was often accompanied with intense retching. Diarrhea complicated four of the cases in this series, all of which occurred in children under 4 years of age. They averaged from three to seven movements daily. The stools were thin, highly offensive, and contained a large amount of mucus which was occasionally streaked with blood. Fibrinous pleurisy complicated two cases and frontal sinusitis and purulent otitis media each complicated a single case.

It is strange to say that pneumonia did not develop in any of these cases, although Dr. C. F. Blake informs me that he has recently seen two cases follow an acute catarrhal condition, which answered the description of this disease.¹²

COURSE AND RELAPSE.

The average duration of the acute stage is from one week to ten days. The cough frequently persists much longer, and in some instances it continued for five or six weeks. Six of the patients had definite relapses; five occurred in young children and one in an adult. The interval between the subsidence of the primary attack and the onset of the relapse ranged from one to two weeks.

DIFFERENTIAL DIAGNOSIS.

If this is an independent affection, then it is necessary to differentiate it from: 1, measles; 2, whooping-cough; 3, influenza; 4, an epidemic catarrh due to the *micrococcus catarrhalis*, and, 5, epidemic conjunctivitis.

Measles.—The mere fact that there is a uniform absence of Koplik spots and rash should suffice. If, however, in addition, we consider the fact that having had measles does not lessen the susceptibility of the individual to an attack of this disease, we can safely exclude the possibility of measles.

Whooping-cough.—While it is true that in a small percentage of cases there is a distinct whoop, yet this appears as early as the second day of the illness and usually disappears within a period of one week. Moreover, the most typical cases that I saw, developed in children who have had whooping-cough.

Influenza.—With the exception of the absence of a mental and physical depression and severe nerve pains, the condition presents a close resemblance to influenza. Other differentiating features to be considered in the present epidemic are the tendency to the development of pseudomembrane and conjunctivitis; the peculiarly characteristic cough; the more or less persistent hoarseness, and, above all, the absence of the Pfeiffer bacillus.

Epidemic Catarrh.—This is generally a mild catarrhal affection and is not characterized by any gastrointestinal symptoms. The characteristic organism, *Micrococcus catarrhalis*, was not found in any of the patients examined.

Epidemic Conjunctivitis.—As far as the eye symptoms are concerned, there are no distinguishing features to separate it from the epidemic pneumococcic conjunctivitis which was prevalent in a number of large com-

munities in continental Europe in 1896, as described in the report of Kölle and Wassermann. These authors, unfortunately, merely called attention to the fact that there is associated an acute catarrhal condition, and look on the eye symptoms as the essential lesion. In this epidemic, the eye symptoms are subsidiary, the primary seat being almost invariably in the upper air passages, and, as already stated, only 37.5 per cent. developed pneumococcic conjunctivitis.

TREATMENT.

The cough is very intractable and little influenced by ordinary cough mixtures. Antipyrin and the bromids diminish the intensity and frequency of the paroxysms. Much relief is obtained by a thorough cleansing of the mucous membrane of the nose and throat by the use of an alkaline spray. The mild forms of conjunctivitis yield readily to the local application of boracic acid solution. The severe mucopurulent and membranous forms were successfully treated with 10 per cent. argyrol solution. Quinin administered in large doses (twenty-five to thirty-five grains daily) has proved an efficient remedy. It has the effect of promptly reducing the temperature and lessening the cough and coryza.

From the clinical and bacteriologic study of these cases, we believe that this affection is an acute pneumococcic disease of a decidedly epidemic character.

BACTERIOLOGIC STUDY.

Before describing the findings in the bacteriologic study of this acute, inflammatory, communicable infection of the conjunctiva and upper air passages, it is proper to mention the various bacteria which are found in diseased conditions of these mucous membranes.

The conjunctival epidemic inflammations are usually produced either by the Koch-Weeks bacillus or the Morax-Axenfeld organism. The former bacillus is very small, decolorizes by Gram's method, is found in and outside of the pus cells, and will only grow on blood agar or other similar media. The latter organism is a much larger, thicker bacillus which often occurs in pairs and also decolorizes by Gram's stain. It may be cultivated on blood serum. Among the other bacteria may be mentioned *B. diphtheriae*, *Micrococcus gonorrhoea*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Meningococcus intracellularis*, *Bacillus coli*, *Bacillus pyocyaneus* and *Diplococcus pneumoniae*. Any of these organisms also may produce inflammation of the mucous membranes of the throat and nose.

In most of the cases which we examined bacteriologically we found the *Diplococcus pneumoniae* at times in combination with the pyogenic micrococci, and the pneumococcus has previously been reported by several investigators as the cause of epidemic conjunctivitis. These outbreaks are described in Kölle and Wassermann's Bacteriology, 1903.

The first cases of pneumococcic conjunctivitis were reported by Parinaud¹³ and Morax¹⁴ in 1894 as occurring in newborn infants and young children, and these cases were accompanied by lachrymation, coryza and purulent and fibrinous inflammation of the conjunctival mucous membranes.

Gasparrini¹⁵ and Axenfeld¹⁶ in 1896 reported many cases of pneumococcic conjunctivitis in children and

13. Ann. d'ocul., 1894.

14. Recherches bacter. sur l'etiol. des conjunctivites aigues. Thèse de Paris, 1894.

15. Annali d'ottalm., 1893, xxii, 6, and 1896, xxv.

16. Verhandlungen der ophth. Gesellsch. Heidelberg, 1896; Deutsche med. Wochschr., 1898, No. 1.

12. The association of pneumococcus conjunctivitis and of pneumonia was pointed out by Pymowicz in 1902 (*Postep. Okul*, No. 11, Polish).

adults, and they considered the infection communicable.

The geographic distribution of this disease is widespread, and epidemics have been described as occurring in several cities in Germany, Austria, Italy, Denmark, Switzerland, Russia, Egypt and the United States. In these cases the inflammation was limited to the conjunctiva, and we have been able to find only one case reported by Denig¹⁷ in which this condition was accompanied by an angina, or fibrinous inflammation of the throat.

It is difficult to produce an inflammation of the conjunctiva in rabbits and other animals by means of the pneumococcus, although this has been accomplished after scarifying the conjunctiva. The history of the outbreaks among human beings, however, makes it highly probable that the condition is communicable, and this was proven by Gifford¹⁸ and von Pichler¹⁹, who produced the disease in healthy persons by inoculating their eyes with pure cultures of the pneumococcus obtained from cases of epidemic conjunctivitis. Hauenschild²⁰ produced the disease by inoculating the pus from the eye into healthy eyes, and proved an incubation period of about two days.

From what has been said it can be seen that the clinical picture of this condition has almost entirely related to the conjunctiva. Dr. Beck, however, has demonstrated that the mucous membranes of the throat and nose are also frequently affected both by purulent and fibrinous inflammations, and in our bacteriologic studies we have made routine cultures from the nose, throat and eye of all the cases examined. In addition, in most of the cases, we have examined cover slips from the eye, nose and throat, and the cultures have frequently been inoculated into white mice or rabbits to prove the virulence of the pneumococcus.

So far we have studied 33 cases, and by microscopic examination have found pus cells and pneumococci in 7 specimens from the nose, 9 from the throat and 10 from the eye. Capsules were stained by Welch's method in 1 preparation from the nose, 3 from the throat and 2 from the eye. We failed to find pus cells and pneumococci in 2 specimens from the nose, 1 from the throat and 3 from the eye. No microscopic examinations were made of the other cases.

The pneumococcus was found in cultures in 8 cases from the nose, in 2 of which fibrinous membranes were present, and in 23 cases from the throat, in 4 of which membranes were present. In 7 out of 17 cases the cultures from the eye contained pneumococci, and in the rest of the 33 cases the eyes were not examined. The original cultures from most of our cases were inoculated into white mice, and the pneumococci often proved virulent, killing the animals with general septicemia in from 1 to 11 days. The cultures from the eye produced pneumococcic septicemia 3 times and those from the throat and nose each 5 times.

In the second series of 16 cases we inoculated 8 rabbits intravenously with pure cultures from the throats of the patients, and 5 died of general pneumococcic septicemia. In 29 out of 33 cases the pneumococcus was obtained either from the nose, throat or eye by cover-slips, cultures or animal experiments.

In order to exclude the Koch-Weeks and influenza bacilli, cultures were made from the throat and eye in 5 cases on blood agar with negative results. We also

failed to find any of those very fine, Gram-decolorizing bacilli in the microscopic examinations of cultures from the eyes, nose and throat, and think that this organism may be excluded as a causative agent. In many of the cases we obtained the pneumococcus in pure cultures, but some of the throat and nose cultures contained the *Staphylococcus aureus* and *albus* and the xerosis bacillus. Several cultures from the eye showed the presence of these pyogenic micrococci. We think that these organisms are secondary invaders and that the inflammatory condition can be produced by the pneumococcus alone.

In several of the cultures from the pus from the eyes and membranes in the nose and throat and in cover-slips from these materials only pneumococci were found. Frequently the pneumococci were within the pus cells showing phagocytosis. We consider this, therefore, the primary infectious agent.

These cultures gave the usual cultural and staining characteristics of the *Diplococcus pneumoniae*, although we made no attempt to distinguish between the two forms of the organism, in respect to its effect on culture materials containing inulin. The diagnosis was based on the presence of capsules, the coagulation and acidulation of milk, the clouthing of bouillon, and its failure to grow well in gelatin.

In ascribing to the pneumococcus the causative rôle in this disease, we have not disregarded its presence in the normal mouth and throat. The "Report of the Medical Commission for the Investigation of Acute Respiratory Diseases of the Health Department of the City of New York" contains several investigations which show that the pneumococcus is an almost constant inhabitant of the throat and mouth. Park and Williams found either typical or atypical pneumococci in 51 out of 80 normal mouth secretions examined, and Longcope and Fox found typical pneumococci in 19 out of 40 normal throats which they examined. Duval and Lewis isolated this organism in all of 24 healthy persons examined, and Berger found it in 110 out of 283 normal throats examined. The pneumococcus, therefore, can be considered an almost constant inhabitant of the mouth and throat.

But when this organism is discovered in diseased conditions of the mucous membranes, usually in pure culture; frequently virulent, and so distributed in purulent and fibrinous inflammations as to explain these conditions, we have no hesitancy in considering it as the cause of the condition. The photomicrographs from typical cases also show the organism in large numbers, often within the pus cells, showing active phagocytosis, and this is a further point in favor of its being the cause of the disease.

CONCLUSIONS.

This disease exists as a distinct entity, characterized by purulent or fibrinous inflammation of the mucous membranes of the eye, nose and throat.

The infection can be communicated from diseased to healthy persons.

The infection is caused by the pneumococcus, resembling in its cultural and pathogenic properties the cause of lobar pneumonia.

Physical Morality.—Perhaps nothing will so much hasten the time when body and mind will both adequately be cared for, as a diffusion of the belief that the preservation of health is a duty. Few seem conscious that there is such a thing as physical morality.—Herbert Spencer.

17. Ztschr. f. Augenheilk., 1900, 213.
18. Arch. of ophthal., 1896, xxv, 314.
19. Beitr. z. Augenheilk., 1896, xxiv, 19.
20. Ztschr. f. Augenheilk., 1900, iii, 1.