ABSTRACT OF DISCUSSION

DR. FRED WISE, New York: While the theory of the tuberculous origin of erythema nodosum is very interesting, it is scarcely in accord with the short duration and practically spontaneous disappearance of the lesions in some of these cases. Lesions which are caused by the presence or activity of the tubercle bacillus in the tissues are almost invariably chronic and persistent in nature. How, then, should we reconcile the presence of tubercle bacilli in the lesions of erythema nodosum, when they are so evanescent in character?

DR. ERNEST L. McEWEN, Chicago: Dr. E. C. Rosenow of Chicago has recently succeeded in isolating an organism from the lesions of erythema nodosum in a number of cases. In one of these, of which I have personal knowledge, the patient was a young, foreign-born woman with indefinite joint-pains; in the lesions of the legs Dr. Rosenow was able to find this particular organism and to show its close relationship to the streptococcus which he has demonstrated to be the cause of articular rheumatism. If Rosenow is on the right track these cases of erythema nodosum associated with tuberculosis must be explained as Dr. Foerster has suggested, namely, that some acute infectious disease, like measles, lights up a latent tuberculosis, while coincidentally an infection with Rosenow's organism occurs. It is quite remarkable that so many cases of erythema nodosum associated with tuberculosis are reported in the literature. We have reason to expect that the future will reveal to us some wonderful things in connection with the transmutation of organisms. We know that certain strains of organisms will select certain tissues and leave others alone, a fact which is as important as it is striking. Rosenow, I understand, has been able to change his streptococcus of rheumatism, through a series of definite stages, into a pneumococcus, and back again, by animal inoculations under certain conditions. This field of research, I believe, will prove to be an extremely rich one in the near future.

DR. GEORGE M. MacKEE, New York: I believe, with Dr. McEwen, that Rosenow is on the right track. I also feel as does Dr. Wise with regard to the short duration and spontaneous recovery of erythema nodosum lesions, and I cannot conceive of a tuberculous process of so brief a duration. In erythema nodosum we have lesions that last only a few days, and it is almost inconceivable that they should be due to the toxins of tuberculosis.

DR. W. C. BROWNSON, Asheville, N. C.: With regard to the value of the von Pirquet test from a diagnostic point of view, the pediatricists have come to the conclusion that it is of doubtful value except in very young children. Still we constantly read of the von Pirquet test as an aid in the diagnosis of tuberculosis verrucosa cutis or blastomycosis. As a matter of fact, the test is of little value except in young children.

DR. J. B. KESSLER, Iowa City, Iowa: This discussion reminds me of a patient I had last winter who was suffering from erythema nodosum. The case had been regarded as one of lues until that diagnosis had been disproved by several negative Wassermann tests. The paper of Dr. Foerster and the work of Dr. Rosenow certainly show us the way to make a correct diagnosis in these cases.

DR. A. RAVOGLI, Cincinnati: I am not inclined to believe that erythema nodosum has any relationship to tuberculosis. I have seen only a few cases of this affection, usually occurring in young girls, ranging in age from 12 to 19, and giving a history of some acute antecedent disease, usually a sore throat together with arthritic pains. The attacks were of brief duration, the eruption being usually confined to the lower extremities, in the form of hard, oval, red erythematous nodules, which disappeared in a short time, leaving the patient in an apparently normal condition. For these eruptions I think Rosenow has given us a very good explanation, namely, that the staphylococcus or some hybrid form of streptococcus from the tonsils enters the circulation and when it reaches the small ramification of the capillary blood-vessels, produces a kind of thrombi which causes exudation and the erythematous eruption. Bazin's disease, erythema

induratum, I have always regarded as a localized tuberculosis, as it often ends in ulcerations.

DR. RICHARD L. SUTTON, Kansas City, Mo.: I have recently done some work in connection with sarcoids (Boeck and Spiegler-Fendt), erythema nodosum and the erythema induratum of Bazin. Beginning with nodular tuberculosis of the hypoderm (Wende), we can readily separate the various types of lesions into definite clinical groups. There are undoubtedly three distinct varieties of "erythema nodosum," and perhaps four: erythema nodosum directly due to bacteria (Rosenow), erythema nodosum due to the tubercle bacillus or its products, and erythema nodosum occurring as a result of the ingestion of certain foods or drugs. The tuberculous type of erythema nodosum I think should be designated as "erythema indurata disseminata," to distinguish it from the other varieties.

DR. O. H. FOERSTER, Milwaukee: I wish to emphasize that many of the cases reported in the literature have been in children in whom the erythema nodosum has been followed by a tuberculous meningitis or other tuberculous manifestation.

DISEASES AND NEW GROWTHS OF LYMPHATIC ORIGIN *

PROF. G. ARNDT, M.D.

Privat-docent in Dermatology, University of Berlin

BERLIN, GERMANY

When I was first asked to read a paper before the Section on Dermatology of the American Medical Association I hesitated to accept because I did not know anything new to communicate. Then my friend, Udo Wile, proposed that I speak about the changes of the skin in lymphatic leukemia, a subject I have written about. Though many of you will be quite familiar with what I am going to say, there may be some who have not had much opportunity to study this question.

Skin troubles may be the first prominent symptom of lymphadenosis and the skin specialist familiar with these troubles will, in a good many cases, be able to make the diagnosis in a very early stage of the disease. Concerning the nomenclature I think the old names lymphatic and myeloid leukemia should be replaced by aleukemic, subleukemic and leukemic lymphadenosis and myelosis. The essential pathologic feature of lymphadenosis is the generalized hyperplasia of the lymphatic hematopoietic tissue and the essential feature of the myelosis is a generalized hyperplasia of the myeloid tissue. According to the blood-changes, which represent only a symptom, although a very important one—we distinguish, in speaking of lymphadenosis, an aleukemic form, if the blood is normal, a subleukemic form if the number of the white blood-cells is normal or very little increased, but the proportion of the lymphocytes is augmented, and a leukemic form when there is an absolute, permanent, progressive increase of lymphocytes and the total number of leukocytes is augmented.

The changes of the skin that may be observed in the beginning or in the later course of lymphadenosis can, from clinical and histologic points of view, be divided into changes not characteristic at all and not distinguishable from clinically and histologically similar affections of other origin on the one hand, and on the other, changes representing proper localizations of the lymphadenotic process in the skin. The former more frequently met, are called leukemids, a name

* Read before the Section on Dermatology at the Sixty-Fifth Annual Session of the American Medical Association, Atlantic City, N. J. June, 1914.

suggested by Audry; the latter are the lymphadenoses of the skin in the proper sense of the word.

LYMPHADENIDS (LEUKEMIDS)

Among the skin lesions not characteristic in themselves are to be mentioned the purpura met especially in the so-called acute mostly large-celled leukemia, first described by Friedreich, Ebstein and A. Fraenkel. In those cases, multiple swelling of the lymph-nodes, enlargement of the liver and spleen may be completely absent, and the most striking features of the clinical picture are either petechial or more diffuse hemorrhages of skin and mucous membranes, the latter often terminating in necrotic lesions of the gums and the mucous membrane of the mouth and suggesting scurvy.

If there is fever or severe troubles of the general health one first thinks of a septic affection leading to the hemorrhagic diathesis and only the blood-picture enables one to make the diagnosis, which might be quite impossible in the aleukemic forms of this type. The purpura can be combined with the formation of blisters, described by Shattuck in this country.

Then there must be mentioned the so-called prurigo lymphatica (first described by E. Wagner, Buschke, Breseendorff). Like the prurigo of Hebra the prurigo lymphatica is characterized by numerous pinhead- to hemp-seed-sized semiglobular or more conical, pale, elastic, edematous papules each bearing on its top a deep-seated vesicle that soon gets scratched off and is replaced by a crust. The skin between these papules becomes thickened, edematous, as the effect of the continual scratching. The surface is dry and scaly. Secondary pus infection (impetigo, furuncles) complicate the picture. In a word, there is, morphologically, a very great resemblance between prurigo lymphatica and the prurigo of Hebra. Whereas, Hebra's disease, however, always starts in early childhood, recurs in more severe attacks during the winter or summer, the prurigo lymphatica starts at every age and shows exacerbations in certain seasons. In prurigo lymphatica one does not find the typical localization of the prurigo of Hebra.

Prurigo lymphatica is not only observed in aleukemic and leukemic lymphadenosis, but also in other more or less generalized affections of the lymphatic apparatus, parenchymatous as well as granulomatous, in lymphosarcoma (Wunderlich, Dubreuilh), in some cases of tuberculosis of the lymphatic glands, in hyperplastic conditions of the lymphatic glands observed in malaria, in lymphogranulomatosis (Hodgkin's disease). Neither the clinical nor the histologic picture (thickening of the epithelium, formation of rare vesicles, status spongiosus, chronic inflammatory changes of the superficial layers of the derma) is sufficient to prove the lymphadenotic nature of the condition. The diagnosis has to be based on the blood-finding or the histologic examination of an excised lymph-node.

In other cases you might observe an intense itching, a pruritus, without any visible lesions of the skin or secondary changes, due to scratching or pus infection. This pruritus is not of itself characteristic in any way. Very instructive and convincing of the connection between affections of the lymphatic system and pruritus is the case published by Blaschko. A patient with lymphosarcoma of the supraclavicular glands suffered from very severe itching. After removal of the glands the pruritus disappeared but came back four years later with a recurrence of the lymphatic new-growth.

In other cases of lymphadenosis, one can observe common urticaria, urticaria papulosa or papulovesiculosa (lichen urticatus), multiple formation of furuncles and carbuncles.

More interesting are the cases of universal exfoliative erythrodermia observed in the course of lymphadenosis by Elsberg, Peter, Wassermann, Nicolau. Those cases show a close resemblance to the so-called pityriasis rubra of Hebra and many observations of cases published under this name may be allied to a lymphadenotic condition. The "leukemic and pseudo-leukemic erythrodermias" are characterized by a sub-acute or chronic progressive reddening of the skin spreading in a shorter or longer lapse of time over the whole, or almost the whole body, by formation of bran-like or flaky scales and enlargement of the lymph-nodes. The latter symptom, however, can be absent as in Wassermann's case, in which Ehrlich had made the diagnosis of lymphatic leukemia on account of the blood-finding and the post-mortem detected a considerable enlargement of the retroperitoneal glands caused by a pure lymphoid hyperplasia.

There may be swelling of the liver and the spleen, but these symptoms are very often absent and the diagnosis can be proved only if an increase of lymphocytes in the blood is shown, the total number of leukocytes being normal or augmented. In case the blood-picture is aleukemic, a histologic examination of an enlarged lymph-node is to be made. To some extent the skin-lesions vary. There may be more or less complete loss of hair and nails; the latter may be cracked and thickened; in one case the skin is thickened, edematous, while in another it is thinned, atrophic, leading to retraction (ectropion of the eyelids, for instance). Fever is frequent but may be absent. Among the subjective symptoms must be mentioned the feeling of chill common to all universal erythrodermias and due to the augmented loss of heat and the itching, mostly not very marked, but sometimes rather intense.

The histologic changes consist in a continuous diffuse or more circumscribed infiltration of the upper layers of the derma, the stratum papillare and sub-papillare, the blood-vessels of which are much dilated. The infiltration is formed by small deeply stained round cells—morphologically and after their tinctorial properties identical with the small lymphocytes of the lymph-nodes and larger elements with pale oval nuclei—proliferated connective-tissue cells. The lymphocytes surround the vessel follicles, sweat-glands and sebaceous glands; the fibroblasts are more diffusely spread between. Except the cell-forms mentioned you find mast-cells and pigment cells, the latter sometimes in very large numbers. In Nicolau's case they formed at the lower border of the infiltration a band consisting of three or four rows of cells. The rete malpighii is mostly atrophic and covered by a more or less hypertrophic horny layer.

A differentiation of these "pseudoleukemic and leukemic erythrodermias" from other universal chronic exfoliative reddenings of the skin of unknown origin, especially from the genuine pityriasis rubra of Hebra, is quite impossible by means of clinical and histologic study of the skin only. It has already been mentioned, and especially Jadassohn and Nicolau insist on the point, that some of the cases of Hebra's disease belong to the lymphadenosis (lymphatic leukemia and pseudo-leukemia of the old nomenclature). A critical analysis in that respect is quite impossible; so far as I know in no case published as pityriasis rubra of Hebra

has a reliable blood-count been made. In every case of chronic universal exfoliative erythrodermia the blood must be examined at different times, repeatedly, with special regard to the differential count, and the diagnosis of pityriasis rubra of Hebra can be made only after excluding a leukemic and a tuberculous erythrodermia (cases of the latter described by Jadasohn, Brunsgaard, Wertheim-Finger, as you know). As the histologic findings of the universal exfoliative erythrodermias encountered in lymphadenosis are not especially characteristic, common chronic inflammation (perivascular round-cell collections and proliferation of the connective-tissue cells), they must be differentiated, at least for the present, from the clinically, not so much, as the histologically absolutely characteristic, universal, lymphadenotic (lymphatic-leukemic) infiltrations of the skin.

Suppose the cause, the pathologic agent or agents of lymphadenosis will be discovered one day, suppose the bacteriologic facts found by E. Fraenkel and Much, and by myself—not very convincing now—will be confirmed by other authors and put on a solid experimental basis—suppose all that, and the more or less artificial borders put up between the true clinically and histologically well-characterized lymphadenosis of the skin and the non-characteristic so-called lymphadenitis (leucémids—Audry) will disappear.

THE PROPER LYMPHADENOSIS OF THE SKIN

The proper (leukemic, subleukemic and aleukemic) lymphadenosis of the skin represents a pathologic condition completely equivalent to the changes of the inner organs caused by a generalized hyperplasia of the lymphatic tissue. The lymphadenosis of the skin may be divided (after Riehl, Paltauf, and F. Pinkus) into a diffuse or better, universal form, and a circumscribed form causing flat infiltration, nodes or tumor-like swellings.

In the so-called myelosis (myeloid leukemia)—characterized by a generalized hyperplastic proliferation of myeloid tissue—a diffuse universal participation of the skin with typical histologic structure (myeloid tissue) has not been observed. There is a circumscribed myelosis of the skin, however, first perhaps described by Hindenburg and lately by Brunsgaard from Christiania, in a very remarkable paper. Histologically myelosis cutis shows perivascular myeloid cell-formations, that mean neutrophil and eosinophil myelocytes, myeloblasts, giant-cells (megacaryocytes) and nucleated red corpuscles. In comparison with the circumscribed lymphadenosis the circumscribed myelosis of the skin is extremely rare and there have been observed only a few cases.

UNIVERSAL LYMPHADENOSIS OF THE SKIN

The conception of the universal form is not very well defined and has not been accepted by all authors, a fact probably due to the extreme rarity of the cases. Some authors—for instance, Felix Pinkus, who has written an excellent article on lymphatic leukemia of the skin—mix the universal form up with the so-called lymphoderma perniciosum.

The name "lymphoderma perniciosum" has made great confusion and has been applied to quite different processes showing certain clinical similarities. Kaposi first used this name for a case studied by himself and well known from the literature. No exactly analogous case has ever been observed since, by Kaposi or any other author. Tallant thinks that the pathologic

changes in the original case ought to be considered as an atypical malignant proliferation of the lymphatic tissue-leukosarcomatosis, in the sense of Sternberg. Most of the cases reported afterward as lymphoderma perniciosum correspond to the type called by the French authors (E. Besnier, Vidal, Hallopeau, Jeanselme, Leredde) *érythrodermie prémycosique*; some of them are perhaps identical with the universal form of lymphadenosis.

A satisfactory critical analysis of all the material referring to that subject is scarcely possible, as most of the publications are of an older date and do not give a sufficient description of the hematologic and histologic picture. By clinical observation alone a differentiation of all these processes cannot be made. Clinically their similarity may be striking. Universal reddening and more or less considerable tumefaction of the skin in connection with swelling of all the palpable lymph-nodes, and with enlargement of liver and spleen may be found in generalized purely hyperplastic and perhaps also in aggressive malignant proliferation of the lymphatic apparatus as well as in certain—principally in the lymphatic tissue—localized infectious granulomas (mycosis fungoides and perhaps lymphogranulomatosis). As we do not know very much about the cause of these different conditions they must be differentiated by their pathologic picture. Their diagnosis has to be based on the histologic findings. The blood-picture may be of great value for the diagnosis if it shows characteristic alteration, but if it is normal or only slightly changed it is of no use in the classification of those affections. Therefore, I suggested to divide the universal infiltrating erythrodermas observed in the course of the different affections of the lymphadenoid tissue, on account of the histologic changes, into four groups: (1) lymphadenotic (lymphatic-leukemic), (2) lymphosarcomatous, (3) mycotic, and (4) lymphogranulomatous erythrodermias.

The number of the cases of universal infiltrating erythrodermia, in which there can be no doubt, clinically, hematologically or histologically—about their lymphadenotic nature, and which should be called lymphadenotic erythroderma (diffuse lymphadenosis of the skin) is very small and amounts only to four, including one personal observation. The first case has been described by Riehl of Vienna, to whom we are indebted for the knowledge of this disease and the second one has been observed and published by Linser of Tübingen, the third one, by Rodler-Zykin in Jadasohn's clinic, differs rather from the classic type.

I shall not bother you with the history of my own case. You will find it in my publication on this subject. There was a universal intensely itching swelling and reddening of the skin, enlargement of all the palpable lymph-nodes especially the axillary, inguinal and crural. The blood-picture was that of lymphatic leukemia, total number of leukocytes 40,000, lymphocytes 60 per cent. (among them 11 per cent. of large atypical forms). The histologic examination of an excised piece of skin showed the characteristic lesions—a continuous, diffuse, pure lymphocytic infiltration of the two upper thirds of the derma, going down to the lower third of the cutis propria and the subcutaneous tissue in form of perivascular and periglandular cell-collections.

The infiltration seems to be very monomorphous—at least at the first glance—and contains numerous newly formed much-dilated vessels. If the infiltrating cells

are examined with a higher-power lens it may be remarked that they are all lymphocytes, but that the common so-called small lymphocytes are rather scarce and that the large forms with pale round or oval, sometimes kidney-shaped and lobulated nuclei, the so-called lymphoblasts and atypical forms, prevail.

Mitoses are very numerous. There is no proliferation of the connective tissue. The original connective tissue is mechanically dislocated and forms a sort of network, the meshes of which are filled in by the lymphocytes. The epidermis is in some parts proliferated, in others atrophic. The papillary and subpapillary layers show a very marked edema, dilatation of blood-vessels, but very little infiltration, so that regarding the sections with a low power, the uppermost layer of the derma seems to be normal. About the bacteriologic findings I shall speak later.

In the later course the number of leukocytes increased considerably (up to 120,000, 78 per cent. of which were lymphocytes). The patient died one year and nine months after the first trouble of the skin had appeared. The post-mortem showed diffuse lymphatic leukemic changes of the skin, adhesive pericarditis, fatty degeneration of the heart-muscle, aneurysm of the aorta descendens, hydrothorax (left side) and edema of the lungs, leukemic infiltration of the liver, spleen, kidneys and lymphatic glands.

CIRCUMSCRIBED LYMPHADENOSIS OF THE SKIN

While the clinical picture of the diffuse lymphadenosis of the skin is not very well characterized, the diagnosis of the circumscribed form is not so difficult and can be made with more or less probability. The two forms never go into each other. The circumscribed type neither terminates in a universal swelling of the cutis nor are there well-defined tumor-like growths to be observed in the course of the lymphadenosis universalis. The circumscribed patches and tumors are observed in the aleukemic, subleukemic and leukemic lymphadenosis.

My personal observations of circumscribed lymphadenosis of the skin concern three cases of subleukemic and two cases of aleukemic lymphadenosis (in the latter the blood examined at different times being absolutely normal).

These facts are of certain value and prove the identity of processes that show pathologically the same changes but differ as to the blood-picture and were formerly strictly separated as lymphatic, leukemic and pseudoleukemic. Ehrlich and his pupil, F. Pinkus, the well-known dermatologist in Berlin, showed first that the lymphatic, leukemic and pseudoleukemic are the same process. Long before them, J. Orth had, on account of the pathologic findings, pointed out their relationship.

DIAGNOSIS OF THE CIRCUMSCRIBED LEUKEMIC AND ALEUKEMIC LYMPHADENOSIS OF THE SKIN

An absolutely certain clinical diagnosis of these lesions is impossible considering the rarity of the affection on one side and certain similarities with other processes on the other side. Nevertheless, you can, in the great majority of cases, conclude on the nature of the skin lesions, and I shall try to sketch the most important clinical features. First of all, the localization is very characteristic, the face being the seat of all predilection that is taken either exclusively or generally. It is quite an exception that it does not show any changes like in the much discussed case observed

by Nékam. There are other skin diseases, of course, that prefer the face (lupus vulgaris, lupus erythematosus), but the appearance of chronic tumor-like swellings of the face reminding one of sarcoma, must always raise the suspicion of lymphadenosis.

Especially the nose and the adjoining parts of the cheeks, the eye-lids, lips, chin, ears and ear-lobes are involved. The infiltration of the eye-brows and glabella gives the face a most characteristic appearance recalling the facies leontina of the lepers. Frequently the scalp and upper extremities, especially the backs of the hands, are affected at the same time as the face.

Among the less frequent localizations are to be mentioned the skin of the breast, especially about the mammae (Pfeiffer, second of my cases), abdomen, scrotum (one of my observations), glans penis (infiltration of the corpora cavernosa in one of my cases). Occasionally every part of the body may be involved, but the cases with numerous nodules all over the body (head, trunk and extremities), cases described by Biesiadecki, Hochsinger and Schiff, Kreibich (second case Feger), are extremely rare.

Concerning the shape of the lesions it must be mentioned that more or less well-defined, macular, slightly elevated patches and semiglobular projecting tumors develop close together and turn one into the other. I think that this fact is very characteristic and of value, making the diagnosis with the genuine new growths (sarcoma).

The formation of ringed, festooned lesions either by central involution and peripheral spreading of a single patch or by juxtaposition of different smaller patches is quite unusual in lymphadenosis, whereas it is frequently seen in mycosis fungoides. The flat patches may persist for years without transformation into tumor-like new growths. The size varies within the widest limits. The patches vary from the size of a silver dollar to an area corresponding to two palm's breadths and more. The tumors are from pea-size up to the size of two fists and are sometimes lobulated, showing that they have formed by coalescence of different smaller new growths.

The color is mostly a peculiar yellowish-brown or yellowish-brown red, rarely bluish-red, on pressure translucent, reminding one of lupus vulgaris. If there is a hemorrhagic diathesis at the same time, the tumors may be the seat of hemorrhages and look like melanotic new growths (dark blue, bluish-black). If the infiltrations are localized only in the deeper layers of the cutis propria and in the subcutaneous tissue, the surface may appear quite normal or only show a slight bluish tinge.

Sometimes the tumors are firm to the touch, sometimes soft and always of a somewhat elastic edematous consistence. The changes remain always localized to the derma and hypoderma or to the subcutis alone; they never take the underlying tissues. The visible mucous membranes are, as a rule, free from lesions except some slight edema. There are only three cases of lymphadenosis (observed by Fröhlich, Pfeiffer, Kirkoroff) with typical changes of the mucous membranes of mouth, pharynx and larynx.

Most of the cases of so-called leukemia and pseudo-leukemia cutis, with formation of nodes about the pharynx, the uvula, the soft palate, etc., or with diffuse infiltrations of the mucous membranes of the pharynx and larynx are lymphosarcoma (Case Fano and Liebmänn, case Schultz). A very striking feature is the absolute want of any signs of a regressive metamor-

phosis (softening disintegration). Even if rather large tumors disappear suddenly, for instance, after an erysipelas, or a pneumonia, there is only a slight atrophy of the skin left. True scarring is never observed. It is quite exceptional that the new growths ulcerate. Only in one case (Nicolau) has there been a superficial ulceration. The course is very characteristic. After growing rather rapidly to a certain size they may remain stationary for years or grow but very slowly. The subjective symptoms are, in the majority of cases, very trifling and consist in some burning sensations recurring from time to time, and in a certain pain on pressure.

Of great importance in diagnosis are, of course, multiple symmetrical swellings of the lymph-nodes especially of the retromaxillary and submaxillary regions, about the neck, the axillae and the groins. These enlarged glands may be found in other diseases of the lymphatic system, however, leading equally to tumor-like conditions of the skin, and on the other hand, in lymphadenosis the palpable lymph-nodes may be of normal size. In three of my cases nobody would have thought of lymphatic leukemia by examining the lymphatic glands. The same is true of enlargement of the liver and the spleen.

The blood-picture is a great help if it shows the characteristic changes of leukemic and subleukemic lymphadenosis, but if it is aleukemic, that means normal, it cannot influence the diagnosis. The blood-finding alone is not always reliable, since leukemic blood-findings may be seen in atypical new growths of the lymphatic system (lymphosarcoma, leukosarcomatosis) and in certain forms of mycosis fungoides. Therefore, a histologic examination should be made in every case.

The histologic picture is absolutely typical and shows a pure, lymphocytic infiltration of the skin and the subcutaneous tissue, stopping with a very sharp line just beneath the papillary layer. The epithelioma (normal or flattened) is always separated from the tumor-like infiltration by a narrow, somewhat edematous zone, in which one can only with a high-power lens detect some newly formed cells. This distribution is very characteristic. The original collagenous tissue is dislocated and forms a kind of reticulum; the elastic tissue has been destroyed in the center of the large tumors. In the smaller patches, it may be rather well conserved.

In the large tumors absence of follicles, sebaceous glands and sweat-glands may be observed. In the smaller patches one can see that the process very often starts in the neighborhood of the hair follicles, the sebaceous glands and sweat-coils, which are well conserved. This perifollicular, periglandular localization is due to the fact that follicles, etc., are surrounded by a very well-developed system of capillaries.

At the first glance and employing the ordinary stains, it looks as though there was only one cell-form, small, more or less round elements with a very narrow border of protoplasm or without protoplasm, and a dark-stained nucleus showing sometimes a sort of wheel structure employing certain stains (polychrome, methylene blue, methyl green, pyronin and a combination of Much, Grünwald, Gram), one is able to differentiate between the different forms of cells.

Most of the infiltrating cells are of the type of the so-called small lymphocytes of the lymph-nodes but there are also irregularly scattered lymphoblasts and

lymphoblastic plasma-cells (cells with a large, pale, round or kidney-shaped, lobulated nucleus, two or more very distinct nucleoli and a narrow border of protoplasm, that takes the pyronin more or less intensely). At the periphery there are some proliferated connective-tissue cells.

Lymphocytic plasma-cells (type Marschalko) are absent in the majority of cases but sometimes they are found, as in one of my cases, in which they were rather numerous in some sections. Mitoses are very rare.

PATHOGENESIS

There are two different theories about the origin of the lymphadenotic infiltrations of the skin. Some authors (Kreibich, Nékam, Nicolau) thought that innumerable lymphocytes emigrated and were deposited in the neighborhood of the blood-vessels; that there was no local proliferation of lymphocytes at all or only to a very small amount. In favor of this theory, the above-mentioned authors noted the absence of inflammatory changes and proliferative lesions of the connective tissue, the lack of mitoses, in spite of the extraordinary quantity of newly formed cells, the presence of small lymphocytes in the lumen, the walls and the neighborhood of the same vessel and the universal distribution in some cases.

This theory was opposed by Felix Pinkus. Basing his theory on Ehrlich's doctrine that the lymphocytes had no self-movement, he said it was quite impossible or at least very improbable that the numerous cells forming the lymphadenotic infiltrations came exclusively from the blood. Concerning the lack of mitoses, he found something analogous in the lymphatic glands of his own case—very few mitoses and absolute absence of germinating centers.

He thinks that in the early stages of lymphatic leukemia there is an overproduction of lymphocytes but later on a sort of stasis takes place. Based on Ribbert's theory that the lymphatic tissue is spread all over the body, he explains the origin of the lymphadenosis cutis by a local proliferation of the preformed lymphatic tissue of the skin, produced by the leukemic condition, analogous to the lymphatic proliferations in the lymph-nodes, the spleen, marrow, lungs and kidneys. Suppose that, normally, there is lymphatic tissue in the skin, the theory of Pinkus seems to be the most plausible. There can be no doubt that in some cases the cell-proliferation by indirect division exists. Nékam and Nicolau insisted on the lack of mitoses. I found karyokineses in two of my cases of circumscribed lymphadenosis cutis and in my case of universal lymphadenosis they were quite numerous.

On the other hand, it is possible that some of the lymphocytes come from the blood. We know from the researches of Max Schultze, Hirschfeld, Wolf, Maximow, Jolly, Helly, Schwarz, Schridde and Zieler, that the lymphocytes are capable of moving or at least acquiring this power of self-movement under certain pathologic conditions.

Schridde lately contested the existence of rudimentary lymph-follicles in the fetal connective tissue, considering the perivascular cell-collections found by Ribbert to be myeloid formations (consisting of myelocytes, myeloblasts, nucleated red cells, megakaryocytes) which are wide-spread in the embryonic organism. For the explanation of the perivascular formation of the lymphadenotic infiltrations of the skin the preexistence of preformed lymphatic tissue is not nec-

essary. It is quite possible that the lymphatic new growths originate from the perivascular lymphocytes of the adventitia, the so-called *Adventitiazellen* (Marchand) or from the endothelial cells of the lymphcourses changed by metaplasia (Schridde) or from the so-called *Gefahrsrandzellen* (Pappenheim).

ETIOLOGY

We know nothing absolutely certain concerning the cause of the generalized hyperplasias of the lymphatic system. The so-called acute leukemia is perhaps of infectious origin, although it is sometimes quite impossible to exclude a secondary infection in these cases, and many authors think that the acute forms are quite another disease and have nothing to do with the chronic lymphadenosis. Concerning the cause of the latter, some pathologists (Orth, Pappenheim) suggested that it was of an infectious nature; others think lymphatic leukemia must be placed among the genuine neoplasms; others consider it as a special trouble quite different from the true new growths and from the infectious granulomatous conditions.

E. Fraenkel and Hatluch inoculated the emulsion of a lymphatic leukemic lymph-node into six guinea-pigs and found, on smears of two swollen lymph-glands, large, partially granular rods, three or four times as thick as tubercle bacilli and partially or completely stained by Gram stain. E. Fraenkel and H. Much found the same Gram-positive, thick, partially granular rods in the antiformin sediment of spleen and lymphatic glands in other cases of lymphatic leukemia. There were no acid-fast rods to be found. The leukemic bacilli of E. Fraenkel and H. Much could not be confirmed by other authors and it is, of course, absolutely necessary to control their findings on a larger material and by animal inoculations.

Then there should be mentioned the very interesting case published by Coley and Ewing in the *Transactions of the New York Pathological Society* under the title of "Acute Lymphatic Tuberculosis (Pure Generalized Lymphatic Hyperplasia Caused by Tubercle Bacilli)." In one of my cases of circumscribed leukemia of the skin, I found, in the sections of skin and spleen, partially extracellular, acid-fast, Gram-positive bacilli. Some of them looked like tubercle bacilli, some of them were thicker. In the case of universal lymphadenosis of the skin I found the same organisms, acid-fast Gram-positive and antiformin-fast bacilli. As I had not at all expected to find acid-fast bacilli in lymphatic leukemia, I had, unfortunately, not inoculated guinea-pigs or monkeys and I cannot say anything about the nature of these micro-organisms. They show, however, that we will have, in every case of lymphadenosis, to look for them and to put these findings on an experimental basis.

The prognosis of the lymphadenosis of the skin coincides with the prognosis of lymphatic leukemia in general.

ABSTRACT OF DISCUSSION

DR. H. H. HAZEN, Washington, D. C.: We American dermatologists have known for a long time of the splendid work of Professor Arndt. We have read some of his articles on skin diseases due to lymphatic derangements and are amazed at the depth and breadth of his pathologic knowledge. This proves that in order to be a good dermatologist, one must first of all be a good general pathologist. We have in this group of diseases the sarcomas, including that of Kaposi, and the curious tumors described by Pollard and others. Next come the Spiegler-Fendt sarcomas, and then the various other

sarcomas, between which there is often no hard-and-fast dividing line. On the other hand, and likewise grading into the sarcomas, we find the lymphatic diseases, which may roughly be divided into leukemic and aleukemic. In the first group belong acute and chronic lymphatic leukemia, which merge one into the other, and the myeloid leukemia and also chloroma. In the aleukemic group we may place multiple myeloma, Mikulicz' disease, Hodgkin's disease, mycosis fungoides and certain lymphomas such as those described by Winfield. Nearly all of the aleukemic growths may become leukemic. Wende has described a typical case of Hodgkin's disease or pseudoleukemia that terminated in leukemia, and the case of mycosis fungoides reported by Pardee and Zeit, although the blood-counts might not bear critical examination, ended in leukemia so far as the blood-picture and the necropsy findings could be interpreted. We must bear in mind that when there is any visceral complication in mycosis fungoides, it is limited to the lymphatic glands or the bone-marrow. In addition, certain cases of lymphoderma perniciosum (of Kaposi) seem to be a direct connecting link between leukemia and mycosis fungoides, while the leukemias are connected with the sarcomas through the so-called leuko-sarcoma of Strenberg, a condition of which most Americans have heard very little. Certain other diseases also show a relationship to the sarcomas. Some years ago White and Proeschner described a spirochete as the cause of some of these diseases, but their work has not been confirmed. More recently, however, Bunting and Yates seem to have isolated a bacillus as the exciting organism in Hodgkin's disease. In diagnosis, Dr. Arndt has rightly emphasized the fact that leukemic conditions may have aleukemic periods, and at such times a blood examination would be positively misleading. He calls attention to the necessity, in such cases, of studying tissue obtained by biopsy, but it seems to me that there might be some danger of confusing a leukemic condition with a benign lymphoma, as in the case of Winfield. In the treatment of these conditions, the Roentgen ray should be used, possibly combined with benzol. Will Dr. Arndt give the German conception of lymphogranulomatosis, concerning which some of us are a trifle hazy?

DR. JOHN A. FORDYCE, New York: Dr. Arndt's lucid description of these lymphatic affections of the skin will lead us to study our cases more carefully from both a clinical and a histologic point of view. In the last few years I have had two cases of universal lymphadenosis similar to the condition described by Dr. Arndt. One case was that of a woman with a generalized erythrodermia without much exfoliation, together with enlargement of the lymphatic glands. That case undoubtedly belonged to the group described by Dr. Arndt under the name of universal lymphadenosis. After trying all the usual remedies in this case, we resorted to the Roentgen rays, to which the lesions yielded rather rapidly, and at present the patient shows only a few recurrent lesions, with some thickened patches of skin. It has been found impossible completely to eradicate the condition. The picture never looked anything like mycosis fungoides. Several blood examinations were made without any definite findings, but after listening to Dr. Arndt's paper we will proceed to make a more careful examination. Some years ago I had another case of generalized thickening of the skin, with lymphatic enlargement, which at the time was regarded as one of premycotic erythrodermia.

DR. A. RAVOGLI, Cincinnati: While studying under Hebra years ago, I saw many cases of prurigo, and he usually referred to them as scrofulodermata and prescribed cod-liver oil with excellent results. The lymphatic exudation in the derma constituting the small papula of prurigo offers a good explanation for its persistency, and for the itching sensation, which it causes. In granuloma fungoides, although the leukemic condition has been advocated by Brocq and others, yet there may be a peculiar infection, which we so far do not know, which causes the proliferation and the fungous growths of the lesions. It is possible that the lymphatic system acts as an important factor in some cases of erythroderma, of which we have, so far, no idea of their etiology.

DR. MARTIN F. ENGMAN, St. Louis: The more we study and follow the literature on the differences between the so-called granuloma and malignant growths, the more we must realize that too much dependence cannot be placed on the morphologic aspect of the subject. The experimental work done by Rous and others on the dog and chicken tumors shows that in one there was produced, apparently, a granuloma and in the other a sarcoma. The original irritation, be it a micro-organism or of chemical nature, may cause certain lymph-changes and produce certain objective clinical symptoms, and as these increase, we may have later the typical blood-changes, which in the earlier types may be absent, while in other cases they follow very quickly. In the literature on mycosis fungoides, we are given the blood-findings, yet a differential count is unique. In skin affections, leukocytosis alone means nothing. We reported six cases of mycosis fungoides, with indefinite blood-findings; later, we obtained a leukocytosis, but the cells were composed of polymorphonuclear leukocytes, without any increase in the lymphocytes. In the tuberculid group we have one case now under observation, and we have found that all these cases run a high mononuclear count. The whole classification depends largely on the blood-findings.

DR. S. POLLITZER, New York: The entire subject of the leukemic diseases of the skin is relatively new and has been studied only since the establishment of hematology on a scientific basis by Ehrlich. These cases are, of course, very rare, although they are not so rare as their scant report in the literature would lead us to believe. I have been looking for leukemic affections of the skin for several years, and I am sure that in the past I overlooked some cases. I have at present under observation a man with typical lesions of leukemia, and with both a relative and absolute lymphocytosis. A relatively uncommon feature in this case is the occurrence of bullae of the deep-seated type, that when ruptured, healed slowly. Their cell-content is largely lymphocytic. I also have in mind a woman with nodular fawn-colored lesions on the extremities, pea-sized and smaller, many of them excoriated, who came to me because of a pruritus which was confined mainly to the extremities. I suspected a leukemic affection, but a blood-count showed a normal blood-picture. I then made a biopsy and was surprised to find that I was dealing with a rather unusual form of leprosy. I think it is quite impossible to separate the generalized erythrodermia of mycosis fungoides and pityriasis rubra from erythrodermia leukemica on clinical grounds. Jadassohn, who was the first to range pityriasis rubra with the tuberculids, did not by any means maintain that all cases of pityriasis rubra belonged to that class. It is evident that erythrodermia is a symptom of many different systemic disorders of a toxemic character. Cases of mycosis fungoides, which do not properly belong to the group described by Dr. Arndt, are far more frequent than those of the leukemic affections. In the extension statistics collected by the American Dermatological Association covering three-quarters of a million cases of skin diseases, we have thus far made no place for the leukemic affections of the skin, whereas mycosis fungoides has been reported in the proportion of one case to every three thousand.

PROF. G. ARNDT, Berlin: I purposely omitted any discussion of various affections of the lymphatic tissue, such as lymphosarcoma, lymphogranuloma and leukosarcoma. Lymphogranulomatosis, concerning which Dr. Hazen asks, is essentially the same as Hodgkin's disease, or malignant granuloma, with swelling of the lymph-nodes, the liver, spleen, and so on. The blood-picture in these cases is either normal, or there may be hyperleukocytosis, sometimes with eosinophilia; no lymphocytosis. The histologic picture in these cases is quite characteristic, and differs from that in ordinary hyperplasia of the skin: here we have small and large lymphocytes, with giant-cells of a peculiar type, sometimes reminding one of the giant-cells of sarcoma, with one huge, bizarre nucleus. We know nothing about the etiology of Hodgkin's disease. Many regard it as a form of tuberculosis; others think it is an infectious granuloma of

unknown origin; others, again, that the lesions may be caused by different pathologic agents, such as those of tuberculosis, leprosy or rachitis. Many different opinions exist and it is quite impossible to be positive. We see in lymphogranulomatosis, skin lesions, such as those of prurigo, pruritus and many others. The histologic features of lymphogranulomatosis are quite characteristic, and so long as we do not know the cause of those diseases, we have to differentiate them by their pathologic and histologic pictures as strictly as we can. One day, doubtless, we shall learn the cause of mycosis fungoides, but so long as we are ignorant of it, we must insist on a strict differentiation based on histologic and pathologic grounds. We often get a leukocytosis in the blood-picture in mycosis fungoides, and there is a group of cases—I have seen two or three—which present a rather characteristic blood-picture, with an increase of the large mononuclear cells and of the eosinophils, a picture which we never get in leukemia.

PARESIS PATIENTS TREATED WITH INTRASPINAL INJECTIONS OF SAL- VARSANIZED SERUM

A BRIEF REPORT*

L. B. PILSBURY, M.D.

LINCOLN, NEB.

Paresis is such a hideous and destructive disease that physicians are always on the *qui vive* for a hopeful method of treatment. Mercury, iodids and the various arsenical preparations seem to be of very little use as ordinarily administered in paresis and many cases of tabes and cerebrospinal syphilis. This fact led Swift and Ellis¹ to experiment in animals with intraspinal injections of salvarsanized and neosalvarsanized blood-serum. Later these methods were applied to man with results already familiar to you. Swift and Ellis, however, specifically declare that they are presenting their method, not as a substitute, but rather as an adjuvant to the generally accepted forms of treatment.

Concerning the rationale of the method there is perhaps some difference of opinion. Myerson² warns against expecting too much of it, partly on the ground that the flow of the cerebrospinal fluid is centrifugal rather than centripetal as regards the tissues of the central nervous system. At the same time he urges further trial on the basis of his own results. The finding of the treponema in the central nervous system has of course increased the theoretic hopefulness of the method. It is well known that meningococcus antitoxin is effectual only when injected into the subdural space or ventricles. Viton³ thinks that he can increase the permeability of the meninges by intraspinal injections of a weak solution of mercuric cyanid. If, as Mott⁴ believes, the cerebrospinal fluid plays the part of a lymph in the central nervous system acting in part as a nutrient fluid, then it should be able to convey medicinal substances to some, at least, of the brain-cells. This is in agreement with the view of

* Read before the Section on Nervous and Mental Diseases at the Sixty-Fifth Annual Session of the American Medical Association, Atlantic City, N. J., June, 1914.

1. Swift, Homer F., and Ellis, Arthur W. M.: New York Med. Jour., 1912, xcvi, 53; The Treatment of Syphilitic Affections of the Central Nervous System, with Especial Reference to the Use of Intraspinal Injections, Arch. Int. Med., Sept. 15, 1913, p. 331.

2. Myerson, A.: Intraspinal Injection of Salvarsanized Serum, Correspondence, THE JOURNAL A. M. A., Jan. 31, 1914, p. 396.

3. Viton, J. J.: Reactivation of the Permeability of the Meninges in Treatment of Syphilis of the Nervous System, Semaine méd., Dec. 11, 1913; abstr., THE JOURNAL A. M. A., Feb. 14, 1913, p. 582.

4. Mott, referred to in: The Production of Cerebrospinal Fluid, Editorial, THE JOURNAL A. M. A., March 28, 1914, p. 1018.