This article was downloaded by: [UQ Library]

On: 14 March 2015, At: 02:52

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Transactions of the Royal Society of South Africa

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/ttrs20>

ON THE DEVELOPMENT OF PIROPLASMA PARVUM (PROTOZOA) IN THE VARIOUS ORGANS OF CATTLE

Dr. R. Gonder

Published online: 26 Mar 2010.

To cite this article: Dr. R. Gonder (1910) ON THE DEVELOPMENT OF PIROPLASMA PARVUM (PROTOZOA) IN THE VARIOUS ORGANS OF CATTLE, Transactions of the Royal Society of South Africa, 2:1, 63-68, DOI: [10.1080/00359191009519363](https://doi.org/10.1080/00359191009519363)

To link to this article: <http://dx.doi.org/10.1080/00359191009519363>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

ON THE DEVELOPMENT OF *PIROPLASMA PARVUM*
(PROTOZOA) IN THE VARIOUS ORGANS OF CATTLE.

BY DR. R. GONDER.

(Read June 15, 1910.)

Since the investigations of R. Koch on East Coast Fever of cattle in East Africa, the cause of this disease has formed the subject of much discussion.

At present there exist two different views. Some investigators hold that East Coast Fever is due to an invisible organism, others declare *P. parvum* to be the cause, but neither party has been able to prove to the satisfaction of the other that their investigations were free of errors. This is due to the fact that the cattle of Africa are, as a rule, infected with various parasites; with *P. bigeminum* and the more or less harmless parasite *P. mutans*, which, morphologically, is very difficult to distinguish from *P. parvum*. Moreover, there is much controversy on the so-called plasma granules or Koch's bodies, also known as blue bodies, which are found in the organs of cattle suffering from East Coast Fever.

In this short publication, which will be succeeded by a more detailed one, I wish to communicate some investigations on the blue bodies of Koch which will throw some light on their nature. For the present I shall leave out details, neither can I enter into the literature on the subject, which will have to be considered in the more exhaustive report.

Experiments were carried out since March of this year in the Transvaal Veterinary Bacteriological Laboratory. The material used consisted in cattle whose origin we know and in ticks with whose history we were fully acquainted; accordingly double infections could be excluded from the very start. The ticks placed on control cattle imported direct from England proved that they were exclusively infected with East Coast Fever. In addition to this, Dr. Theiler gave me permission to utilise the cattle used in various experiments on East Coast Fever which had been infected experimentally through the implantation and inoculation of different organs. For my zoological studies, therefore, a considerable amount of material was available.

Before discussing the question of Koch's bodies, I wish to make some remarks on *P. parvum*. Last year Ollwig, at the Congress of Microbiologists in Berlin, stated that *P. parvum* and *P. mutans* were identical. My experiments leave no doubt that *P. parvum* and *mutans* are two different organisms, which view has been held for a number of years by different authors. Blood containing *P. parvum* in extraordinarily large numbers was injected subcutaneously and into the organs of various animals, and in no instance was an infection transmitted. On the contrary, it was an easy matter to transmit *P. mutans* to a healthy beast. An emulsion of 1 c.c. red corpuscles in 10 c.c. saline solution was sufficient to do so. As a further proof, it may be stated that *Rhipicephalus appendiculatus* in the Transvaal transmit *P. mutans* only in exceptional cases.

P. parvum and *P. mutans* are therefore, with regard to their pathogenic

effect, two different parasites which also show differences in their cycle of development.

Together with the daily blood examinations, systematic punctures of glands and spleen were made. On comparing preparations from cattle suffering from East Coast Fever with those infected with *P. mutans* no doubts can be left about the cause of East Coast Fever. Shortly before the parasites appear in the blood one will notice segmenting forms of Koch's bodies which throw off a great number of parasites. For this investigation the large shoulder glands are the most suitable. Neither an infection with *P. bigeminum* nor one with *P. mutans*, *P. equi*, or *P. canis* shows similar evolution forms, whereby it is proved that Koch's bodies represent a stage in the development of *P. parvum*, as already Koch him-

self had surmised. Accordingly the view of Martin Mayer, that the so-called Koch's granules are not specific for East Coast Fever, falls to the ground.

The blue bodies represent, indeed, a certain stage in the cycle of P. parvum which corresponds to the shizogony in the cycle of the malarial and other parasites.

These bodies appear both intracellularly and extracellularly, more particularly in the lymphocytes. Quite exceptionally polynuclear leucocytes are found to be infected. The development takes place more or less in the following manner. In the blue body (Agamont) the nucleus is first formed; this grows and becomes richer in chromatin and plasma; subsequently it divides successively into a great number of smaller nuclei

similarly to what occurs in the cycle of hæmoproteus which takes place in the organs of the pigeon and paddy. The cells thus formed break up into as many subdivisions as there are nuclei (Figs. 1-6). One encounters but rarely a second one, to which I will refer hereafter. The explanation of this fact may be that this shizogonous stage, which I call, with Hartmann, Agamogony, repeats itself. Accordingly agametes, resulting from the agamogonous stage, would grow into agamonts. I have, however, not been able to completely follow up the further development of these later forms.

The second mode of multiplication ends with the segmentation into a great number of small parasites, undoubtedly resulting from the agamo-

gonous multiplication (Figs. 7-13). The young parasites enter into the red corpuscles, and represent now the forms known as *P. parvum*. Thus we have in the organs, particularly in the lymphatic glands and the spleen, two different phases, which morphologically are not difficult to distinguish from each other. The nuclei of the one phase (agamogonous) when young show a loose structure, and are irregularly shaped. The nuclei of the second phase, that is, of the second generation or gamogonous, take the nuclear stains more deeply; they undergo reductions, and accordingly appear smaller. Before the segmentation of the nuclei has taken place they are oval-shaped. One is struck by the fact that the extracellular segmenting forms of the gamogonous generation produce parasites in far smaller numbers (Figs. 14-15) than those found in the cells themselves

(Figs. 12-13). I have not been able to give an explanation of this observation. In the blood no multiplication has as yet been seen, but my studies on these parasites are not yet concluded. Plasma granules were found in the blood, but this fact by no means necessitates a change of the above technology. It may also be possible that two agamogonous stages have to be distinguished, and the parasites which develop in the red corpuscles would have to be considered as the gamogenous generation. I am, however, of the opinion that the two stages in the cycle which I have called provisionally agamogony and gamogony, and which finally lead to the formation of the forms seen in the blood, succeed each other within the cell. Accordingly it is clear that the number of parasites increase as the disease develops. The forms which result from the segmentation of the last generation differ morphologically in no way from the parasites

seen in the blood at the commencement of an infection. They possess, in proportion to the whole parasite, a large nucleus, which contains a distinctive Karyosome. In using Giemsa's stain one can see differences due to the staining in the plasma of the bodies belonging to the agamogonous stage as well as of the gamogonous one. There are dark blue forms along with light blue ones containing larger alveoli. It is possible that from the very start we have to deal with a sexual differentiation. The above nuclear phenomenon can easily be studied in preparations stained according to the dry and moist methods of Giemsa, as well as in those stained with hæmatoxylin. Microscopic examination of the various organs, particularly of the spleen, does not always allow of definite conclusions. These organs represent a filter, which retains the broken-down parasites usually encountered in all protozoal diseases. When using the vital method I was never able to recognise the blue bodies, so that it is impossible for me to believe that these represent an assimilation product of the nuclei. The presence of Koch's bodies in the kidneys is of secondary importance. In East Coast Fever experimentally produced, either through the implantation of organs or in cases brought on by ticks, which finish rapidly with death, the formation of infarcts in the kidneys may not take place at all. My drawings of, and the communication referring to, Koch's granules, concern fresh material obtained with a syringe from the lymphatic glands and the spleen. In cases of East Coast Fever where fever is present without parasites being present in the blood, the diagnosis can be made by examining juice from the spleen and lymphatic glands. If in the juice obtained from the puncture of the lymphatic glands or the spleen are seen Koch's bodies containing only larger nuclei of a loose structure, and no forms of the second or gamogonous generation, it is possible that this latter does not take place at all, and no parasites are then found in the blood. Whether ticks which have been sucking the blood of such animals transmit the disease will be shown by later experiments.

It remains to explain the fact that the blood of animals suffering from East Coast Fever injected into healthy animals does not produce the disease. It is possible that the blood contains forms which can only develop in the tick, and which, injected into the animal, die. In the transplantation of organs undertaken by Theiler and Meyer, Koch's granules are directly transmitted. These represent, under natural conditions, the stages which are formed only after the tick has transmitted the disease. The infection of the various glands in the body of an animal takes place through the blood current, and accordingly it becomes feasible to expect that the inoculation of blood also ought to produce the disease. A suitable occasion would probably arise when the plasma granules are found in the blood. Experiments to elucidate this will be carried out as

soon as opportunity occurs. In an animal which was infected by the transplantation of organs I found Koch's granules in the blood. Dr. Meyer showed me blood-smears sent in from the practice in which a great number of mononuclear leucocytes contained the plasma bodies.

Concerning the place of the East Coast Fever parasite in protozoology, the proposition of Bettencourt and others to separate it from the piroplasms and to substitute a new name (*Theileria parva*) is justified.

DESCRIPTION OF THE FIGURES.

- FIG.
1-6. Agamogonous generation.
3. Intracellular form.
7-13. Intracellular gamogonous forms.
12-13. Gamogonous forms in segmentation.
14-15. Free gamogonous forms.
15. Free gamogonous form in segmentation.
16. Free parasites.
17. Parasites in the blood corpuscle.